

12. Risk Assessment (Annex III of the Cartagena Protocol on Biosafety)

COMMON FORMAT FOR Risk Assessment

(In accordance with Annex III of the Cartagena Protocol on Biosafety)

Risk assessment details	
1. Country Taking Decision:	South Africa
2. Title:	Application for Commodity Clearance of Genetically Modified Organisms (GMO) - MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7
3. Contact details:	<p>Monsanto Company, represented by Monsanto S.A.(Pty) Ltd</p> <p>Monsanto Company 800 N. Lindbergh Boulevard St. Louis, Missouri 63167 USA</p> <p>Monsanto House, Building No. 4 Fourways Office Park Maizeer Fourways Boulevard and Roos Streets Fourways South Africa</p>
LMO information	
4. Name and identity of the living modified organism:	Multi-event stack MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7
5. Unique identification of the living modified organism:	MON-87427-7× MON-89034-3 × DAS-01507-1 × MON-87411-9 × DAS-59122-7
6. Transformation event:	MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7
7. Introduced or Modified Traits:	<p>Chemical tolerance - Herbicide tolerance</p> <p>Pest resistance - Insect resistance</p>
8. Techniques used for modification:	MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 maize was obtained by conventional breeding of five single maize products: MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7. MON 87427, MON 89034 and MON 87411 have been developed by Monsanto Company while TC1507 and DAS 59122-7 were developed by Pioneer Hi-Bred International and Mycogen Seeds c/o Dow AgroSciences LLC.

NON-COMPANY CONFIDENTIAL INFORMATION

9. Description of gene modification:	<p>Conventional breeding techniques was used to develop MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7, a combined-trait maize product that has a glyphosate-induced, non-viable pollen phenotype to produce hybrid seed and confers herbicide tolerance and insect resistance. Each biotechnology-derived trait contributes specific benefits to the final combined product as follows:</p> <p>MON 87427 utilizes a specific promoter and intron combination to drive CP4 EPSPS protein expression in vegetative and female reproductive tissue. Little to no CP4 EPSPS protein is expected to be produced in MON 87427 pollen, thus providing maize lines with tissue-selective glyphosate tolerance to facilitate the production of viable hybrid maize seed.</p> <p>MON 89034 produces two insecticidal proteins that protect against feeding damage caused by European corn borer (<i>Ostrinia nubilalis</i>) and other lepidopteran insect pests. Cry1A.105 is a modified <i>Bacillus thuringiensis</i> (Bt) Cry1A protein and Cry2Ab2 is a Bt (subsp. <i>kurstaki</i>) protein. The combination of the two insecticidal proteins provides enhanced insect control and offers an additional insect-resistance management tool.</p> <p>TC1507 expresses the Cry1F insecticidal proteins, providing protection against certain lepidopteran insect pests such as European corn borer (<i>Ostrinia nubilalis</i>) and the phosphinothricin-N-acetyltransferase (PAT) protein that provides tolerance to glufosinate-ammonium.</p> <p>MON 87411 contains a suppression cassette that expresses an inverted repeat sequence designed to match the sequence of western corn rootworm (WCR; <i>Diabrotica virgifera virgifera</i>). The expression of the suppression cassette results in the formation of a double-stranded RNA (dsRNA) transcript containing a 240 bp fragment of the WCR Snf7 gene (DvSnf7). Upon consumption, the plant-produced dsRNA in MON 87411 is recognized by the CRW's RNA interference (RNAi) machinery, resulting in down-regulation of the targeted DvSnf7 gene and leading to CRW mortality. MON 87411 also contains a cry3Bb1 gene that produces a modified <i>Bacillus thuringiensis</i> (subsp. <i>kumamotoensis</i>) Cry3Bb1 protein to protect against CRW larval feeding. In addition, MON 87411 contains the cp4 epsps gene from <i>Agrobacterium</i> sp. strain CP4 that encodes for the 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) protein, which confers tolerance to glyphosate, the active ingredient in Roundup® agricultural herbicides.</p> <p>DAS 59122-7 expresses the Cry34/35Ab1 binary insecticidal proteins that also provide protection against certain coleopteran insect pests belonging to the <i>Chrysomelidae</i> family (corn rootworm, <i>Diabrotica</i> spp.) and the PAT protein that provides tolerance to glufosinate-ammonium.</p>
	Characteristics of modification
10. Vector characteristics (Annex III.9(c)):	Not applicable MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 was obtained through conventional breeding methods.

® Roundup and Roundup Ready are registered trademarks of Monsanto Technology LLC

NON-COMPANY CONFIDENTIAL INFORMATION

11. Insert or inserts (Annex III.9(d)):	<p>Like MON 87427, MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7, expresses the CP4 EPSPS protein isolated from <i>Agrobacterium</i> sp. strain CP4, which confers tolerance to glyphosate, the active ingredient in the Roundup® family of agricultural herbicides.</p> <p>Like MON 89034, MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 expresses the Cry1A.105 and Cry2Ab2 insecticidal proteins derived from <i>B. thuringiensis</i>, providing protection against certain lepidopteran insect pests such as maize stalk borers.</p> <p>Like TC1507, MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 expresses the Cry1F insecticidal proteins, providing protection against certain lepidopteran insect pests such as European corn borer (<i>Ostrinia nubilalis</i>) and the phosphinothricin-N-acetyltransferase (PAT) protein that provides tolerance to glufosinate-ammonium.</p> <p>Like MON 87411, MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 contains a suppression cassette that expresses an inverted repeat sequence designed to match the sequence of western corn rootworm (WCR; <i>Diabrotica virgifera virgifera</i>). The expression of the suppression cassette results in the formation of a double-stranded RNA (dsRNA) transcript containing a 240 bp fragment of the WCR Snf7 gene (DvSnf7). Upon consumption, the plant-produced dsRNA in MON 87411 is recognized by the CRW's RNA interference (RNAi) machinery, resulting in down-regulation of the targeted DvSnf7 gene and leading to CRW mortality. MON 87411 also contains a <i>cry3Bb1</i> gene that produces a modified <i>Bacillus thuringiensis</i> (subsp. <i>kumamotoensis</i>) Cry3Bb1 protein to protect against CRW larval feeding. In addition, MON 87411 contains the <i>cp4 epsps</i> gene from <i>Agrobacterium</i> sp. strain CP4 that encodes for the 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) protein, which confers tolerance to glyphosate, the active ingredient in Roundup® agricultural herbicides.</p> <p>Like DAS 59122-7, MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 expresses the Cry34/35Ab1 binary insecticidal proteins that also provide protection against certain coleopteran insect pests belonging to the <i>Chrysomelidae</i> family (corn rootworm, <i>Diabrotica</i> spp.) and the PAT protein that provides tolerance to glufosinate-ammonium.</p>
---	---

Recipient organism or parental organisms (Annex III.9(a)):

12. Taxonomic name/status of recipient organism or parental organisms:	<p>Common name: Maize</p> <p>Family name: Gramineae</p> <p>Genus: Zea</p> <p>Species: mays (2n+20)</p>
13. Common name of recipient organism or parental organisms:	Maize or Corn

® Roundup and Roundup Ready are registered trademarks of Monsanto Technology LLC

NON-COMPANY CONFIDENTIAL INFORMATION

14. Point of collection or acquisition of recipient or parental organisms:	MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 was produced using elite parent lines into which the MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7 events have been introgressed independently, using conventional breeding techniques. The original transformations that produced the individual events used privately owned germplasm acquired for this purpose.
15. Characteristics of recipient organism or parental organisms related to biosafety:	<p>Maize is the world's third leading cereal, following rice and wheat, in terms of production and area harvested. It has a long history of safe use as a raw material for processed products, and direct uses as a human food or animal feed. Today, maize is produced on every continent except Antarctica, and is exported and imported as viable grain for use as foods or feeds, or directly in processing, without risk to the environment.</p> <p>According to the OECD [Consensus Document on the Biology of <i>Zea mays</i> subsp. <i>mays</i> (Maize), 2003], "<i>Maize has lost the ability to survive in the wild due to its long process of domestication, and needs human intervention to disseminate its seed</i>". In addition, "<i>maize is incapable of sustained reproduction outside of domestic cultivation</i>", and "<i>maize plants are non-invasive in natural habitats</i>." Despite the fact that maize frequently appears as a volunteer plant in a subsequent rotation, it has no inherent ability to persist or propagate. In all regions of the world, volunteer plants are managed with herbicides, tillage, or manual removal of plants. As such, maize is not considered to be a pest anywhere in the world. When it occurs outside of cultivation, it has no impact on the conservation and sustainable use of biological diversity.</p> <p>Gene flow from maize occurs through dispersal of seed and pollen-mediated exchange of genes to sexually compatible plants. Since maize has no biological mechanism to scatter seed, low-level, incidental dispersal of viable grain occurs as a result of human-based activities such as transport and harvesting operations. As was noted by the OECD, the few plants that might result from incidental release will not persist or meaningfully reproduce without human intervention. Gene flow via pollen is only possible to other maize plants throughout the world except in Mexico and Guatemala where wild relatives occur. Maize reproduces sexually, is a wind-pollinated, monoecious species with separate staminate (tassels) and pistillate (silk) flowers, which encourages natural cross-pollination between maize plants. The distance that viable pollen can travel depends on prevailing wind patterns, humidity, and temperature. Generally, the pollen dissemination period lasts three to seven days. Because incidental release of maize during importation occurs at very low levels, and because maize is not competitive, pollen-mediated gene flow between local maize and rare volunteers has had no effect on the conservation and sustainable use of biological diversity.</p>

<p>16. Centre(s) of origin of recipient organism or parental organisms:</p>	<p>Maize is thought to have its origin in Mexico, from where it spread northward to Canada and southward to Argentina. Although secondary centres of origin in South America are possible, the oldest archaeological evidence of domesticated maize (5000 B.C.) was discovered in the valley of Tehuacan in Mexico (Benson and Pearce, 1987). Several theories on the origin of maize have been proposed; the two theories most adhered to being that either teosinte (a wild relative of maize that is endemic to parts of Mexico and Guatemala) or a wild pod maize that is now extinct was the wild ancestor of maize (Benson and Pearce, 1987; Brown et al., 1984).</p> <p>Maize is a member of the genus <i>Zea</i>, which is broken into 2 sections: ZEA and LUXURIANTES. The section ZEA includes one species (<i>mays</i>), which includes three subspecies: ssp. <i>mays</i>, ssp. <i>mexicana</i> (formerly <i>Euchlaena mexicana</i>), and ssp. <i>parviglumis</i>. The former subspecies is known as maize while the latter comprise a portion of the complex known as teosinte. Furthermore, ssp. <i>mexicana</i> and ssp. <i>parviglumis</i> are further separated into several races (OECD, 2003). Section LUXURIANTES encompasses 3 species: an annual <i>Z. luxurians</i>, and perennials <i>Z. diploperennis</i> and <i>Z. perennis</i>. While the classification of <i>Zea</i> continues to be modified, teosintes are the only know wild relatives of maize capable of forming hybrids in nature. Outcrossing and gene exchange between teosinte and maize has been reported with annual teosinte (<i>Zea mays</i> ssp. <i>mexicana</i>) ($2n = 20$) and maize (<i>Zea mays</i> L.) ($2n = 20$). A frequency of one F1 hybrid (maize \times teosinte) for every 500 maize plants or 20 to 50 teosinte plants in the Chalco region of the Valley of Mexico was reported. However, newer information shows that annual teosintes may be a separate species because of the level of genetic isolation and that hybrids that do form are highly unsuccessful in introgressing genetic material (OECD, 2003). Regardless, Mexico and parts of Central America are regarded as the centre of genetic diversity for maize. The natural distribution of teosinte is limited to the seasonally dry, suropical zone with summer rain along the western escarpment of Mexico and Guatemala and the Central Plateau of Mexico.</p> <p>The belief that Central America and southern Mexico are both the centre of origin and a centre of diversity for maize was supported by Vavilov (1992).</p>
<p>17. Centres of genetic diversity, if known, of recipient organism or parental organisms:</p>	<p>Refer to point 16 above</p>

NON-COMPANY CONFIDENTIAL INFORMATION

18. Habitats where the recipient organism or parental organisms may persist or proliferate:	<p>As noted by OECD (2003), maize is not invasive of natural habitats, does not persist or disperse anywhere in the world without the human intervention. Early domestication and diversification through selection occurred in Meso-America. Maize is grown across a wide range of ecological conditions including soil types, altitude and rainfall. Currently, maize is grown over a wide range of conditions because of its many divergent types that have been bred for this purpose. The bulk of the maize is produced between latitudes 30° and 55°, with relatively little grown at latitudes higher than 47° latitude anywhere in the world. The greatest maize production occurs where the warmest month isotherms range between 21°C and 27°C and the frost-free season lasts 120 to 180 days. A summer rainfall of 15 cm is approximately the lower limit for maize production without irrigation.</p> <p>Experience with maize imported for use as foods or feeds, or directly in processing, has demonstrated that stable populations do not establish, persist or proliferate because of this practice.</p>
Donor organism or organisms (Annex III.9(b)):	
19. Taxonomic name/status of donor organism(s)	<p>MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 was obtained from conventional breeding of the single events MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7.</p> <p>The donor organisms used in development of the single events MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7 were <i>Bacillus thuringiensis</i>, <i>Agrobacterium tumefaciens</i> strain CP4,.</p>
20. Common name of donor organism(s):	Bacteria
21. Point of collection or acquisition of donor organism(s):	The organisms are ubiquitous in nature
22. Characteristics of donor organism(s) related to biosafety:	Not applicable, since the donor organisms are ubiquitous in nature and therefore do not pose a threat to biodiversity.
Intended use and receiving environment	
23. Intended use of the LMO (Annex III 9(g)):	The intent of this application is for the commodity import of maize containing MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7
24. Receiving environment (Annex III.9(h)):	This is an application for Commodity Clearance therefore the product will not be cultivated in South Africa.
Risk assessment summary	

NON-COMPANY CONFIDENTIAL INFORMATION

- | | |
|--|--|
| 25. Detection/Identification method of the LMO (Annex III.9(f)): | MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 was produced by crossing plants containing MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7, using conventional breeding methods. Therefore, MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 is detectable using the combination of the single event-specific PCR method for detecting the introduced DNA present in the single events MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7. |
|--|--|

For all plants in which two or more events are combined by conventional breeding, the events involved will segregate in the grain; therefore such detection methods when applied to individual grains harvested from fields of MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 will detect any combination of the four events.

Event specific detection methods for detection of MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7 DNA have been validated by the European Commission Joint Research Centre (EU JRC) and are available on the EU JRC website

- | | |
|---|--|
| 26. Evaluation of the likelihood of adverse effects (Annex III.8(b)): | |
|---|--|

Based on the nature of the recipient species (unable to proliferate) and the lack of related and wild species with which MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS-59122-7 can outcross, the likelihood of adverse effects from outcrossing to other related species is negligible.

Transgenic maize hybrids with similar genes have been grown around the world for several years and in South Africa for 19 years without any recorded impact on the environment other than those created by conventional maize production.

Any volunteers could, like conventional maize, be removed by current agricultural practices such as ploughing and the use of herbicides.

NON-COMPANY CONFIDENTIAL INFORMATION

27. Evaluation of the consequences (Annex III.8(c)):	<p>Studies conducted with MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 confirmed that this event is agronomically and compositionally equivalent to conventional maize and has no increased tendency towards weediness or an increased susceptibility of tolerance to insects normally associated with maize. Thus, should any of the potential risks materialize, the consequences would be negligible.</p> <p>No potentially adverse effects were detected based on extensive characterization of MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7, which included molecular analysis, expression analysis, compositional analyses and phenotypic evaluation.</p> <p>Testing of MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 demonstrated no changes in its ability to persist in the environment without human intervention or to become invasive compared to conventional maize. As such, the potential consequences to biodiversity resulting from MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 for commercial use, including food, feed or processing, are the same as with conventional maize.</p> <p>Any volunteer seed germinating in subsequent growing seasons would be detected in the fields that were planted and destroyed using chemical or mechanical means.</p>
28. Overall risk (Annex III.8(d)):	Considering the potential risks and the consequences should the potential risks materialize the overall risk of importing, cultivating or field testing MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 is extremely low.
29. Recommendation (Annex III.8(e)):	No risks have been identified and therefore other than the containment parameters that might apply through the permit conditions, no additional actions need to be taken.
30. Actions to address uncertainty regarding the level of risk (Annex III.8(f)):	The potential risks for the specific product is negligible; hence no additional actions are required except compliance with the conditions contained in the permit.
Additional information	
31. Availability of detailed risk assessment information:	More information regarding the safety of MON 87427 × MON 89034 × TC1507 × MON 87411 × DAS 59122-7 is contained in the application preceding this section. Additional information can be obtained from the individual event dossiers and Annexes submitted for MON 87427, MON 89034, TC1507, MON 87411 and DAS 59122-7.
32. Any other relevant information:	None
33. Attach document:	<i>Not applicable to applicant</i>
34. Notes:	None