



US009453234B2

(12) **United States Patent**
Conner et al.

(10) **Patent No.:** **US 9,453,234 B2**
(45) **Date of Patent:** ***Sep. 27, 2016**

(54) **CHIMERIC PROMOTERS COMPRISING A RICE ACTIN 1 PROMOTER AND 35S ENHANCERS FOR USE IN PLANTS**

(71) Applicant: **Monsanto Technology LLC**, St. Louis, MO (US)

(72) Inventors: **Timothy W. Conner**, Chesterfield, MO (US); **Stanislaw Flasinski**, Chesterfield, MO (US); **Sheng Z. Pang**, Chesterfield, MO (US); **Jinsong You**, Manchester, MO (US)

(73) Assignee: **Monsanto Technology LLC**, St. Louis, MO (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 495 days.
This patent is subject to a terminal disclaimer.

(21) Appl. No.: **13/861,951**

(22) Filed: **Apr. 12, 2013**

(65) **Prior Publication Data**

US 2013/0324411 A1 Dec. 5, 2013

Related U.S. Application Data

(60) Division of application No. 12/876,071, filed on Sep. 3, 2010, now Pat. No. 8,436,161, which is a division of application No. 12/023,350, filed on Jan. 31, 2008, now Pat. No. 7,838,652, which is a continuation of application No. 11/038,981, filed on Jan. 20, 2005, now Pat. No. 7,371,848.

(60) Provisional application No. 60/537,793, filed on Jan. 20, 2004.

(51) **Int. Cl.**
C12N 15/82 (2006.01)

(52) **U.S. Cl.**
CPC **C12N 15/82** (2013.01); **C12N 15/823** (2013.01); **C12N 15/8216** (2013.01); **C12N 15/8225** (2013.01); **C12N 15/8275** (2013.01); **C12N 15/8286** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,853,530 A 12/1974 Franz
5,097,025 A * 3/1992 Benfey et al. 536/23.6
5,362,865 A 11/1994 Austin
5,530,196 A 6/1996 Fraley et al.

5,641,876 A 6/1997 McElroy et al.
6,051,753 A 4/2000 Comai et al.
6,429,357 B1 8/2002 McElroy et al.
6,448,476 B1 9/2002 Barry
6,462,258 B1 10/2002 Fincher et al.
6,489,542 B1 12/2002 Corbin et al.
6,569,122 B2 5/2003 Fischer et al.
6,660,911 B2 12/2003 Fincher et al.
6,670,467 B2 12/2003 Barbour et al.
6,689,880 B2 2/2004 Chen et al.
6,759,575 B2 7/2004 Michiels et al.
6,919,495 B2 7/2005 Fincher et al.
7,371,848 B2 5/2008 Conner et al.
7,622,632 B2 11/2009 Ursin et al.
7,838,652 B2 11/2010 Conner et al.
8,436,161 B2 * 5/2013 Conner et al. 536/24.1

FOREIGN PATENT DOCUMENTS

WO WO 00/11185 * 3/2000 C12N 15/82
WO WO 00/11185 A2 3/2000
WO WO 00/29596 * 5/2000 C12N 15/82
WO WO 00/29596 A1 5/2000
WO WO 01/44457 A2 6/2001

OTHER PUBLICATIONS

Benfey et al., "Tissue-specific expression from CaMV 35S enhancer subdomains in early stages of plant development," *EMBO J.*, 9(6):1677-1684, 1990.

Kay et al., "Duplication of CaMV 35S promoter sequences created a strong enhancer for plant genes," *Science*, 236:1299-1302, 1987.

Lam et al., "Site-specific mutations alter in vitro factor binding and change promoter expression pattern in transgenic plants," *Proc. Natl. Acad. Sci. USA*, 86:7890-7894, 1989.

Omirulleh et al., "Activity of a chimeric promoter with the double CaMV 35S enhancer element in protoplast-derived cells and transgenic plants in maize," *Plant Mol. Biol.*, 21:415-428, 1993.

* cited by examiner

Primary Examiner — Cathy Kingdon Worley

(74) *Attorney, Agent, or Firm* — Dentons US LLP; Carine M. Doyle, Esq.

(57) **ABSTRACT**

The present invention provides polynucleotide molecules useful for expressing transgenes in plants, including novel chimeric promoters made by fusing the rice actin 1 promoter and selected CaMV 35S promoter enhancer domains. The present invention also provides expression constructs containing the polynucleotide molecules useful for expressing transgenes in plants. The present invention also provides transgenic plants and seeds containing the polynucleotide molecules useful for expressing transgenes in plants.

6 Claims, 3 Drawing Sheets




Figure 1




Figure 2




Figure 3

**CHIMERIC PROMOTERS COMPRISING A
RICE ACTIN 1 PROMOTER AND 35S
ENHANCERS FOR USE IN PLANTS**

This application is a divisional of U.S. application Ser. No. 12/876,071 filed Sep. 3, 2010, now U.S. Pat. No. 8,436,161, which is a divisional of U.S. application Ser. No. 12/023,350 filed Jan. 31, 2008, now U.S. Pat. No. 7,838,652, which is a continuation of U.S. application Ser. No. 11/038,981 filed Jan. 20, 2005, now U.S. Pat. No. 7,371,848, which claims the benefit of U.S. application 60/537,793 filed Jan. 20, 2004, each of the disclosures of which are incorporated herein by reference in their entirety.

INCORPORATION OF SEQUENCE LISTING

Two copies of the sequence listing (Seq. Listing Copy 1 and Seq. Listing Copy 2) and a computer-readable form of the sequence listing, all on CD-ROMs, each containing the file named pa_01117.rpt, which is 61,440 bytes (measured in MS-DOS) and was created on Jan. 18, 2005 all of which are incorporated herein by reference.

FIELD OF THE INVENTION

The invention relates to the field of plant molecular biology and plant genetic engineering and polynucleotide molecules useful for modulating gene expression in plants.

BACKGROUND

One of the goals of plant genetic engineering is to produce plants with agronomically desirable characteristics or traits. The proper expression of a desirable transgene in a transgenic plant is one way to achieve this goal. Promoters are non-coding polynucleotide molecules which play an integral role in the overall expression of genes in living cells. Isolated promoters that function in plants are useful for modifying plant phenotypes through the methods of genetic engineering.

Many constitutive promoters are available and are useful for providing good overall gene expression. For example, constitutive promoters such as P-FMV, the promoter from the ³⁵S transcript of the Figwort mosaic virus, (U.S. Pat. No. 6,051,753); P-CaMV 35S, the promoter from the 35S RNA transcript of the Cauliflower mosaic virus, (U.S. Pat. No. 5,530,196); P-Rice Actin 1, the promoter from the actin 1 gene of *Oryza sativa*, (U.S. Pat. No. 5,641,876); and P-NOS, the promoter from the nopaline synthase gene of *Agrobacterium tumefaciens* are known to provide some level of gene expression in most or all of the tissues of a plant during most or all of the plant's lifespan. Alternately, many promoters are available with more specific expression patterns such as tissue specificity, temporal specificity, or developmental specificity. These promoters are useful for the targeted expression of a transgene in plants.

Optimal expression of a transgene is useful for producing plants with agronomically desirable characteristics or traits. Such optimal expression often requires a promoter having a specific expression pattern which may not be readily available in known promoters. One example of such a specific expression pattern is a high level of transgene expression in both vegetative and reproductive tissues. The present invention solves this problem by producing novel chimeric promoters containing elements from known promoters. These

novel chimeric promoters can then be tested in plants to determine whether the desired expression pattern is indeed achieved.

SUMMARY

In one embodiment the invention provides novel chimeric promoters provided as SEQ ID NO: 9-35 comprising a caulimovirus promoter enhancer fused with a plant actin gene promoter and useful for modulating gene expression in plants. In another embodiment the invention provides constructs comprising the novel chimeric promoter and useful for modulating gene expression in plants. In another embodiment the invention provides a transgenic plant comprising the novel chimeric promoter and the seed of the transgenic plant. In another embodiment the invention provides a method of inhibiting weed growth in a field of transgenic glyphosate tolerant crop plants comprising planting the transgenic plants transformed with an expression cassette comprising the novel chimeric promoter operably linked to a DNA molecule encoding a glyphosate tolerance gene and applying glyphosate to the field at an application rate that inhibits the growth of weeds.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 represents a section of the CaMV 35S promoter with enhancer domains marked. Also diagrammatically represented are five enhancer domains constructed for use in creating chimeric actin promoters.

FIG. 2 represents the native rice actin 1 promoter and novel chimeric promoters made by fusing the rice actin 1 promoter and selected CaMV 35S promoter enhancer domains.

FIG. 3 represents the native *Arabidopsis* actin 1 promoter and novel chimeric promoters made by fusing the *Arabidopsis* actin 1 promoter and selected CaMV 35S promoter enhancer domains.

DETAILED DESCRIPTION

The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

The invention disclosed herein provides novel combinations of polynucleotide molecules for use in constructing novel chimeric promoters. The design, construction, and use of chimeric or hybrid promoters comprising one or more of the enhancer domains of a caulimovirus 35S promoter and a plant actin gene promoter is one object of this invention. The novel chimeric promoter sequences thereof of SEQ ID NO: 9-35, are capable of transcribing operably linked DNA sequences in multiple tissues and therefore can selectively regulate expression of transgenes in multiple tissues.

As used herein, the term "polynucleotide molecule" refers to the single- or double-stranded DNA or RNA of genomic or synthetic origin, i.e., a polymer of deoxyribonucleotide or ribonucleotide bases, respectively, read from the 5' (upstream) end to the 3' (downstream) end.

As used herein, the term "polynucleotide sequence" refers to the sequence of a polynucleotide molecule. The nomenclature for DNA bases as set forth at 37 CFR §1.822 is used.

As used herein, the term "gene regulatory activity" refers to the ability to affect transcription or translation of an

operably linked transcribable polynucleotide molecule. An isolated polynucleotide molecule having gene regulatory activity may provide temporal or spatial expression or modulate levels and rates of expression of the operably linked transcribable polynucleotide molecule. An isolated

polynucleotide molecule having gene regulatory activity may comprise a promoter, intron, leader, or 3' transcription termination region.

As used herein, the term "gene expression" refers to the transcription of a DNA molecule into a transcribed RNA molecule. Gene expression may be described as related to temporal, spatial, developmental, or morphological qualities as well as quantitative or qualitative indications.

As used herein, the term "regulatory element" refers to a polynucleotide molecule that may affect the transcription or translation of an operably linked transcribable polynucleotide molecule. Regulatory elements such as promoters, leaders, introns, and transcription termination regions are non-coding polynucleotide molecules having gene regulatory activity which play an integral part in the overall expression of genes in living cells. Isolated regulatory elements that function in plants are therefore useful for modifying plant phenotypes through the methods of genetic engineering.

As used herein, the term "promoter" refers to a polynucleotide molecule that is involved in recognition and binding of RNA polymerase II and other proteins (trans-acting transcription factors) to initiate transcription. A plant promoter is a native or non-native promoter that is functional in plant cells. A promoter can be used as a 5' regulatory element for modulating expression of an operably linked transcribable polynucleotide molecule. Promoters may be defined by their temporal, spatial, or developmental expression pattern.

As used herein, the term "enhancer domain" refers to a cis-acting transcriptional regulatory element, a.k.a. cis-element, which confers an aspect of the overall control of gene expression. An enhancer domain may function to bind transcription factors, trans-acting protein factors that regulate transcription. Some enhancer domains bind more than one transcription factor, and transcription factors may interact with different affinities with more than one enhancer domain. Enhancer domains can be identified by a number of techniques, including deletion analysis, i.e., deleting one or more nucleotides from the 5' end or internal to a promoter; DNA binding protein analysis using DNase I footprinting, methylation interference, electrophoresis mobility-shift assays, in vivo genomic footprinting by ligation-mediated PCR, and other conventional assays; or by DNA sequence similarity analysis with known cis-element motifs by conventional DNA sequence comparison methods. The fine structure of an enhancer domain can be further studied by mutagenesis (or substitution) of one or more nucleotides or by other conventional methods. Enhancer domains can be obtained by chemical synthesis or by isolation from promoters that include such elements, and they can be synthesized with additional flanking nucleotides that contain useful restriction enzyme sites to facilitate subsequent manipulation. Thus, the design, construction, and use of enhancer domains according to the methods disclosed herein for modulating the expression of operably linked transcribable polynucleotide molecules are encompassed by the present invention

As used herein, the term "chimeric" refers to the product of the fusion of portions of two or more different polynucleotide molecules. As used herein, the term "chimeric promoter" refers to a promoter produced through the manipulation of known promoters or other polynucleotide

molecules. Such chimeric promoters may combine enhancer domains that can confer or modulate gene expression from one or more promoters, for example, by fusing a heterologous enhancer domain from a first promoter to a second promoter with its own partial or complete regulatory elements. The novel chimeric promoters of the present invention desirably contain at least one enhancer domain fused to a plant actin promoter. Thus, the design, construction, and use of chimeric promoters according to the methods disclosed herein for modulating the expression of operably linked transcribable polynucleotide molecules are encompassed by the present invention.

As used herein, the term "percent sequence identity" refers to the percentage of identical nucleotides in a linear polynucleotide sequence of a reference ("query") polynucleotide molecule (or its complementary strand) as compared to a test ("subject") polynucleotide molecule (or its complementary strand) when the two sequences are optimally aligned (with appropriate nucleotide insertions, deletions, or gaps totaling less than 20 percent of the reference sequence over the window of comparison). Optimal alignment of sequences for aligning a comparison window are well known to those skilled in the art and may be conducted by tools such as the local homology algorithm of Smith and Waterman, the homology alignment algorithm of Needleman and Wunsch, the search for similarity method of Pearson and Lipman, and preferably by computerized implementations of these algorithms such as GAP, BESTFIT, FASTA, and TFASTA available as part of the GCG® Wisconsin Package® (Accelrys Inc., Burlington, Mass.). An "identity fraction" for aligned segments of a test sequence and a reference sequence is the number of identical components which are shared by the two aligned sequences divided by the total number of components in the reference sequence segment, i.e., the entire reference sequence or a smaller defined part of the reference sequence. Percent sequence identity is represented as the identity fraction multiplied by 100. The comparison of one or more polynucleotide sequences may be to a full-length polynucleotide sequence or a portion thereof, or to a longer polynucleotide sequence.

As used herein, the term "substantial percent sequence identity" refers to a percent sequence identity of at least about 80% sequence identity, at least about 90% sequence identity, or even greater sequence identity, such as about 98% or about 99% sequence identity. Thus, one embodiment of the invention is a polynucleotide molecule that has at least about 80% sequence identity, at least about 90% sequence identity, or even greater sequence identity, such as about 98% or about 99% sequence identity with a polynucleotide sequence described herein. Polynucleotide molecules that are capable of regulating transcription of operably linked transcribable polynucleotide molecules and have a substantial percent sequence identity to the polynucleotide sequences of the promoters provided herein are encompassed within the scope of this invention.

Promoter Isolation and Modification Methods

Any number of methods well known to those skilled in the art can be used to isolate fragments of a promoter disclosed herein. For example, PCR (polymerase chain reaction) technology can be used to amplify flanking regions from a genomic library of a plant using publicly available sequence information. A number of methods are known to those of skill in the art to amplify unknown polynucleotide molecules adjacent to a core region of known polynucleotide sequence. Methods include but are not limited to inverse PCR (IPCR), vectorette PCR, Y-shaped PCR, and genome walking

approaches. Polynucleotide fragments can also be obtained by other techniques such as by directly synthesizing the fragment by chemical means, as is commonly practiced by using an automated oligonucleotide synthesizer. For the present invention, the polynucleotide molecules were isolated by designing PCR primers based on available sequence information.

Novel chimeric promoters can be designed or engineered by a number of methods. For example, a chimeric promoter may be produced by fusing an enhancer domain from a first promoter to a second promoter. The resultant chimeric promoter may have novel expression properties relative to the first or second promoters. Novel chimeric promoters can be constructed such that the enhancer domain from a first promoter is fused at the 5' end, at the 3' end, or at any position internal to the second promoter. The location of the enhancer domain fusion relative to the second promoter may cause the resultant chimeric promoter to have novel expression properties relative to a fusion made at a different location.

Those of skill in the art are familiar with the standard resource materials that describe specific conditions and procedures for the construction, manipulation, and isolation of macromolecules (e.g., polynucleotide molecules, plasmids, etc.), as well as the generation of recombinant organisms and the screening and isolation of polynucleotide molecules.

Constructs

As used herein, the term "construct" refers to any recombinant polynucleotide molecule such as a plasmid, cosmid, virus, autonomously replicating polynucleotide molecule, phage, or linear or circular single-stranded or double-stranded DNA or RNA polynucleotide molecule, derived from any source, capable of genomic integration or autonomous replication, comprising a polynucleotide molecule where one or more transcribable polynucleotide molecule has been operably linked.

As used herein, the term "operably linked" refers to a first polynucleotide molecule, such as a promoter, connected with a second transcribable polynucleotide molecule, such as a gene of interest, where the polynucleotide molecules are so arranged that the first polynucleotide molecule affects the function of the second polynucleotide molecule. The two polynucleotide molecules may be part of a single contiguous polynucleotide molecule and may be adjacent. For example, a promoter is operably linked to a gene of interest if the promoter regulates or mediates transcription of the gene of interest in a cell.

As used herein, the term "transcribable polynucleotide molecule" refers to any polynucleotide molecule capable of being transcribed into a RNA molecule. Methods are known for introducing constructs into a cell in such a manner that the transcribable polynucleotide molecule is transcribed into a functional mRNA molecule that is translated and therefore expressed as a protein product. Constructs may also be constructed to be capable of expressing antisense RNA molecules, in order to inhibit translation of a specific RNA molecule of interest. For the practice of the present invention, conventional compositions and methods for preparing and using constructs and host cells are well known to one skilled in the art, see for example, *Molecular Cloning: A Laboratory Manual, 3rd edition Volumes 1, 2, and 3* (2000). J. F. Sambrook, D. W. Russell, and N. Irwin, Cold Spring Harbor Laboratory Press.

Constructs of the present invention would typically contain a promoter operably linked to a transcribable polynucleotide molecule operably linked to a 3' transcription termination polynucleotide molecule. In addition, constructs may include but are not limited to additional regulatory polynucleotide molecules from the 3'-untranslated region (3'

UTR) of plant genes (e.g., a 3' UTR to increase mRNA stability of the mRNA, such as the PI-II termination region of potato or the octopine or nopaline synthase 3' termination regions). Constructs may include but are not limited to the 5' untranslated regions (5' UTR) of an mRNA polynucleotide molecule which can play an important role in translation initiation and can also be a genetic component in a plant expression construct. For example, non-translated 5' leader polynucleotide molecules derived from heat shock protein genes have been demonstrated to enhance gene expression in plants (see for example, U.S. Pat. No. 5,659,122 and U.S. Pat. No. 5,362,865, all of which are incorporated herein by reference). These additional upstream and downstream regulatory polynucleotide molecules may be derived from a source that is native or heterologous with respect to the other elements present on the construct.

Thus, one embodiment of the invention is a promoter such as provided in SEQ ID NO: 9-35, operably linked to a transcribable polynucleotide molecule so as to direct transcription of said transcribable polynucleotide molecule at a desired level or in a desired tissue or developmental pattern upon introduction of said construct into a plant cell. In one embodiment, the transcribable polynucleotide molecule comprises a protein-coding region of a gene, and the promoter provides for transcription of a functional mRNA molecule that is translated and expressed as a protein product. In another embodiment, the transcribable polynucleotide molecule comprises an antisense region of a gene, and the regulatory element affects the transcription of an antisense RNA molecule or other similar inhibitory RNA in order to inhibit expression of a specific RNA molecule of interest in a target host cell.

Exemplary transcribable polynucleotide molecules for incorporation into constructs of the present invention include, for example, polynucleotide molecules or genes from a species other than the target species or genes that originate with or are present in the same species, but are incorporated into recipient cells by genetic engineering methods rather than classical reproduction or breeding techniques. The type of polynucleotide molecule can include but is not limited to a polynucleotide molecule that is already present in the plant cell, a polynucleotide molecule from another plant, a polynucleotide molecule from a different organism, or a polynucleotide molecule generated externally, such as a polynucleotide molecule containing an antisense message of a gene, or a polynucleotide molecule encoding an artificial, synthetic, or otherwise modified version of a transgene.

The regulatory elements of the present invention can be incorporated into a construct using marker genes as described and tested in transient or stable plant analyses to provide an indication of the regulatory element's gene expression pattern in stable transgenic plants. As used herein the term "marker gene" refers to any transcribable polynucleotide molecule whose expression can be screened for or scored in some way. Marker genes for use in the practice of the present invention include, but are not limited to transcribable polynucleotide molecules encoding β -glucuronidase (GUS described in U.S. Pat. No. 5,599,670, which is incorporated herein by reference) and green fluorescent protein (GFP described in U.S. Pat. No. 5,491,084 and U.S. Pat. No. 6,146,826, both of which are incorporated herein by reference), proteins that confer antibiotic resistance, or proteins that confer herbicide tolerance. Useful antibiotic resistance markers, including those encoding proteins conferring resistance to kanamycin (nptII), hygromycin B (aph IV), streptomycin or spectinomycin (aad, spec/strep) and gentamycin (aac3 and aacC4) are known in the art. Herbicides for which transgenic plant tolerance has been demonstrated and the method of the present invention can be

applied, include but are not limited to: glyphosate, glufosinate, sulfonylureas, imidazolinones, bromoxynil, delapton, cyclohezanedione, protoporphyrinogen oxidase inhibitors, and isoxasflutole herbicides. Polynucleotide molecules encoding proteins involved in herbicide tolerance are known in the art, and include, but are not limited to a polynucleotide molecule encoding 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS described in U.S. Pat. No. 5,627,061, U.S. Pat. No. 5,633,435, U.S. Pat. No. 6,040,497 and in U.S. Pat. No. 5,094,945 for glyphosate tolerance, all of which are incorporated herein by reference); polynucleotides encoding a glyphosate oxidoreductase and a glyphosate-N-acetyl transferase (GOX described in U.S. Pat. No. 5,463,175 and GAT described in U.S. Patent publication 20030083480, both of which are incorporated herein by reference); a polynucleotide molecule encoding bromoxynil nitrilase (Bxn described in U.S. Pat. No. 4,810,648 for Bromoxynil tolerance, which is incorporated herein by reference); a polynucleotide molecule encoding phytoene desaturase (crtI) described in Misawa et al. (1993) *Plant J.* 4:833-840 and Misawa et al. (1994) *Plant J.* 6:481-489 for norflurazon tolerance; a polynucleotide molecule encoding acetohydroxyacid synthase (AHAS, aka ALS) described in Sathasivan et al. (1990) *Nucl. Acids Res.* 18:2188-2193 for tolerance to sulfonylurea herbicides; a polynucleotide molecule encoding a dicamba-degrading oxygenase enzyme (described in U.S. Patent Publications US20030135879 and US20030115626, for dicamba tolerance, all of which are incorporated herein by reference); and the bar gene described in DeBlock, et al. (1987) *EMBO J.* 6:2513-2519 for glufosinate and bialaphos tolerance. The regulatory elements of the present invention can express transcribable polynucleotide molecules that encode for phosphinothricin acetyltransferase, glyphosate resistant EPSPS, aminoglycoside phosphotransferase, hydroxyphenyl pyruvate dehydrogenase, hygromycin phosphotransferase, neomycin phosphotransferase, dalapon dehalogenase, bromoxynil resistant nitrilase, anthranilate synthase, glyphosate oxidoreductase and glyphosate-N-acetyl transferase.

Thus, in one embodiment of the invention, a polynucleotide molecule of the present invention as shown in SEQ ID NO: 9-35 is incorporated into a DNA construct such that a polynucleotide molecule of the present invention is operably linked to a transcribable polynucleotide molecule that provides for a selectable, screenable, or scorable marker. The constructs containing the regulatory elements operably linked to a marker gene may be delivered to the tissues and the tissues analyzed by the appropriate mechanism, depending on the marker. The quantitative or qualitative analyses are used as a tool to evaluate the potential expression profile of a regulatory element when operatively linked to a gene of agronomic interest in stable plants. Any marker gene can be used in a transient assay. Methods of testing for marker gene expression in transient assays are known to those of skill in the art. Transient expression of marker genes has been reported using a variety of plants, tissues, and DNA delivery systems. For example, types of transient analyses can include but are not limited to direct gene delivery via electroporation or particle bombardment of tissues in any transient plant assay using any plant species of interest. Such transient systems would include but are not limited to electroporation of protoplasts from a variety of tissue sources or particle bombardment of specific tissues of interest. The present invention encompasses the use of any transient expression system to evaluate regulatory elements operably linked to any transcribable polynucleotide molecule, including but not limited to marker genes or genes of agronomic interest. Examples of plant tissues envisioned to test in transients via an appropriate delivery system would

include but are not limited to leaf base tissues, callus, cotyledons, roots, endosperm, embryos, floral tissue, pollen, and epidermal tissue.

In one embodiment of the invention, a polynucleotide molecule as shown in SEQ ID NO: 9-35 is incorporated into a construct such that a polynucleotide molecule of the present invention is operably linked to a transcribable polynucleotide molecule that is a gene of agronomic interest. As used herein, the term "gene of agronomic interest" refers to a transcribable polynucleotide molecule that includes but is not limited to a gene that provides a desirable characteristic associated with plant morphology, physiology, growth and development, yield, nutritional enhancement, disease or pest resistance, or environmental or chemical tolerance. The expression of a gene of agronomic interest is desirable in order to confer an agronomically important trait. A gene of agronomic interest that provides a beneficial agronomic trait to crop plants may be, for example, including, but not limited to genetic elements comprising herbicide resistance (U.S. Pat. Nos. 6,803,501; 6,448,476; 6,248,876; 6,225,114; 6,107,549; 5,866,775; 5,804,425; 5,633,435; 5,463,175; and U.S. Patent Publications US20030135879 and US20030115626), increased yield (U.S. Pat. Nos. RE38,446; 6,716,474; 6,663,906; 6,476,295; 6,441,277; 6,423,828; 6,399,330; 6,372,211; 6,235,971; 6,222,098; 5,716,837), insect control (U.S. Pat. Nos. 6,809,078; 6,713,063; 6,686,452; 6,657,046; 6,645,497; 6,642,030; 6,639,054; 6,620,988; 6,593,293; 6,555,655; 6,538,109; 6,537,756; 6,521,442; 6,501,009; 6,468,523; 6,326,351; 6,313,378; 6,284,949; 6,281,016; 6,248,536; 6,242,241; 6,221,649; 6,177,615; 6,156,573; 6,153,814; 6,110,464; 6,093,695; 6,063,756; 6,063,597; 6,023,013; 5,959,091; 5,942,664; 5,942,658; 5,880,275; 5,763,245; 5,763,241), fungal disease resistance (U.S. Pat. Nos. 6,653,280; 6,573,361; 6,506,962; 6,316,407; 6,215,048; 5,516,671; 5,773,696; 6,121,436; 6,316,407; 6,506,962), virus resistance (U.S. Pat. Nos. 6,617,496; 6,608,241; 6,015,940; 6,013,864; 5,850,023; 5,304,730), nematode resistance (U.S. Pat. No. 6,228,992), bacterial disease resistance (U.S. Pat. No. 5,516,671), plant growth and development (U.S. Pat. Nos. 6,723,897; 6,518,488), starch production (U.S. Pat. Nos. 6,538,181; 6,538,179; 6,538,178; 5,750,876; 6,476,295), modified oils production (U.S. Pat. Nos. 6,444,876; 6,426,447; 6,380,462), high oil production (U.S. Pat. Nos. 6,495,739; 5,608,149; 6,483,008; 6,476,295), modified fatty acid content (U.S. Pat. Nos. 6,828,475; 6,822,141; 6,770,465; 6,706,950; 6,660,849; 6,596,538; 6,589,767; 6,537,750; 6,489,461; 6,459,018), high protein production (U.S. Pat. No. 6,380,466), fruit ripening (U.S. Pat. No. 5,512,466), enhanced animal and human nutrition (U.S. Pat. Nos. 6,723,837; 6,653,530; 6,5412,59; 5,985,605; 6,171,640), biopolymers (U.S. Pat. Nos. RE37,543; 6,228,623; 5,958,745 and U.S. Patent Publication No. US20030028917), environmental stress resistance (U.S. Pat. No. 6,072,103), pharmaceutical peptides and secretable peptides (U.S. Pat. Nos. 6,812,379; 6,774,283; 6,140,075; 6,080,560), improved processing traits (U.S. Pat. No. 6,476,295), improved digestibility (U.S. Pat. No. 6,531,648) low raffinose (U.S. Pat. No. 6,166,292), industrial enzyme production (U.S. Pat. No. 5,543,576), improved flavor (U.S. Pat. No. 6,011,199), nitrogen fixation (U.S. Pat. No. 5,229,114), hybrid seed production (U.S. Pat. No. 5,689,041), fiber production (U.S. Pat. Nos. 6,576,818; 6,271,443; 5,981,834; 5,869,720) and biofuel production (U.S. Pat. No. 5,998,700); the genetic elements, methods, and transgenes described in the patents listed above are incorporated herein by reference.

Alternatively, a transcribable polynucleotide molecule can effect the above mentioned plant characteristic or phenotypes by encoding a RNA molecule that causes the targeted inhibition of expression of an endogenous gene, for

example via antisense, inhibitory RNA (RNAi), or cosuppression-mediated mechanisms. The RNA could also be a catalytic RNA molecule (i.e., a ribozyme) engineered to cleave a desired endogenous mRNA product. Thus, any transcribable polynucleotide molecule that encodes a transcribed RNA molecule that affects a phenotype or morphology change of interest may be useful for the practice of the present invention.

The constructs of the present invention are generally double Ti plasmid border DNA constructs that have the right border (RB or AGRtu.RB) and left border (LB or AGRtu.LB) regions of the Ti plasmid isolated from *Agrobacterium tumefaciens* comprising a T-DNA, which along with transfer molecules provided by the *Agrobacterium* cells, permit the integration of the T-DNA into the genome of a plant cell. The constructs may also contain the plasmid backbone DNA segments that provide replication function and antibiotic selection in bacterial cells, for example, an *Escherichia coli* origin of replication such as ori322, a broad host range origin of replication such as oriV or oriRi, and a coding region for a selectable marker such as Spec/Strp that encodes for Tn7 aminoglycoside adenytransferase (aadA) conferring resistance to spectinomycin or streptomycin, or a gentamicin (Gm, Gent) selectable marker gene. For plant transformation, the host bacterial strain is often *Agrobacterium tumefaciens* AB1, C58, or LBA4404, however, other strains known to those skilled in the art of plant transformation can function in the present invention.

Transformed Plants and Plant Cells

As used herein, the term "transformed" refers to a cell, tissue, organ, or organism into which has been introduced a foreign polynucleotide molecule, such as a construct. The introduced polynucleotide molecule may be integrated into the genomic DNA of the recipient cell, tissue, organ, or organism such that the introduced polynucleotide molecule is inherited by subsequent progeny. A "transgenic" or "transformed" cell or organism also includes progeny of the cell or organism and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a foreign polynucleotide molecule. A plant transformation construct containing a promoter of the present invention may be introduced into plants by any plant transformation method. Methods and materials for transforming plants by introducing a plant expression construct into a plant genome in the practice of this invention can include any of the well-known and demonstrated methods including electroporation (as illustrated in U.S. Pat. No. 5,384,253; microprojectile bombardment as illustrated in U.S. Pat. No. 5,015,580; U.S. Pat. No. 5,550,318; U.S. Pat. No. 5,538,880; U.S. Pat. No. 6,160,208; U.S. Pat. No. 6,399,861; and U.S. Pat. No. 6,403,865; *Agrobacterium*-mediated transformation as illustrated in U.S. Pat. No. 5,824,877; U.S. Pat. No. 5,591,616; U.S. Pat. No. 5,981,840; and U.S. Pat. No. 6,384,301; and protoplast transformation as illustrated in U.S. Pat. No. 5,508,184, all of which are incorporated herein by reference.

Methods for specifically transforming dicots are well known to those skilled in the art. Transformation and plant regeneration using these methods have been described for a number of crops including, but not limited to, cotton (*Gossypium hirsutum*), soybean (*Glycine max*), peanut (*Arachis hypogaea*), and members of the genus *Brassica*.

Methods for transforming monocots are well known to those skilled in the art. Transformation and plant regeneration using these methods have been described for a number of crops including, but not limited to, barley (*Hordeum vulgare*); maize (*Zea mays*); oats (*Avena sativa*); orchard grass (*Dactylis glomerata*); rice (*Oryza sativa*, including indica and japonica varieties); sorghum (*Sorghum bicolor*); sugar cane (*Saccharum* sp); tall fescue (*Festuca arundina-*

cea); turfgrass species (e.g. species: *Agrostis stolonifera*, *Poa pratensis*, *Stenotaphrum secundatum*); wheat (*Triticum aestivum*), and alfalfa (*Medicago sativa*). It is apparent to those of skill in the art that a number of transformation methodologies can be used and modified for production of stable transgenic plants from any number of target crops of interest.

The transformed plants are analyzed for the presence of the genes of interest and, the expression level and/or profile conferred by the promoters of the present invention. Those of skill in the art are aware of the numerous methods available for the analysis of transformed plants. For example, methods for plant analysis include, but are not limited to Southern blots or northern blots, PCR-based approaches, biochemical analyses, phenotypic screening methods, field evaluations, and immunodiagnostic assays.

The seeds of this invention can be harvested from fertile transgenic plants and be used to grow progeny generations of transformed plants of this invention including hybrid plant lines comprising the construct of this invention and expressing a gene of agronomic interest.

The present invention also provides for parts of the plants of the present invention. Plant parts, without limitation, include seed, endosperm, ovule and pollen. In a particularly preferred embodiment of the present invention, the plant part is a seed.

Still yet another aspect of the invention is a method of inhibiting weed growth in a field of transgenic crop plants comprising first planting the transgenic plants transformed with an expression cassette comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 9-35 operably linked to a transcribable polynucleotide molecule encoding a glyphosate tolerance gene and then applying glyphosate to the field at an application rate that inhibits the growth of weeds, wherein the growth and yield of the transgenic crop plant is not substantially affected by the glyphosate application. The glyphosate application rate is the effective rate necessary to control weeds in a particular glyphosate tolerant crop; these rates may range from 8 ounces/acre to 256 ounces/acre, preferably 16 ounces/acre to 128 ounces/acre, and more preferably 32 ounces/acre to 96 ounces/acre. The glyphosate is applied at least once during the growth of the glyphosate tolerant crop and may be applied 2, 3, or 4 times during the growth of the crop or more as necessary to control weeds in the field.

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent techniques discovered by the inventors to function well in the practice of the invention. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention, therefore all matter set forth or shown in the accompanying drawings is to be interpreted as illustrative and not in a limiting sense.

EXAMPLES

Example 1

Construction of Chimeric Promoters

Novel chimeric promoters are constructed by fusing at least one enhancer domain from a Caulimovirus promoter with a plant actin gene promoter. A brief description of the sequences referred to herein is provided in Table 1 below.

TABLE 1

Brief Listing of the SEQ ID NO		
SEQ ID NO	Sequence Name	Brief Description
1	P-CaMV.35S	35S promoter sequence from the Cauliflower mosaic virus
2	4xB3	P-CaMV.35S Enhancer Domain -- four tandem copies of the B3 domain (-208 to -155) as described in U.S. Pat. No. 5,097,025
3	4xAS-1	P-CaMV.35S Enhancer Domain -- four tandem copies of the "activation sequence" (-83 to -62) as described in U.S. Pat. No. 5,097,025
4	2xB1-B2	P-CaMV.35S Enhancer Domain -- two tandem copies of the B1-B2 domain (-148 to -90) as described in U.S. Pat. No. 5,097,025
5	2xA1-B3	P-CaMV.35S Enhancer Domain -- two tandem copies of the A1-B3 domain (-208 to -46) as described in U.S. Pat. No. 5,097,025
6	2xB1-B5	P-CaMV.35S Enhancer Domain -- two tandem copies of the B1-B5 domain (-343 to -90) as described in U.S. Pat. No. 5,097,025
7	P-Os.Act1	Rice actin 1 promoter
8	P-At.Act1	<i>Arabidopsis</i> Actin 1 promoter
9	P-4xB3/P-Os.Act1-1	Chimeric promoter -- 4xB3 fused to the rice Act1 promoter at the -848 nt position
10	P-4xB3/P-Os.Act1-2	Chimeric promoter -- 4xB3 fused to the rice Act1 promoter at the -462 nt position
11	P-4xB3/P-Os.Act1-3	Chimeric promoter -- 4xB3 fused to the rice Act1 promoter at the -80 nt position
12	P-4xAS-1/P-Os.Act1-1	Chimeric promoter -- 4xAS-1 fused to the rice Act1 promoter at the -848 nt position
13	P-4xAS-1/P-Os.Act1-2	Chimeric promoter -- 4xAS-1 fused to the rice Act1 promoter at the -462 nt position
14	P-4xAS-1/P-Os.Act1-3	Chimeric promoter -- 4xAS-1 fused to the rice Act1 promoter at the -80 nt position
15	P-2xB1-B2/P-Os.Act1-1	Chimeric promoter -- 2xB1-B2 fused to the rice Act1 promoter at the -848 nt position
16	P-2xB1-B2/P-Os.Act1-2	Chimeric promoter -- 2xB1-B2 fused to the rice Act1 promoter at the -462 nt position
17	P-2xB1-B2/P-Os.Act1-3	Chimeric promoter -- 2xB1-B2 fused to the rice Act1 promoter at the -80 nt position
18	P-2xA1-B3/P-Os.Act1-1	Chimeric promoter -- 2xA1-B3 fused to the rice Act1 promoter at the -848 nt position
19	P-2xA1-B3/P-Os.Act1-2	Chimeric promoter -- 2xA1-B3 fused to the rice Act1 promoter at the -462 nt position
20	P-2xA1-B3/P-Os.Act1-3	Chimeric promoter -- 2xA1-B3 fused to the rice Act1 promoter at the -80 nt position
21	P-2xB1-B5/P-Os.Act1-1	Chimeric promoter -- 2xB1-B5 fused to the rice Act1 promoter at the -848 nt position
22	P-2xB1-B5/P-Os.Act1-2	Chimeric promoter -- 2xB1-B5 fused to the rice Act1 promoter at the -462 nt position
23	P-2xB1-B5/P-Os.Act1-3	Chimeric promoter -- 2xB1-B5 fused to the rice Act1 promoter at the -80 nt position
24	P-2xA1-B3/At.Act1/ArvII	Chimeric promoter -- 2xA1-B3 fused to the <i>Arabidopsis</i> Act1 promoter at the ArvII position
25	P-2xA1-B3/At.Act1/BstZI	Chimeric promoter -- 2xA1-B3 fused to the <i>Arabidopsis</i> Act1 promoter at the BstZI position
26	P-2xA1-B3/At.Act1/BstZI-R	Chimeric promoter -- 2xA1-B3 fused to the <i>Arabidopsis</i> Act1 promoter at the BstZI position in reverse orientation
27	P-2xA1-B3/At.Act1/NsiI	Chimeric promoter -- 2xA1-B3 fused to the <i>Arabidopsis</i> Act1 promoter at the NsiI position
28	P-2xA1-B3/At.Act1/NsiI-R	Chimeric promoter -- 2xA1-B3 fused to the <i>Arabidopsis</i> Act1 promoter at the NsiI position in reverse orientation
29	P-2xA1-B3/At.Act1/BsmFI	Chimeric promoter -- 2xA1-B3 fused to the <i>Arabidopsis</i> Act1 promoter at the BsmFI position
30	P-4xAS-1/At.Act1/ArvII	Chimeric promoter -- 4xAS-1 fused to the <i>Arabidopsis</i> Act1 promoter at the ArvII position
31	P-4xAS-1/At.Act1/ArvII-R	Chimeric promoter -- 4xAS-1 fused to the <i>Arabidopsis</i> Act1 promoter at the ArvII position in reverse orientation
32	P-4xAS-1/At.Act1/BstZI	Chimeric promoter -- 4xAS-1 fused to the <i>Arabidopsis</i> Act1 promoter at the BstZI position
33	P-4xAS-1/At.Act1/BstZI-R	Chimeric promoter -- 4xAS-1 fused to the <i>Arabidopsis</i> Act1 promoter at the BstZI position in reverse orientation
34	P-4xAS-1/At.Act1/NsiI	Chimeric promoter -- 4xAS-1 fused to the <i>Arabidopsis</i> Act1 promoter at the NsiI position
35	P-4xAS-1/At.Act1/BsmFI	Chimeric promoter -- 4xAS-1 fused to the <i>Arabidopsis</i> Act1 promoter at the BsmFI position

The Caulimovirus promoter can be any promoter from a virus in the Caulimovirus family, including but not limited to promoters from Cauliflower mosaic virus (CaMV) such as the promoter from the 35S RNA transcript of the Cauliflower mosaic virus (P-CaMV.35S) (SEQ ID NO: 1) (U.S.

Pat. No. 5,530,196) and promoters from Figwort mosaic virus (FMV) such as the promoter from the 35S transcript of the Figwort mosaic virus, (U.S. Pat. No. 6,051,753), all of which are incorporated herein by reference. Promoter enhancer domains constructed as multimers of enhancer

domains from the P-CaMV.35S promoter include but are not limited to the 4x B3 domain (SEQ ID NO: 2), 4x AS-1 (SEQ ID NO: 3), 2x B1-B2 domain (SEQ ID NO: 4), 2x A1-B3 domain (SEQ ID NO: 5), and 2x B1-B5 domain (SEQ ID NO: 6). See FIG. 1. The monomers comprising the multi-

mers listed above as well as other enhancer domains from P-CaMV.35S are described in U.S. Pat. No. 5,097,025, incorporated herein by reference.

The plant actin gene promoter can be any promoter from a plant actin gene, including but not limited to actin promoters from *Oryza sativa* such as the rice actin 1 promoter (P-Os.Act1) (SEQ ID NO: 7) (U.S. Pat. No. 5,641,876), actin promoters from *Arabidopsis thaliana* such as the *Arabidopsis* Actin 1 promoter (P-At.Act1) (SEQ ID NO: 8), and actin promoters from *Zea mays* such as the Actin-2 promoter (U.S. Pat. No. 6,670,467), all of which are incorporated herein by reference.

The fusion of at least one enhancer domain with a plant actin gene promoter may be to any region of the plant actin gene promoter including but not limited to the 5' end of the plant actin gene promoter, the 3' end of the plant actin gene promoter, or any region internal to the plant actin gene promoter. The enhancer domain may be in either the reverse or the forward orientation.

Enhancer domains derived from P-CaMV.35S were inserted in three locations in the rice actin 1 (relative to rice Actin 1 transcription initiation site): -848 nt (HindIII site), -462 nt (EcoRI site), and -80 nt (FseI site). See FIG. 2. Construction of these chimeric promoters is described in detail below.

The P-4x B3/P-Os.Act1 chimeric promoters were created by fusing the four tandem copies CaMV 35S B3 enhancer sequence to the rice Act1 promoter (P-Os.Act1). Chimeras were made by fusing the enhancer region at the -848 nt position of P-Os.Act1 (SEQ ID NO: 9), at the -462 nt position of P-Os.Act1 (SEQ ID NO: 10) and at the -80 nt position of P-Os.Act1 (SEQ ID NO: 11).

The P-4x AS-1/P-Os.Act1 chimeric promoter was created by fusing four tandem copies of the CaMV 35S AS-1 enhancer sequence to the rice Act1 promoter (P-Os.Act1). Chimeras were made by fusing the enhancer region at the -848 nt position of P-Os.Act1 (SEQ ID NO: 12), at the -462 nt position of P-Os.Act1 (SEQ ID NO: 13) and at the -80 nt position of P-Os.Act1 (SEQ ID NO: 14).

The P-2xB1-B2/P-Os.Act1 chimeric promoter was created by fusing two tandem copies of the CaMV 35S B1-B2 enhancer sequence to the rice Act1 promoter (P-Os.Act1). Chimeras were made by fusing the enhancer region at the -848 nt position of P-Os.Act1 (SEQ ID NO: 15), at the -462 nt position of P-Os.Act1 (SEQ ID NO: 16) and at the -80 nt position of P-Os.Act1 (SEQ ID NO: 17).

The P-2xA1-B3/P-Os.Act1 chimeric promoters were created by fusing two tandem copies of the CaMV 35S A1-B3 enhancer sequence to the rice Act1 promoter (P-Os.Act1). Chimeras were made by fusing the enhancer region at the -848 nt position of P-Os.Act1 (SEQ ID NO: 18), at the -462 nt position of P-Os.Act1 (SEQ ID NO: 19), and at the -80 nt position of P-Os.Act1 (SEQ ID NO: 20).

The P-2xB1-B5/P-Os.Act1 chimeric promoter was created by fusing two tandem copies of the CaMV 35S B1-B5 enhancer sequence to the rice Act1 promoter (P-Os.Act1). Chimeras were made by fusing the enhancer region at the -848 nt position of P-Os.Act1 (SEQ ID NO: 21), at the -462 nt position of P-Os.Act1 (SEQ ID NO: 22) and at the -80 nt position of P-Os.Act1 (SEQ ID NO: 23).

Enhancer domains derived from P-CaMV.35S were inserted in four locations in the *Arabidopsis* actin 1 promoter

(P-At.Act1) (with the P-At.Act1 transcription initiation site designated as +1). Four unique restriction sites in P-At.Act1 were used as insertion sites: AvrII, BstZI, NsiI, BsmFI. Construction of these chimeric promoters is described in detail below.

The 2xA1B3 and 4xAS-1 fragments were isolated by restriction enzyme digest. Fragments ends were then made blunt by treatment with T4 DNA polymerase. Vectors containing the P-At.Act1 promoter were cut with one of the four unique enzymes and then the fragment ends were made blunt. Each of the two enhancer domains was fused into each of the four blunted restriction enzyme sites. Chimeric promoters were selected with inserts in both the reverse and forward orientations. See FIG. 3.

Selected chimeric promoters were subcloned into a binary vector to operably link the chimeric promoter with a reporter gene such as the GUS reporter gene (β -glucuronidase gene) or the CP4 gene (for glyphosate tolerance). These vectors were used for plant transformation and subsequent promoter characterization. Transgene expression levels of the chimeric promoters were compared with the transgene expression levels of control constructs.

Example 2

Promoter Characterization in Transient Systems

Selected chimeric promoters were used for transient transformation for reporter expression analysis. Transient systems used included tobacco protoplasts, corn protoplasts, wheat ovary, wheat anther, and barley microspores. Cells were harvested and protein extracted for GUS activity analysis. Methods for measuring GUS activity are well known to those skilled in the art, see for instance *Using the Gus Gene as a Reporter of Gene Expression* (1992) edited by Sean R. Gallagher, Academic Press, Inc., San Diego. Monocot Analysis

Constructs containing chimeric P-Os.Act1 promoters were used to transform corn protoplasts and assay for GUS/LUX activity relative to the P-Os.Act1 promoter and the P-CaMV.e35S promoter. Data are provided in Table 2.

TABLE 2

Transient Analysis in Corn Protoplasts			
Chimeric Promoter	SEQ ID NO	Construct	Relative activity
P-Os.Act1	7	pMON25455	1.0
P-2xB1-B5/P-Os.Act1-1	21	pMON38303	1.6
P-2xB1-B5/P-Os.Act1-2	22	pMON38304	1.1
P-2xB1-B5/P-Os.Act1-3	23	pMON38305	3.6
P-2xA1-B3/P-Os.Act1-1	18	pMON38310	4.5
P-2xA1-B3/P-Os.Act1-2	19	pMON38312	4.6
P-2xA1-B3/P-Os.Act1-3	20	pMON38314	11.0
P-2xB1-B2/P-Os.Act1-1	15	pMON38309	1.1
P-2xB1-B2/P-Os.Act1-2	16	pMON38311	1.5
P-2xB1-B2/P-Os.Act1-3	17	pMON38313	2.2
P-4xAS-1/P-Os.Act1-1	12	pMON38300	1.8
P-4xAS-1/P-Os.Act1-2	13	pMON38301	1.3
P-4xAS-1/P-Os.Act1-3	14	pMON38302	8.9
P-4xB3/P-Os.Act1-1	9	pMON38306	1.8
P-4xB3/P-Os.Act1-2	10	pMON38307	3.3
P-4xB3/P-Os.Act1-3	11	pMON38308	5.4
P-CaMV.e35S	1	pMON25456	5.1

All the chimeric promoters tested were found to have higher activity than the native P-Os.Act1 promoter when tested in corn protoplasts. The three chimeric promoters P-2xA1-B3/P-Os.Act1-3 (SEQ ID NO: 20), P-4xAS-1/P-Os.Act1-3 (SEQ ID NO: 14), and P-4xB3/P-Os.Act1-3

15

(SEQ ID NO: 11) also showed increased activity when compared to the CaMV e35S promoter. Two factors contributed the enhanced activity in the chimeric promoter, namely the enhancer domain selected and the fusion location of the enhancer domain. Every enhancer domain tested appeared to be more powerful when the fusion location was closer to the 3' end of the promoter, and attenuated when the fusion location was at the 5' end of the promoter. Among the enhancer domains tested, the 2xAS1-B3 enhancer was found to be the strongest enhancer domain in corn protoplasts. The 4xAS-1 enhancer was found to convey desirable strength to the chimeric promoter. The 4xB3 enhancer was found to be most effective when fused closer to the transcription start site.

One limitation of expression analysis in the corn protoplast system is that it only represents vegetative tissue. An increased expression level in the corn protoplast system is not necessarily indicative of performance in reproductive tissue. A few selected chimeric promoters were therefore further tested for GUS activity (pmol/min in 10 ul) in wheat anther, wheat ovary, barley microspore, and corn pollen transient assay systems for comparison with corn leaf protoplast data. Data are provided in Table 3 below. In comparison to P-Os.Act1 promoter which is known to express well in reproductive tissue, these chimeric promoters did not show any decrease of activity in reproductive tissue.

TABLE 3

Transient Analysis of Selected Chimeric Promoters in Reproductive Tissues						
Promoter	SEQ ID NO	Construct	Wheat Anther	Wheat Ovary	Barley Microspore	Corn Pollen
P-Os.Act1	7	pMON25455	1.0	1.0	1.0	1.0
P-CaMV.E35S	1	pMON25456	3.3	0.5	0.8	0.03
P-4xAS-1/P-Os.Act1-3	14	pMON38302	16.4	1.2	2.0	2.2
P-4xB3/P-Os.Act1-2	10	pMON38307	1.5	2.1	—	6.0
P-4xB3/P-Os.Act1-3	11	pMON38308	6.4	2.5	—	0.6
P-2xAS1-B3/P-Os.Act1-1	18	pMON38310	3.3	1.3	2.0	1.4
P-2xAS1-B3/P-Os.Act1-2	19	pMON38312	7.6	3.3	2.4	0.5
P-2xAS1-B3/P-Os.Act1-3	20	pMON38314	13.2	3.9	7.8	0.05

Dicot Analysis

Constructs containing chimeric P-At.Act1 promoters were used to transform tobacco protoplasts and assay for GUS activity (nM MUG/ μ g total protein). Data are provided in Table 4 below.

TABLE 4

Transient Analysis in Tobacco Protoplasts			
Chimeric Promoter	SEQ ID NO	Construct	GUS Activity
P-At.Act1	8	pMON54945	10
P-2xAS1-B3/At.Act1/ArvII	24	pMON59394	18
P-2xAS1-B3/At.Act1/BstZI	25	pMON59392	20
P-2xAS1-B3/At.Act1/BstZI-R	26	pMON59392-R	16
P-2xAS1-B3/At.Act1/NsiI	27	pMON59386	23
P-2xAS1-B3/At.Act1/NsiI-R	28	pMON59386-R	12
P-2xAS1-B3/At.Act1/BsmFI	29	pMON59384	65
P-4xAS-1/At.Act1/ArvII	30	pMON59393	19
P-4xAS-1/At.Act1/ArvII-R	31	pMON59393-R	15
P-4xAS-1/At.Act1/BstZI	32	pMON59391	16
P-4xAS-1/At.Act1/BstZI-R	33	pMON59391-R	21
P-4xAS-1/At.Act1/NsiI	34	pMON59385	150
P-4xAS-1/At.Act1/BsmFI	35	pMON59383	179
P-e35S	1	pMON26180	365

16

Both the fusion location and choice of enhancer domain was found to produce a significant effect on promoter activity in tobacco protoplasts. The two promoters with the highest GUS activity were the P-4xAS-1/At.Act1/NsiI (SEQ ID NO: 34) and P-4xAS-1/At.Act1/BsmFI (SEQ ID NO: 35). For constructs comprising the 2xAS1B3 enhancer, the highest activity of GUS was shown with the enhancer fused at the BsmFI position. The 2xAS1-B3 chimeric promoters were generally less active in all positions when compared to the 4xAS-1 chimeric promoters. Reverse orientation of the enhancer domain did not change significantly the activity of the chimeric promoter.

Example 3

Characterization of Chimeric Promoters in Transgenic Corn Plants

Selected chimeric promoters were used for stable corn plant transformation for reporter expression analysis. Plants were transformed using *agrobacterium*-mediated methods.

In order to have a direct side-by-side comparison of GUS activity in the cytoplasm and CP4 expression in plastids, and to minimize variations in sampling and environmental factors, a few selected chimeric promoter constructs were built.

Each test construct comprised the test promoter driving the GUS reporter gene (beta-glucuronidase coding sequence from *E. coli*) and the test promoter driving the CP4 gene (bacterial strain CP4 *aroA* gene encoding class II EPSPS enzyme) in a linear array. Four test constructs, pMON46172 (P-2xAS1-B3/P-Os.Act1-3, SEQ ID NO: 20); pMON46173 (P-4xAS-1/P-Os.Act1-3, SEQ ID NO: 14); pMON46174 (P-4xB3/P-Os.Act1-2, SEQ ID NO: 10); and pMON46175 (P-2xAS1-B3/P-Os.Act1-1, SEQ ID NO: 18), and the reference construct, pMON46170 (P-CaMV.E35S driving GUS and P-Os.Act1 driving CP4), were used to transform corn. Transformed corn plants were selected in glyphosate containing medium. Three R0 plants were generated for each transformation event. The first plant was sprayed with the equivalent of 64 ounce/acre of Roundup® Ultra, the second plant was sprayed with the equivalent of 96 ounce/acre of Roundup® Ultra. The third plant from each event was left as unsprayed control. Ten to fourteen days after Roundup® application each plant was rated for % chlorosis and % malformation. At mid-pollen shed, each R0 plant was rated for male fertility. Positive R0 plants were pollinated with LH198 pollen grains to produce F1 seed.

Transgenic events with single or lowest copy numbers of cassette based on CP4 copy number estimation by Taqman

were selected for F1 corn analysis. A total of 5 events from each construct with triplicate plants in each data point were used. The plants were grown in greenhouse from F1 seeds, and selected for positive segregates via glyphosate spray at 16 oz/acre when plants reach V-2 stage. Leaf tissue was taken at V-4 stage, and V-4 again at V-8/V-9 stage. Also at V-8/V-9 stage, tissue was collected from V-8 leaf, root tip, and immature tassel ranging from 0.5 cm-3 cm. Pollen was collected when shedding. Embryo and endosperm were harvested at 12 days after pollination. Several positive embryos were pooled from GUS positive individual kernels. Samples were extracted and used for both GUS quantitative analysis and CP4 ELISA. The P-2x \times A1-B3/P-Os.Act1-3 and P-4x \times AS-1/P-Os.Act1-3 chimeric promoters provided GUS gene expression which was as good as or better than that provided by P-CaMV.E35S in most of the tissues analyzed. The P-4x \times B3/P-Os.Act1-2 and P-2x \times A1-B3/P-Os.Act1-1 chimeric promoters provided high levels of expression in pollen with moderate or low levels of expression in other tissues as compared to that provided by P-CaMV.E35S. The Data are provided for GUS activity (pmole/min/mg protein) as mean and standard error measurements in Table 5 below.

TABLE 5

F1 Transgenic Corn GUS Activity (pmole/min/mg protein)										
Tissue/Stage	pmon46170 P- CaMV.E35S		pmon46172 P-2x \times A1-B3/ P-Os.Act1-3		pmon46173 P-4x \times AS-1/P- Os.Act1-3		pmon46174 P-4x \times B3/P- Os.Act1-2		pmon46175 P-2x \times A1-B3/ P-Os.Act1-1	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Young V4	6.54	0.29	12.28	3.00	5.11	0.75	1.02	0.35	5.70	0.60
Aged V4	2.82	0.85	6.47	1.43	6.78	1.59	0.10	0.02	4.14	1.09
V8	0.67	0.20	1.27	0.51	0.64	0.12	0.17	0.05	0.13	0.01
Tassel	0.57	0.19	0.49	0.25	0.50	0.23	0.19	0.07	0.13	0.07
Root Tip	0.38	0.04	0.84	0.10	1.11	0.43	0.19	0.04	0.97	0.38
Pollen	1.44	0.11	2.27	0.26	2.87	0.52	15.26	2.84	13.76	2.05
Embryo	2.19	0.22	2.35	0.68	5.68	1.64	1.42	0.18	2.53	0.65
Endosperm	1.73	0.15	9.21	4.35	6.18	1.59	1.40	0.51	2.09	0.37

All four chimeric promoters provided CP4 gene expression which was as good as or better than that provided by P-Os.Act1 in all of the tissues analyzed with the exception of pollen. Expression levels in pollen for chimeric promoter constructs were approximately 20% to 73% that of expression levels in pollen for P-Os.Act1 constructs. Data are provided for CP4 expression levels (μ g CP4 protein/g total protein) as mean and standard error measurements in Table 6 below.

TABLE 6

F1 Transgenic Corn CP4 Expression levels (μ g CP4 protein/g total protein)										
Tissue/Stage	pmon46170 and P-Os.Act1		pmon46172 P-2x \times A1-B3/ P-Os.Act1-3		pmon46173 P-4x \times AS-1/P- Os.Act1-3		pmon46174 P-4x \times B3/P- Os.Act1-2		pmon46175 P-2x \times A1-B3/ P-Os.Act1-1	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Young V4	31	3	35	2	408	122	280	66	952	540
Aged V4	20	2	2187	656	1802	411	524	257	688	298
V8	17	2	1055	298	870	272	311	35	284	110
Tassel	225	27	1334	515	1002	178	408	31	1072	303
Root Tip	89	8	745	185	765	173	184	22	279	68
Pollen	797	97	165	13	217	33	370	16	582	74
Embryo	241	54	623	69	1101	187	305	69	566	84
Endosperm	229	21	792	212	889	229	237	36	375	46

The F1 progenies were also used for field tests. Three F1 populations derived from three R0 plants of each event were tested. Commercial Roundup Ready[®] corn lines GA21 and NK603 were used as positive controls. Three Roundup[®] rates were used in this test: 0, 96, and 128 oz/A. Roundup[®] was applied at V4 leaf stage. Data obtained were % transformation efficiency (TE), number of single copy events per total events generated, Leaf CP4 levels (μ gCP4/g fresh weight tissue), Pollen CP4 levels (μ gCP4/g fresh weight tissue), percent chlorosis (CHL), percent malformation (MAL) at 10-14 DAT (days after treatment), and male fertility score (MFR) measured 1-5 with 5 the highest. Data represent the average score collected across the events generated for each construct and only tasseled plants were included for some measurements. Plants transformed with the P-4x \times B3/P-Os.Act1-2 or the P-4x \times AS-1/P-Os.Act1-3 chimeric promoter constructs had chlorosis scores equivalent to the reference construct (pMON46170). P-4x \times AS-1/P-Os.Act1-3 had a malformation score lower than that of the reference construct. All chimeric promoters tested had male fertility rates comparable to that of plants transformed with the reference construct. Data are provided in Table 7 below.

TABLE 7

Roundup ® Tolerance in Transgenic Corn Field Tests										
Promoter	Construct	% TE	Single Copy	Leaf CP4	Pollen CP4	Pollen GUS	% CHL	% MAL	MFR	
P-CaMV.E35S and P-Os.Act1	pMON46170	12	7/15	22	1471	4.4	1	1	5	
P-2x A1-B3/P-Os.Act1-3	pMON46172	10	5/15	116	313	5.8	7	4	5	
P-4xAS-1/P-Os.Act1-3	pMON46173	16	9/15	319	475	9.7	1	0	5	
P-4x B3/P-Os.Act1-2	pMON46174	24	8/15	78	674	24.3	1	3	5	
P-2x A1-B3/P-Os.Act1-1	pMON46175	—	1/15	77	767	34.1	2	3	5	

Example 4

Characterization of Chimeric Promoters in Transgenic Dicots

Several constructs were evaluated in transgenic *Arabidopsis* plants for GUS expression. GUS expression in leaf and flower tissue was measured as pM MUG/µg total protein and results were averaged for all the events produced for construct. The CaMV e35S promoter (P-CaMV.e35S) and *Arabidopsis* actin 1 promoter (P-At.Act1) were used as controls. Data are provided in Table 8 below

TABLE 8

Transgenic <i>Arabidopsis</i> analysis with chimeric At-Act1 promoters				
Promoter	SEQ ID		Flower	Leaf
	NO	Construct		
P-At.Act1	8	pMON59382	139	62
P-CaMV.e35S	1	pMON59381	374	176
P-2x A1-B3/At.Act1/BstZI	25	pMON59378	0	24
P-4xAS-1/At.Act1/BstZI	32	pMON59377	12	47
P-4xAS-1/At.Act1/ArvII	30	pMON59379	58	18
P-4xAS-1/At.Act1/NsiI	34	pMON59375	46	36
P-4xAS-1/At.Act1/BstZI-R	33	pMON59371	66	21
P-2x A1-B3/At.Act1/NsiI	27	pMON59376	113	56
P-2x A1-B3/At.Act1/BstZI-R	26	pMON59372	149	114
P-4xAS-1/At.Act1/BsmFI	35	pMON59373	29	256
P-2x A1-B3/At.Act1/ArvII	24	pMON59380	88	199
P-2x A1-B3/At.Act1/BsmFI	29	pMON59374	153	193

GUS expression analysis in *Arabidopsis* showed that P-2x A1-B3/At.Act1/BstZI-R (SEQ ID NO: 26) and P-2x A1-B3/At.Act1/BsmFI (SEQ ID NO: 29) provided transgene expression in flowers comparable to that of the P-At.Act1 promoter. The P-2x A1-B3/At.Act1/BstZI-R (SEQ ID NO: 26), P-4xAS-1/At.Act1/BsmFI (SEQ ID NO: 35), P-2x A1-B3/At.Act1/ArvII (SEQ ID NO: 24), and P-2x A1-B3/At.Act1/BsmFI (SEQ ID NO: 29) promoters provided transgene expression in leaves higher than or comparable to the P-At.Act1 and P-CaMV.e35S.

Example 4

Insect Control Analysis in Corn

Two constructs (pMON38858 and pMON38859) were used to generate transgenic corn plants. Transformations were performed using *Agrobacterium* mediated methods. Both constructs contained the 2x A1-B3/P-OsAct1-3 promoter (SEQ ID NO: 20) operably linked to a nucleotide sequence encoding a Cry2Ab insecticidal protein (U.S. Pat. No. 6,489,542). Several transgenic corn plant lines produced from each construct were analyzed for Cry2Ab protein levels. The 2x A1-B3/P-OsAct1-3 promoter was found to express the Cry2Ab protein at high levels in corn leaf tissue when compared with standard controls.

Example 5

Glyphosate Tolerance Analysis in Wheat

Three chimeric promoters were tested in transgenic wheat plants. Transgenic wheat plants were generated from each of the single cassette constructs pMON43646 (P-4xAS-1/P-Os.Act1-3, SEQ ID NO: 14), pMON43647 (P-2x A1-B3/P-Os.Act1-2, SEQ ID NO: 19), and pMON43648 (P-2x A1-B3/P-Os.Act1-3, SEQ ID NO: 20). Transformations were performed with *Agrobacterium* in immature Bobwhite embryos. All three constructs contain a single copy of the CP4 EPSPS gene for glyphosate tolerance. The distinguishing element in each construct is the promoter. Events from each single cassette construct were analyzed for vegetative and reproductive tolerance to glyphosate equivalent to the double cassette lead event 33391 (U.S. Patent Publication US20020062503) generated from pMON30139 which contains two copies of the CP4 EPSPS gene driven by the P-e35S and P-Os.Act1 promoters, respectively. Plants were analyzed for glyphosate tolerance, phenotype, copy number, molecular profile, and genome location. Results are provided below.

R0 plants were spray tested for vegetative and reproductive tolerance with 64 oz/A Roundup® Ultra (1.68 kg/ha acid equivalents of glyphosate) prior to jointing. Plants with vegetative damage were discarded. Fertility was estimated by counting the number of seeds in 20 florets from the central portion of the head and reported as % fertility. Data are provided in Table 9 below.

TABLE 9

Glyphosate Tolerance in Transgenic R0 Wheat Plants				
Promoter	Construct	R0 Events	# Events with	# Events with >=80%
			Vegetative Tolerance (% of total)	Fertility (% with Veg Tol)
P-Os.Act1	pMON30167	63	4 (6%)	2 (50%)
P-CaMV.e35S and P-Os.Act1	pMON30139	150	104 (69%)	24 (23%)
P-4xAS-1/P-Os.Act1-3	pMON43646	87	65 (75%)	13 (20%)

TABLE 9-continued

Glyphosate Tolerance in Transgenic R0 Wheat Plants				
Promoter	Construct	R0 Events	# Events with Vegetative Tolerance (% of total)	# Events with >=80% Fertility (% with Veg Tol)
P-2xA1-B3/P-Os.Act1-2	pMON43647	83	61 (73%)	16 (26%)
P-2xA1-B3/P-Os.Act1-3	pMON43648	69	51 (74%)	12 (24%)

Data on the vegetative and reproductive glyphosate tolerance of R0 wheat plants transformed with a single cassette vector containing CP4 driven by the rice actin promoter (pMON30167) and a double cassette vector containing two CP4 genes driven by P-CaMV.e35s and P-Os.Act1 are also provided in Table 15. Wheat transformation experiments that used the P-CaMV.e35S cassette alone resulted in plants that were vegetatively tolerant but had low fertility. For example plants generated with pMON42411, comprising only the P-CaMV.e35S promoter driving the CP4 gene, produced only 1 of 37 vegetatively tolerant plants with a fertility of >=80%. This data indicates that the e35S promoter is responsible for vegetative expression of CP4 while the rice actin promoter is responsible for reproductive expression of CP4. Results for the single cassette vector containing the three chimeric promoters (pMON43646, pMON43647, and pMON43648) were similar to those for the double cassette vector pMON30139. 73-75% of the total events generated using these three chimeric promoter constructs had vegetative tolerance, and 20-26% of those events had >=80% fertility. Thus, using one of the three chimeric promoters to drive a single copy of the CP4 EPSPS gene produced similar results in R0 tests as double cassette lead event.

In addition to glyphosate tolerance, transformation efficiencies (TE), leaf CP4 levels (µgCP4/mg total protein from R0 wheat plants), and meristem CP4 levels (µgCP4/mg total protein from R0 wheat plants) were measured in R0 plants. Plants transformed with pMON30139 (the double cassette construct) and pMON30159 (P-ScBV see U.S. Pat. No. 6,489,462) were used as controls. Data are provided in Table 10 below.

TABLE 10

CP4 Expression in Transgenic R0 Wheat Plants				
Promoter	Construct	TE	Leaf CP4	Meristem CP4
P-CaMV.e35S and P-Os.Act1	pMON30139	4%	2.23 (0.93%)	4.54 (1.48%)
P-ScBV	pMON30159	4.50%	6.99 (1.43%)	9.7 (1.71%)
P-4xAS-1/P-Os.Act1-3	pMON43646	3.30%	2.13 (.99%)	3.72 (1.55%)
P-2xA1-B3/P-Os.Act1-2	pMON43647	2.90%	1.41 (1%)	1.66 (1.23%)
P-2xA1-B3/P-Os.Act1-3	pMON43648	2.20%	4.62 (0.77%)	5.79 (1.31%)

R1 seeds were then collected from the R0 plants and advanced to R1 testing. R1 plants were simultaneously tested for glyphosate tolerance and copy number. The transgene copy number for each event was determined by Southern blot and/or TaqMan quantitative PCR analysis. CP4 gene

sequences were used as probes for both assays. A good correlation was observed between single copy calls on Southern and by TaqMan analysis. Single copy events, with good tolerance to high doses of glyphosate, were advanced to the R2 generation.

Seeds from selected wheat lines advanced to the R2 generation were planted in 2" pots and sprayed with 128 oz/A Roundup at the 3 leaf stage. Retained lines were transferred to larger pots and sprayed again with 128 oz/A at the 6 leaf stage. In an attempt to force differentiation among events an extreme pressure test was devised where plants were sprayed with 512 oz/A of Roundup®. This was repeated four times between the 3 leaf stage and emergence of the flag leaf. Results for the chimeric promoter lines were compared with results for non-transgenic Bobwhite plants, R4 generation lead event plants (line 33391 generated from pMON30139), and R3 generation plants containing the P-ScBV promoter (line TA_S2520 generated from pMON30159). Yield data (grams) and fertility data (as % of total plants) were collected for each line. Data are provided in Table 11 below.

TABLE 11

Pressure Test of R2 Wheat Plants				
Promoter	Construct	Line	Yield (g)	Fertility (%)
N/A	N/A	Non-transgenic Bobwhite	23.1	93.5
P-CaMV.e35S and P-Os.Act1	pMON30139	33391	24.1	91.2
P-ScBV	pMON30159	TA_S2520	21.0	87.5
P-2xA1-B3/P-Os.Act1-2	pMON43647	TA_S7535	22.2	91.5
P-2xA1-B3/P-Os.Act1-2	pMON43647	TA_S7890	23.0	95.8
P-2xA1-B3/P-Os.Act1-2	pMON43647	TA_S9215	21.5	95.2
P-4xAS-1/P-Os.Act1-3	pMON43646	TA_S9240	20.6	95.0

From this analysis it is apparent that the use of the P-4xAS-1/P-Os.Act1-3 (SEQ ID NO: 14), P-2xA1-B3/P-Os.Act1-2 (SEQ ID NO: 19), and P-2xA1-B3/P-Os.Act1-3 (SEQ ID NO: 20) promoters in plant CP4 expression cassettes confers glyphosate tolerance to vegetative tissues in wheat plants without negatively impacting fertility. In wheat, cassettes containing the rice actin promoter without these elements produce plants with low vegetative tolerance and are not useful for the production of glyphosate tolerant plants. Therefore one advantage of the chimeric promoters in wheat is that vegetative and reproductive tolerance can be achieved without the use of a double CP4 cassette.

Field trails were performed with R4 generation plants to assess the performance of selected lines. Field trails were done at eight US sites with 4 replications each and treatments of 0, 64, and 128 oz/A Roundup® Ultra equiva-

lent. Each treatment was arranged as a separate, randomized complete block in order to collect equivalence data on unsprayed non-transformed Bobwhite plants. All events had vegetative and reproductive tolerance comparable to the lead event 33391 at doses up to 128 oz/A of Roundup® Ultra. Yield was measured in tons/hectar (T/ha). Data are provided in Table 12 below.

TABLE 12

Yield Data from Field Trials of R4 Plants (T/ha)					
Promoter	Construct	Line	128 oz/A	64 oz/A	No spray
P-CaMV.e35S and P-Os.Act1	pMON30139	33391	3.52	3.58	3.63
P-ScBV	pMON30159	TA_S2520	3.78	3.79	3.69
P-2xA1-B3/ P-Os.Act1-2	pMON43647	TA_S7535	3.72	3.68	3.51
P-2xA1-B3/ P-Os.Act1-2	pMON43647	TA_S7890	3.79	3.72	3.70

TABLE 12-continued

Yield Data from Field Trials of R4 Plants (T/ha)					
Promoter	Construct	Line	128 oz/A	64 oz/A	No spray
P-2xA1-B3/ P-Os.Act1-2	pMON43647	TA_S9215	3.59	3.57	3.42
P-4xAS-1/ P-Os.Act1-3	pMON43646	TA_S9240	3.58	3.81	3.66

Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims. All publications and published patent documents cited in this specification are incorporated herein by reference to the same extent as if each individual publication or patent application is specifically and individually indicated to be incorporated herein by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 35

<210> SEQ ID NO 1
 <211> LENGTH: 612
 <212> TYPE: DNA
 <213> ORGANISM: Cauliflower mosaic virus

<400> SEQUENCE: 1

```

gggccgattg agacttttca acaaagggta atatccggaa acctcctcgg attccattgc      60
ccagctatct gtcactttat tgtgaagata gtggaaaagg aagggtggctc ctacaaatgc      120
catcattgcg ataaggaaa ggccatcggt gaagatgcct ctgccgacag tggteccaaa      180
gatggacccc caccacagag gagcatcgtg gaaaaagaag acgttccaac cacgtcttca      240
aagcaagtgg attgatgtga tgggccgatt gagacttttc aacaaagggt aatatccgga      300
aacctcctcg gattccattg occagctatc tgtcacttta ttgtgaagat agtggaagag      360
gaagtggtct cctacaaatg ccatcattgc gataaaggaa aggccatcgt tgaagatgcc      420
tctgccgaca gtggtcccaa agatggaccc ccaccacga ggagcatcgt ggaaaaagaa      480
gacgttccaa ccacgtcttc aaagcaagtg gattgatgtg atatctccac tgacgtaagg      540
gatgacgcac aatcccacta tccttcgcaa gaccttcct ctatataagg aagttcattt      600
catttggaga gg                                          612
    
```

<210> SEQ ID NO 2
 <211> LENGTH: 240
 <212> TYPE: DNA
 <213> ORGANISM: Cauliflower mosaic virus

<400> SEQUENCE: 2

```

catcgttgaa gatgcctctg ccgacagtgg tcccaaagat ggacccccac ccacctcgac      60
catcgttgaa gatgcctctg ccgacagtgg tcccaaagat ggacccccac ccacctcgac      120
catcgttgaa gatgcctctg ccgacagtgg tcccaaagat ggacccccac ccacctcgac      180
catcgttgaa gatgcctctg ccgacagtgg tcccaaagat ggacccccac ccacctcgac      240
    
```

<210> SEQ ID NO 3
 <211> LENGTH: 221

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Cauliflower mosaic virus

<400> SEQUENCE: 3
agcttctgac gtaagggatg acgcacctga cgtaagggat gacgcacctg acgtaagggg    60
tgacgcacct gacgtaaggg atgacgcact cgagatcccc atctccactg acgtaagggg    120
tgacgcacaa tcccactatc cttcgcaaga ccttcctct atataaggaa gttcatttca    180
tttgagagg acacgctgac aagctagctt ggctgcaggt a                               221

<210> SEQ ID NO 4
<211> LENGTH: 125
<212> TYPE: DNA
<213> ORGANISM: Cauliflower mosaic virus

<400> SEQUENCE: 4
catcgtggaa aaagaagacg ttccaaccac gtcttcaaag caagtggatt gatgtgataa    60
gctcctcatc gtggaaaaag aagacgttcc aaccacgtct tcaaagcaag tggattgatg    120
tgata                                               125

<210> SEQ ID NO 5
<211> LENGTH: 332
<212> TYPE: DNA
<213> ORGANISM: Cauliflower mosaic virus

<400> SEQUENCE: 5
catcgttgaa gatgcctctg cgcacagtgg toccaagat ggacccccac ccacgaggag    60
catcgttgaa aaagaagacg ttccaaccac gtcttcaaag caagtggatt gatgtgatat    120
ctccactgac gtaagggatg acgcacaatc cactatcct tcgaggcctc atcgttgaag    180
atgcctctgc cgacagtggc cccaagatg gacccccacc cagcaggagc atcgtggaaa    240
aagaagacgt tccaaccacg tcttcaaagc aagtggattg atgttatatc tccactgacg    300
taagggatga cgcacaatcc cactatcctt cg                               332

<210> SEQ ID NO 6
<211> LENGTH: 524
<212> TYPE: DNA
<213> ORGANISM: Cauliflower mosaic virus

<400> SEQUENCE: 6
ggtcgcatgt gagacttttc aacaaaggtt aatatccgga aacctcctcg gattccattg    60
cccagctatc tgtcacttta ttgtgaagat agtggaaaag gaagtggtgct cctacaaatg    120
ccatcattgc gataaaggaa aggccatcgt tgaagatgcc tctgccgaca gtggtcccaa    180
agatggacce ccaccacgga ggagcatcgt ggaaaaagaa gacgttccaa ccacgtcttc    240
aaagcaagtg gattgatgtg atggtccgat gtgagacttt tcaacaaagg gtaatatccg    300
gaaacctcct cggattccat tgcccagcta tctgtcactt tattgtgaag atagtggaaa    360
aggaagtggt ctctacaaa tgccatcatt cgcataaagg aaaggccatc gttgaagatg    420
cctctgccga cagtggtecc aaagatggac cccccccac gaggagcatc gtggaaaaag    480
aagacgttcc aaccacgtct tcaaagcaag tggattgatg tgat                               524

<210> SEQ ID NO 7
<211> LENGTH: 921
<212> TYPE: DNA
<213> ORGANISM: Oryza sativa

```

-continued

<400> SEQUENCE: 7

```

tcgaggatcat tcatatgctt gagaagagag tgggatagat ccaaaataaa acaaaggtaa    60
gattacctgg tcaaaagtga aaacatcagt taaaaggtag tataaagtaa aatatacggta    120
ataaaaggtag gcccaaagtg aaatttactc ttttctacta ttataaaaat tgaggatggt    180
tttgtcggta ctttgatacg tcatttttgt atgaattggg ttttaagttt attcgctttt    240
ggaaatgcat atctgtatct gagtcggggt ttaagtctgt ttgcttttgt aaatacagag    300
ggatttgtat aagaaatctc tttagaaaaa cccatagctc aatttgacat aatttttgag    360
aaaaatatat attcaggcga attctcacia tgaacaataa taagattaaa atagctttcc    420
cccgttcgag cgcattggga ttttttctag taaaaataaa agataaactt agactcaaaa    480
catttcaaaa aacaaccctc aaagtctcta aagcccaaag tgctatccac gatccatagc    540
aagcccagcc caaccacaac caaccagcc caccacagtc cagccaactg gacaatagtc    600
tccacacccc cccactatca ccgtgagttg tccgcacgca ccgcacgtct cgcagccaaa    660
aaaaaaaaaga aagaaaaaaaa agaaaaagaa aaaacagcag gtgggtccgg gtcgtggggg    720
ccggaaacgc gaggaggatc gcgagccagc gacgaggccg gccctccctc cgcttccaaa    780
gaaacgcccc ccatcgccac tatatacata cccccccctc tcctccctc cccccaacc    840
taccaccacc accaccacca cctccacctc ctccccctc gctgccggac gacgagctcc    900
tccccctcc cctcccgccg c                                     921

```

<210> SEQ ID NO 8

<211> LENGTH: 1577

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 8

```

taaatgacat cagatacacg cttgtgaacc atctttaaag tattgatgga ctcttcaacta    60
tgaaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaatat    120
tagagggtcca aattcaaaaa aatgtogtat tgaatcattc cactactaaa ttgggtcaat    180
gtcagattta aacagcctag ggataattta gtgagatag agattctact tcaacatat    240
actaatccta aatctctagc aactttttat ataagctata aatatcatga aaatgtattt    300
taatcgtttc ataatttatg cagtcacact aatggaaaaa aggccaatta ttattatttt    360
cttcagacta taaatgaaaa cataaattaa aatgcagatt agtttaaaat ttaataagt    420
aagtaaaatg cttatagcct tatacaaaat catatttggg agtttctaac attggtgcaa    480
tttgttatca caaatcacag taatatttgt atactaatta gtaattacaa ctatacacia    540
atttaaatgg gtaatcatat atttgtgtcc agtggattga acaaatatgc tcggcccatg    600
cggaagtaat gccaattttg ggtgagtaaa gcccatgcga aattttcaca taagaaatgc    660
atgctttttg ttttcaacga catgagttgc atgcttttta tcattgctta tatagttgca    720
agtttgcaac tccttgatat tttttttatg tagacactac taccaccaa aacttttggt    780
ctgcttattc ttgtttacta tgtaaaaaaa ataaatgaat tgtttattta ctccgatttg    840
atggagctcg gtttatgagg ttttatagcc tttacagaaa attgatagtt acaaaaatat    900
ttttcaaaaa taaaagggta aaaccgctat ttcaagttgt tattgttttg ggggactgga    960
tttgaatga aatatagaac cggaaaaaaa ggtgagccga agtcgaagcc tttggaccg    1020
tttttatatt tactcctccc attcctctct ccttcaatcc ttccttctc ctctccctt    1080
cttctcttc cctctttca ttttccagcc actacaaact tttctatctc tactttttt    1140

```

-continued

```

cctctcgatt tcaggtactt tttgagacc tttgttgga ttttcgaaca cacaccccaa 1200
ttacgtttga tttttgatcc cgcacogatt tcaattcatc cgtttctgag tttcttttgg 1260
atctgggtgt cttgagctaa tcttttcgat ctggtgttta tcgattttac tcatgcgtat 1320
gttcattaca ccatttgta tttgtttaat caacccaaaag actcatgttt tccaaatgtc 1380
tttaataata tttttctgat tgaattttat aatatttaca tgattctgga tccagaatat 1440
ccttctctct cttccatttt gtcctgtatt gatttgcctt tgaaaaagga ttgttctttg 1500
tatctgtatt ggtgaaaaag gattgttatt tgttgataaa aatttgatct ttaacaatg 1560
tttggttttg cataaag 1577

```

```

<210> SEQ ID NO 9
<211> LENGTH: 1184
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 9

```

```

aagctagctt gtcgaccatc gttgaagatg cctctgccga cagtgggcc aaagatggac 60
ccccaccac ctcgaccatc gttgaagatg cctctgccga cagtgggcc aaagatggac 120
ccccaccac ctcgaccatc gttgaagatg cctctgccga cagtgggcc aaagatggac 180
ccccaccac ctcgaccatc gttgaagatg cctctgccga cagtgggcc aaagatggac 240
ccccaccac ctcgacagct tactcgaggt cattcatatg cttgagaaga gagtcgggat 300
agtccaaaat aaaacaaagg taagattacc tggtaaaaag tgaaacatc agttaaagg 360
tggataaaag taaaatctg gtaataaaag gtggcccaaa gtgaaattta ctcttttcta 420
ctattataaa aattgaggat gttttgtcgc gtaactttgat acgtcatttt tgtatgaatt 480
ggtttttaag tttattcgtc tttggaaatg catatctgta tttgagtcgg gttttaagtt 540
cgtttgcttt tgtaaataca gagggatttg tataagaaat atcttttagaa aaaccatata 600
gctaatttga cataattttt gagaaaaata tatattcagg cgaattotca caatgaacaa 660
taataagatt aaaatagctt tcccccgtyg cagcgcgatgg gtattttttc tagtaaaaat 720
aaaagataaa cttagactca aaacatttac aaaaacaacc ctaaagttc ctaaagccca 780
aagtgcatac cacgatccat agcaagccca gcccaaccca acccaaccca gccacccca 840
gtccagccaa ctggacaata gtctccacac cccccacta tcaccgtgag ttgtccgcac 900
gcaccgcacy tctcgcagcc aaaaaaaaa agaaagaaaa aaaagaaaa gaaaaaacag 960
caggtgggtc cgggtcgtgg gggccgaaa cgcgaggagg atcgcgagcc agcgacgagg 1020
ccggccctcc ctccgcttcc aaagaacgc cccccatcgc cactatatac ataccccccc 1080
ctctctccc atcccccaa ccctaccacc accaccacca ccacctccac ctctcccccc 1140
ctcgtgccc gacgacgagc tctccccccc tccccctcg ccgc 1184

```

```

<210> SEQ ID NO 10
<211> LENGTH: 1184
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 10

```

```

aagcttactc gaggtcattc atatgcttga gaagagatc gggatagtcc aaaataaac 60

```

-continued

```

aaaggaaga ttacctggc aaaagtgaac acatcagtta aaagtggtgta taaagtaaaa 120
tatcggtaat aaaaggtggc ccaaagtgaac atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgtcggctact ttgatcagtc atttttgtat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcataat ctgtatttga gtcggggtttt aagttcgttt gcttttgtaa 300
atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tagcttctcg accatcgttg aagatgcctc 420
tgccgacagt ggtcccaaaag atggaccccc acccacctcg accatcgttg aagatgcctc 480
tgccgacagt ggtcccaaaag atggaccccc acccacctcg accatcgttg aagatgcctc 540
tgccgacagt ggtcccaaaag atggaccccc acccacctcg accatcgttg aagatgcctc 600
tgccgacagt ggtcccaaaag atggaccccc acccacctcg acaattctca caatgaacaa 660
taataagatt aaaatagctt tcccccgctg cagcgcagtg gtatttttct tagtaaaaaat 720
aaaagataaa cttagactca aacattttac aaaaaacaacc cctaaagtct ctaaagccca 780
aagtgtctatc cagcatccat agcaagccca gcccaaccca acccaaccca gcccacccca 840
gtccagccaa ctggacaata gtctccacac cccccacta tcaccgtgag ttgtccgcac 900
gcaccgcagc tctcgcagcc aaaaaaaaaa agaaagaaaa aaaagaaaaa gaaaaaacag 960
caggtgggtc cgggtcgtgg gggccggaac cgcgaggagg atcgcgagcc agcgcagagg 1020
ccggccctcc ctccgcttcc aaagaacgc cccccatcgc cactatatac atccccccc 1080
ctctctccc atccccccaa ccctaccacc accaccacca ccacctccac ctctcccc 1140
ctcgtgccc gagcagcagc tcctcccccc tccccctccg ccgc 1184

```

```

<210> SEQ ID NO 11
<211> LENGTH: 1176
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 11

```

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaac 60
aaaggaaga ttacctggc aaaagtgaac acatcagtta aaagtggtgta taaagtaaaa 120
tatcggtaat aaaaggtggc ccaaagtgaac atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgtcggctact ttgatcagtc atttttgtat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcataat ctgtatttga gtcggggtttt aagttcgttt gcttttgtaa 300
atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tctcacaatg aacaataata agattaaaaat 420
agctttcccc cgttcgagcgc catgggtatt ttttctagta aaaataaaag ataaacttag 480
actcaaaaca tttacaaaaa caaccctaa agttcctaaa gcccaaatg ctatccacga 540
tccatagcaa gccagccca acccaaccca acccagccca cccagtcga gccaaactgga 600
caatagtctc cacaccccc cactatcacc gtgagttgtc cgcacgcacc gcacgtctcg 660
cagccaaaaa aaaaaagaaa gaaaaaaaag aaaaaaaaaa aacagcaggt gggtcgggt 720
cgtgggggccc ggaaaacgca ggaggatcgc gagccagcga cgaggagctt gtcgaccatc 780
ggtgaagatg cctctgccga cagtggtccc aaagatggac cccacccac ctcgaccatc 840
ggtgaagatg cctctgccga cagtggtccc aaagatggac cccacccac ctcgaccatc 900
ggtgaagatg cctctgccga cagtggtccc aaagatggac cccacccac ctcgaccatc 960

```

-continued

```

gttgaagatg cctctgccga cagtgggtccc aaagatggac ccccacccac ctegaccct 1020
ccctccgctt ccaaagaaac gccccccatc gccactatat acataccccc ccctctcttc 1080
ccatccccc aaccctacca ccaccaccac caccacctcc acctcctccc ccctcgtgctc 1140
cggacgacga gctcctcccc cctccccctc cgccgc 1176

```

```

<210> SEQ ID NO 12
<211> LENGTH: 1158
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 12

```

```

aagctagctt ctgacgtaag ggatgacgca cctgacgtaa gggatgacgc acctgacgta 60
agggatgacg cacctgacgt aagggatgac gcaactcgaga tccccatctc cactgacgta 120
agggatgacg cacaatccca ctatccttcg caagaccctt cctctatata aggaagtcca 180
tttcatttgg agaggacacg ctgacaagct agcttggtcg caggtagatc agcttactcg 240
aggtcattca tatgcttgag aagagagtcg ggatagtcca aaataaaaca aaggaagat 300
tacctggtca aaagtgaaaa catcagttaa aaggtggtat aaagtaaaat atcggttaata 360
aaagtggtgc caaagtgaaa tttactcttt tctactatta taaaaattga ggatgttttt 420
gtcggtaactt tgatacgtca tttttgatg aattggtttt taagtttatt cgcttttggg 480
aatgcatatc tgtatttgag tcgggtttta agttcgtttg cttttgtaaa tacagagggg 540
tttgataag aaatatcttt agaaaaacc atatgcta attgacataat ttttgagaaa 600
aatatatatt caggcgaatt ctcaaatga acaataataa gattaaaata gctttcccc 660
gttgacgacg atgggtattt tttctagtaa aaataaaaga taaacttaga ctcaaaacat 720
ttacaaaaac aaccctaaa gttcctaaa cccaaagtgc tatccacgat ccatagcaag 780
cccagcccaa cccaacccaa cccagcccac cccagtcag ccaactggac aatagtctcc 840
acaccccccc actatcacgg tgagttgtcc gcacgcaccg cacgtctcgc agccaaaaaa 900
aaaaagaaag aaaaaaaaga aaaagaaaa acagcaggtg ggtccgggtc gtggggggcg 960
gaaacgcgag gaggatcgcg agccagcgac gaggccggcc ctccctcgc tccaaagaa 1020
acgcccccca tegccactat atacataccc cccctctcc tcccateccc ccaaccctac 1080
caccaccacc accaccacct ccacctctc cccctcgtc gccggacgac gagctcctcc 1140
ccctcccccc tccgcccgc 1158

```

```

<210> SEQ ID NO 13
<211> LENGTH: 1158
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 13

```

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaaac 60
aaagtaaga ttacctggtc aaaagtgaaa acatcagtta aaagtggtg taaagtaaaa 120
tatcggtaat aaaagtggtc ccaaagtgaa atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgcggttact ttgatcgtc atttttgat gaattggttt ttaagtttat 240
tcgcttttgg aatgcatat ctgtatttga gtcgggtttt aagttcgttt gcttttgtaa 300

```

-continued

```

atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tagcttctga cgtaagggat gacgcacctg 420
acgtaagga tgacgcacct gacgtaaggg atgacgcacc tgacgtaagg gatgacgcac 480
tcgagatccc catctccact gacgtaaggg atgacgcaca atcccactat ccttcgcaag 540
acccttctc tatataagga agttcatttc atttggagag gacacgctga caagctagct 600
tgggtgcagg tagatcaatt ctcaaatga acaataataa gattaaaata gctttccccc 660
gttgacgagc atgggtattt tttctagtaa aaataaaaga taaacttaga ctcaaacat 720
ttacaaaaac aaccctaaa gttcctaaag cccaaagtgc tatccacgat ccatagcaag 780
cccagcccaa cccaacccaa ccagccccc cccagtcag ccaactggac aatagtctcc 840
acaccccccc actatcaccc tgagttgtcc gcacgcaccg cacgtctcgc agccaaaaaa 900
aaaaagaaag aaaaaaaga aaaagaaaa acagcaggtg ggtccgggtc gtggggggcg 960
gaaacgcgag gaggatcgcg agccagcgac gaggcgggcc ctccctccgc ttccaaagaa 1020
acgcccccca tcgccactat atacataccc cccctctcc tcccatcccc ccaaccctac 1080
caccaccacc accaccacct ccacctctc cccctcgcct gccggacgac gagctcctcc 1140
ccccccccc tcgcccgc 1158

```

```

<210> SEQ ID NO 14
<211> LENGTH: 1150
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 14

```

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaaac 60
aaaggaaga ttacctggtc aaaagtgaac acatcagtta aaagtggtg taaagtaaaa 120
tatcggtaat aaaaggtggc ccaaagtga atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgcggtact ttgatacgtc atttttgtat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcata ctgtatttga gtcgggtttt aagttcgttt gcttttgtaa 300
atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tctcacaatg aacaataata agattaaaat 420
agctttcccc cgttgcagcg catgggtatt ttttctagta aaaataaaag ataaacttag 480
actcaaaaac ttacaaaaa caaccctaa agttcctaaa gcccaaagtg ctatccacga 540
tccatagcaa gccagccca acccaaccca accagccca cccagtcga gccaaactgga 600
caatagtctc cacaccccc cactatcacc gtgagttgtc cgcacgcacc gcacgtctcg 660
cagccaaaaa aaaaaagaaa gaaaaaaaag aaaaagaaaa aacagcaggt ggtccgggt 720
cgtggggggc ggaaacgcga ggagatcgc gagccagcga cgaggagctt ctgacgtaag 780
ggatgacgca cctgacgtaa gggatgacgc acctgacgta agggatgacg cacctgacgt 840
aaggatgac gcaactgaga tccccatctc cactgacgta agggatgacg cacaatcca 900
ctatccttcg caagaccctt cctctatata aggaagtca tttcatttgg agaggacag 960
ctgacaagct agcttggctg caggtagatc ccctccctcc gcttccaaag aaacgcccc 1020
catcgccact atatacata cccccctct cctcccatcc cccaacctc accaccacca 1080
ccaccaccac ctccacctcc tccccctcg ctgccggagc acgagctcct cccccctcc 1140
cctccgcccgc 1150

```

-continued

```

<210> SEQ ID NO 15
<211> LENGTH: 1065
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 15
aagcttaggc ctcatcgtgg aaaaagaaga cgttccaacc acgtttcaa agcaagtgga    60
ttgatgtgat aagctcctca tcgtggaaaa agaagacgtt ccaaccacgt cttcaaagca    120
agtggattga tgtgataagc ttactcgagg tcattcatat gcttgagaag agagtcggga    180
tagtcacaaa taaaacaaag gtaagattac ctggcacaaa gtgaaaacat cagttaaaag    240
gtggtataaa gtaaaatata ggtaataaaa ggtggcccaa agtgaaatct actcttttct    300
actattataa aaattgagga tgtttttgtc ggtactttga tacgtcattt ttgtatgaat    360
tggtttttaa gttttatcgc ttttggaat gcatactctg atttgagtcg ggttttaagt    420
tcgtttgctt ttgtaaatat agagggattt gtataagaaa tatctttaga aaaaccata    480
tgctaatttg acataathtt tgagaaaaat atatattcag gcgaattctc acaatgaaca    540
ataataagat taaaatagct ttccccctgt gcagcgcgat ggtatttttt ctagtaaaaa    600
taaaagataa acttagactc aaaacattta caaaaacaac ccctaaagtt cctaaagccc    660
aaagtgctat ccacgatcca tagcaagccc agcccaacc aaccacaacc agcccacccc    720
agtccagcca actggacaat agtctccaca cccccccact atcaccgtga gttgtccgca    780
cgcaccgcac gtctcgcagc caaaaaaaaa aagaaagaaa aaaaagaaaa agaaaaaaca    840
gcaggtgggt ccgggtcgtg ggggcccggaa acgcgaggag gatcgcgagc cagcgcagag    900
gccggccctc cctccgcttc caaagaaacg ccccccatcg ccactatata catacccccc    960
cctctctctc catcccccca accctaccac caccaccacc accacctcca cctctctccc   1020
cctcgtgccc ggacgacgag ctctctcccc ctccccctcc gccgc                               1065

```

```

<210> SEQ ID NO 16
<211> LENGTH: 1073
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 16
aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaac    60
aaaggtaaaga ttacctggtc aaaagtgaac acatcagtta aaaggtggta taaagtaaaa    120
tatcggtaat aaaaggtggc ccaaagtgaac atttactctt ttctactatt aaaaaaattg    180
aggatgtttt tgtcggctact ttgatcagtc atttttgtat gaattggttt ttaagtttat    240
tcgcttttgg aaatgcatac ctgtatttga gtcgggtttt aagttcgttt gcttttgtaa    300
atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa    360
tttttgagaa aaatatatat tcaggcgaat tagcttaggc ctcatcgtgg aaaaagaaga    420
cgttccaacc acgtcttcaa agcaagtgga ttgatgtgat aagctcctca tcgtggaaaa    480
agaagacgtt ccaaccacgt cttcaaagca agtggattga tgtgataagc taattctcac    540
aatgaacaat aataagatta aaatagcttt cccccgttgc agcgcagggg tattttttct    600
agtaaaaata aaagataaac ttagactcaa aacatttaca aaaacaaccc ctaaagtccc    660

```

-continued

```

taaagcccaa agtgctatcc acgatccata gcaagcccag cccaacccaa cccaaccag 720
cccccccag tccagccaac tggacaatag tctccacacc cccccactat caccgtgagt 780
tgtcgcacg caccgcacgt ctgcgagcca aaaaaaaaaa gaaagaaaaa aaagaaaaag 840
aaaaaacagc aggtgggtcc gggctcgtggg ggccggaaac gcgaggagga tcgagagcca 900
gcgacgaggc cgccctccc tccgcttcca aagaaacgcc ccccatcgcc actatataca 960
tcccccccc tctctccca tcccccaac cctaccacca ccaccaccac cacctccacc 1020
tcctcccccc tcgctgccgg acgacgagct cctccccct cccccccgc cgc 1073

```

```

<210> SEQ ID NO 17
<211> LENGTH: 1065
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 17

```

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaac 60
aaaggaaga ttacctggtc aaaagtgaac acatcagtta aaaggtgta taaagtaaaa 120
tatcggtaat aaaaggtggc ccaaagtga atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgtcggctact ttgatcgtc atttttgtat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcata ctgtatttga gtcgggtttt aagttcgttt gcttttgtaa 300
atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tctcacaatg aacaataata agattaaaat 420
agctttcccc cgttgcagcg catgggtatt ttttctagta aaaataaaag ataaacttag 480
actcaaaaca tttacaaaa caaccocaa agttcctaaa gcccaaagtg ctatccacga 540
tccatagcaa gccagccca acccaaccca acccagccca cccagtcga gccaaactgga 600
caatagtctc cacaccccc cactatcacc gtgagttgtc cgcacgcacc gcacgtctcg 660
cagccaaaaa aaaaaagaaa gaaaaaaaaa aaaaagaaaa aacagcaggt gggtcggggt 720
cgtgggggccc gaaaacgcga ggaggatcgc gagccagcga cgaggagctt aggccctcctc 780
gtggaaaaag aagacgttcc aaccacgtct tcaaagcaag tggattgatg tgataagctc 840
ctcatcgtgg aaaaagaaga cgttccaacc acgtcttcaa agcaagtgga ttgatgtgat 900
aagtcctctc cctccgcttc caaagaaacg cccccatcg ccaactatata cataccccc 960
cctctctctc catcccccca accctaccac caccaccacc accacctcca cctcctcccc 1020
cctcgtctgc ggacgacgag ctctcctccc ctccccctcc gccgc 1065

```

```

<210> SEQ ID NO 18
<211> LENGTH: 1273
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 18

```

```

aagcttaggc ctcatcgttg aagatgcctc tgccgacagt ggtcccaaag atggaccccc 60
accacagagg agcatcgtgg aaaaagaaga cgttccaacc acgtcttcaa agcaagtgga 120
ttgatgtgat atctccactg acgtaaggga tgacgcacaa tcccactatc cttcgaggcc 180
tcatcgttga agatgcctct gccgacagtg gtcccaaaga tggacccccca cccacgagga 240
gcatcgtgga aaaagaagac gttccaacca cgtcttcaaa gcaagtggat tgatgttata 300

```

-continued

```

tctocactga cgtaagggat gacgcacaat cccactatcc ttcgaagctt actcogaggtc 360
attcatatgc ttgagaagag agtcgggata gtccaaaata aaacaaaggt aagattacct 420
ggtcaaaagt gaaaacatca gttaaaaggt ggtataaagt aaaatatcgg taataaaagg 480
tggcccaaag tgaaatctac tcttttctac tattataaaa attgaggatg tttttgtcgg 540
tactttgata cgtcattttt gtatgaattg gtttttaagt ttattcgctt ttggaaatgc 600
atatctgtat ttgagtcggg ttttaagttc gtttgctttt gtaaatcacag agggatttgt 660
ataagaaata tcttttagaaa aaccatgatg ctaatttgac ataatttttg agaaaaatat 720
atattcaggc gaattctcac aatgaacaat aataagatta aaatagcttt cccccgttgc 780
agcgcatggg tattttttct agtaaaaata aaagataaac ttagactcaa aacatttaca 840
aaaaacaacc ctaaagtctc taaagcccaa agtgctatcc acgatccata gcaagcccag 900
cccaccccaa cccaaccag cccaccccag tccagccaac tggacaatag tctccacacc 960
ccccactat caccgtgagt tgtccgcacg caccgcacgt ctcgcagcca aaaaaaaaaa 1020
gaaagaaaaa aaagaaaaag aaaaaacagc aggtgggtcc gggctcgtggg ggcgggaaac 1080
gcgaggagga tcgagagcca gcgacgaggc cggcctccc tccgcttcca aagaaacgcc 1140
ccccatcgcc actatataca tcccccccc tctcctccca tcccccaac cctaccacca 1200
ccaccaccac cacctccacc tctcccccc tcgctgcccg acgacgagct cctccccct 1260
ccccctccgc cgc 1273

```

```

<210> SEQ ID NO 19
<211> LENGTH: 1281
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 19

```

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaaac 60
aaaggtaaaga ttacctggtc aaaagtgaaa acatcagtta aaaggtggta taaagtaaaa 120
tatcggtaat aaaaggtggc ccaaagtgaa atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgcggttact ttgatcgtc atttttgtat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcatat ctgtatttga gtcgggtttt aagttcgttt gcttttgtaa 300
atacagagggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tagcttaggc ctcacgttg aagatgcctc 420
tgccgacagt ggtcccaaag atggaccccc acccacgagg agcatcgtgg aaaaagaaga 480
cgttccaacc acgtcttcaa agcaagtgga ttgatgtgat atctccactg acgtaagggg 540
tgacgcacaa tcccactatc cttcgaggcc tcatcgttga agatgcctct gccgacagtg 600
gtcccaaaga tggaccccca cccacgagga gcacgtgga aaaagaagac gttccaacca 660
cgtcttcaaa gcaagtgatg tgatgttata tctccactga cgtaagggat gacgcacaat 720
cccactatcc ttcgaageta attctcacia tgaacaataa taagattaaa atagctttcc 780
cccgttgagc cgcattgggta ttttttctag taaaaataaa agataaaactt agactcaaaa 840
catttcaaaa aacaaccctc aaagttccta aagcccaaag tgctatccac gatccatagc 900
aagcccagcc caaccacaac caaccagcc caccacagtc cagccaactg gacaatagtc 960
tccacacccc cccactatca ccgtgagttg tccgcacgca ccgacgtct cgcagccaaa 1020

```

-continued

```

aaaaaaaaaga aagaaaaaaaa agaaaaaaaaa aaaacagcag gtgggtccgg gtcgtggggg 1080
ccggaaacgc gaggaggatc gcgagccagc gacgaggccg gccctccctc cgcttccaaa 1140
gaaacgcccc ccategccac tatatacata cccccccctc tcctcccatc cccccaacct 1200
taccaccacc accaccacca cctccacctc ctccccctc gctgccggac gacgagctcc 1260
tccccctcc cctcccgccg c 1281

```

```

<210> SEQ ID NO 20
<211> LENGTH: 1273
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 20

```

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtc aaaataaaac 60
aaagtaaga ttacctggtc aaaagtgaac acatcagtta aaagtggtgta taaagtaaaa 120
tatcgtaat aaaaggtggc ccaaagtgaac atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgcggtact ttgatcagtc atttttctat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcatac ctgtatttga gtcgggtttt aagttcgttt gcttttgtaa 300
atacagaggg atttgataaa gaaatatctt tagaaaaacc catatgctaa ttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tctcacaatg aacaataata agattaaaat 420
agctttcccc cgttgccagc catgggtatt tttctagta aaaataaaag ataaacttag 480
actcaaaaca ttacaaaaa caaccctcaa agttcctaaa gcccaaagtg ctatccacga 540
tccatagcaa gccagccca acccaaccca acccagccca cccagtcga gccaaactgga 600
caatagtctc cacaccccc cactatcacc gtgagttgtc cgcacgcacc gcacgtctcg 660
cagccaaaaa aaaaaagaaa gaaaaaaaaag aaaaaaagaaa aacagcaggt gggtcggggt 720
cgtggggggc ggaaacgcga ggaggatcgc gagccagcga cgaggagctt aggcctcatc 780
gttgaagatg cctctgccga cagtggtccc aaagatggac cccacccac gaggagcatc 840
gtggaaaaag aagacgttcc aaccacgtct tcaaagcaag tggattgatg tgatatctcc 900
actgacgtaa gggatgacgc acaatccac tatccttcga ggctcatcg ttgaagatgc 960
ctctgccgac agtgggtccc aagatggacc cccacccacg aggagcatcg tggaaaaaga 1020
agacgttcca accacgtctt caaagcaagt ggattgatgt tatatctcca ctgacgtaag 1080
ggatgacgca caatcccact atccttcgaa gctccctccc tccgcttcca aagaaacgcc 1140
ccccatgcc actatataca tcccccccc tctcctccca tcccccaac cctaccacca 1200
ccaccaccac cacctccacc tcctcccccc tcgctgccgg acgacgagct cctccccct 1260
ccccctccgc cgc 1273

```

```

<210> SEQ ID NO 21
<211> LENGTH: 1467
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 21

```

```

aagctagctt ctgcaggctc gatgtgagac ttttcaaaa aggtaatat ccggaacct 60
cctcggatc cattgccag ctatctgtca ctttattgtg aagatagtg aaaaggaag 120
tggtcctac aaatgccatc attgcgataa aggaaggcc atcgttgaag atgctctgc 180

```

-continued

```

cgacagtggg cccaaagatg gacccccacc cacgaggagc atcgtggaaa aagaagacgt 240
tccaaccacg tcttcaaagc aagtggattg atgtgatggg ccgatgtgag acttttcaac 300
aaagggtaat atccggaaac ctccctcgat tocattgccc agctatctgt cactttattg 360
tgaagatagt ggaaaaggaa ggtggctcct acaaatgcca tcattgcatg aaaggaaagg 420
ccatcgttga agatgcctct gccgacagtg gtcccaaaga tggacccccca cccacgagga 480
gcatcgtgga aaaagaagac gttccaacca cgtcttcaa gcaagtggat tgatgtgata 540
gcttactcga ggctattcat atgcttgaga agagagtcgg gatagtccaa aataaaacaa 600
aggtaaagatt acctgggtcaa aagtgaaac atcagttaaa aggtgggtata aagtaaaata 660
tcggtaataa aagggtggccc aaagtgaat ttactctttt ctactattat aaaaattgag 720
gatgtttttg tcggtacttt gatcgtcat ttttgatga attggttttt aagtttattc 780
gcttttgaaa atgcatatct gtatttgagt cgggttttaa gttcgtttgc ttttgtaaat 840
acagagggat ttgtataaga aatatcttta gaaaaacca tatgctaatt tgacataatt 900
tttgagaaaa atatatattc aggcgaattc tcacaatgaa caataataag attaaaatag 960
ctttcccccg ttgcagcgca tgggtatttt ttctagtaaa aataaaagat aaacttagac 1020
tcaaaacatt tacaaaaaca acccctaaag ttcctaaagc ccaaaagtct atccacgatc 1080
catagcaagc ccagcccaac ccaacccaac ccagcccacc ccagtccagc caactggaca 1140
atagtctcca cccccccca ctatcacctg gagttgtccg cacgcaccgc acgtctcgca 1200
gccaaaaaaa aaaagaagaa aaaaaagaa aaagaaaaaa cagcaggtgg gtcggggtcg 1260
tgggggcccg aaacgcgagg aggatcgcga gccagcgagc aggcgggccc tcctccgct 1320
tccaaagaaa cccccccat cgccactata tacatacccc ccctctcct cccatcccc 1380
caacctacc accaccacca ccaccacctc cacctcctcc ccctcctcg cgggacgacg 1440
agctcctccc ccctcccct ccgcccgc 1467

```

<210> SEQ ID NO 22

<211> LENGTH: 1467

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 22

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaaac 60
aaaggaaga ttacctggtc aaaagtgaac acatcagtta aaagtggtgta taaagtaaaa 120
tatcggtaat aaaaggtggc ccaaagtgaac atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgtcgttact ttgatcgtc atttttgtat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcatac ctgtatttga gtcgggtttt aagttcgttt gcttttgtaa 300
atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tagcttctgc aggtccgatg tgagactttt 420
caacaaaggg taatatccgg aaacctcctc ggattccatt gcccagctat ctgtcacttt 480
attgtgaaga tagtggaaaa ggaaggtggc tcctacaaat gccatcattg cgataaagga 540
aaggccatcg ttgaagatgc ctctgccgac agtgggtcca aagatggacc cccaccacg 600
aggagcatcg tggaaaaaga agacgttcca accacgtctt caaagcaagt ggattgatgt 660
gatgggtcca tgtgagactt ttcaacaaag ggtaatatcc ggaaacctcc tcggattcca 720

```

-continued

```

ttgccagct atctgtcact ttattgtgaa gatagtggaa aaggaagggtg gctcctacaa 780
atgccatcat tgcgataaag gaaaggccat cgttgaagat gcctctgccg acagtgggtcc 840
caaagatgga cccccaccca cgaggagcat cgtggaaaaa gaagacgttc caaccacgtc 900
ttcaaagcaa gtggattgat gtgataattc tcacaatgaa caataataag attaaaaatag 960
ctttccccc ttgcagcgca tgggtathtt ttctagtaa aataaaagat aaacttagac 1020
tcaaaacatt tacaaaaaca acccctaaag ttcttaaagc ccaaagtgtc atccacgatc 1080
catagcaagc ccagcccaac ccaaccacac ccagcccacc ccagtccagc caactggaca 1140
atagtctcca cccccccca ctatcacctg gagttgtccg cacgcaccgc acgtctcgca 1200
gccaaaaaaa aaaagaaaga aaaaaaagaa aaagaaaaaa cagcaggtgg gtcgggtcg 1260
tgggggccgg aaacgcgagg aggatcgca gccagcgacg aggccggccc tccctccgct 1320
tccaaagaaa cgcccccat cgccactata tacatacccc cccctctcct cccatcccc 1380
caacctacc accaccacca ccaccacctc cacctctccc cccctcgctg cgggacgacg 1440
agctctccc cctccccct cgcgcgc 1467

```

```

<210> SEQ ID NO 23
<211> LENGTH: 1459
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Artificial promoter sequence

```

```

<400> SEQUENCE: 23

```

```

aagcttactc gaggtcattc atatgcttga gaagagagtc gggatagtcc aaaataaaac 60
aaaggaaga ttacctggtc aaaagtgaac acatcagtta aaagggtgta taaagtaaaa 120
tatcggtaat aaaaggtggc ccaaagtgaac atttactctt ttctactatt ataaaaattg 180
aggatgtttt tgcgggtact ttgatcgtc atttttgtat gaattggttt ttaagtttat 240
tcgcttttgg aaatgcatac ctgtatttga gtcgggtttt aagtctgttt gcttttgtaa 300
atacagaggg atttgtataa gaaatatctt tagaaaaacc catatgctaa tttgacataa 360
tttttgagaa aaatatatat tcaggcgaat tctcacaatg aacaataata agattaaaaat 420
agctttcccc cgttcgacgc catgggtatt ttttctagta aaaataaaag ataaacttag 480
actcaaaaca tttacaaaaa caaccctcaa agttcctaaa gcccaaagtg ctatccacga 540
tccatagcaa gccagccca acccaaccca acccagcca cccagtcga gccaaactgga 600
caatagtctc cacaccccc cactatcacc gtgagttgtc cgcacgcacc gcacgtctcg 660
cagccaaaaa aaaaaagaaa gaaaaaaaag aaaaaaagaa aacagcaggt ggggtccgggt 720
cgtgggggcc ggaacgcga ggagatcgc gagccagcga cgaggagctt ctgcaggtcc 780
gatgtgagac ttttcaaaa agggtaatat ccgaaacct cctcggattc cattgcccag 840
ctatctgtca ctttattgtg aagatagtgg aaaaggaagg tggctcctac aaatgccatc 900
attgcgataa aggaaaggcc atcgttgaag atgcctctgc cgacagtggc cccaaagatg 960
gacccccacc cagcaggagc atcgtggaaa aagaagacgt tccaaccacg tcttcaaagc 1020
aagtggattg atgtgatggt ccgatgtgag acttttcaac aaagggtaat atccgaaac 1080
ctcctcggat tcattgccc agctatctgt cactttattg tgaagatagt ggaaaaggaa 1140
gggtgctcct acaaatgcca tcattgcat aaaggaaagg ccacgttga agatgcctct 1200
gccgacagtg gtcccaaaag tggaccccca cccacgagga gcatcgtgga aaaagaagac 1260
gttccaacca cgtcttcaaa gcaagtggat tgatgtgatc cctccctccg cttccaagaa 1320

```

-continued

```

aacgcccc atcgccacta tatacatacc cccccctctc ctcccatccc cccaaacctc 1380
ccaccaccac caccaccacc tccacctctc cccccctcgc tgccggacga cgagctcctc 1440
ccccctcccc ctccgcgcg 1459

```

```

<210> SEQ ID NO 24
<211> LENGTH: 1939
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

```

```

<400> SEQUENCE: 24

```

```

taaatgacat cagatacacg cttgtgaacc atctttaaag tattgatgga ctcttcacta 60
tgaaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaatat 120
tagaggtcca aattcaaaaa aatgtogtat tgaatcattc cactactaaa ttggttcaat 180
gtcagattta aacagcctag agccttaggcc tcatcgttga agatgcctct gccgacagt 240
gtcccaaaga tggaccccc cccacgagga gcatcggtga aaaagaagac gttccaacca 300
cgtcttcaaa gcaagtggat tgatgtgata tctccactga cgttaaggat gacgcacaat 360
cccactatcc ttcgaggcct catcgttgaa gatgcctctg ccgacagtgg tcccaaagat 420
ggacccccac ccacgaggag catcggtgaa aaagaagacg ttccaaccac gtcttcaaag 480
caagtggatt gatgttatat ctccactgac gtaagggatg acgcacaatc cactatcct 540
tcgaagctct agggataatt tagtgagata tgagattcta ctttcaacat atactaatcc 600
taaatctcta gcaacttttt atataagcta taaatatcat gaaaatgtat tttaatcgtt 660
tcataattta tgcagtcaca ctaatggaag aaaggccaat tattattatt ttcttcagac 720
tataaatgaa aacataaatt aaaatgcaga ttagttttaa attttaataa gtaagtaaaa 780
tgcttatagc cttatacaaa atcatatttg gaagtttcta acattgttgc aatttgttat 840
cacaaatcac agtaatatat gtataactat tagtaattac aactatacac aaatttfaat 900
gggtaatcat atatttgtgt ccagtggatt gaacaaatat gctcggccca tgcggaagta 960
atgccaatth tgggtgagta aagcccctgc gaaatthtca cataagaaat gcatgcttht 1020
tgthttcaac gacatgagtt gcatgcttht taccattgct tatatagttg caagtttgca 1080
actccttgat atthttthta tgtagacact actaccacca aaaactthtg gtctgcttat 1140
tcttgthtac tatgtaaaaa aaataaatga attgthtatt tactccgatt tgatggagtc 1200
tggthtatga ggtthtatag cctthtacaga aaattgatag ttacaaaaat atthttcaaa 1260
aataaaaagg taaaaccgtc atthcaagtt gttattgtht tgggggactg gatttgaaat 1320
gaaatataga accggaaaac aaggtgagcc gaagtcaag cctthggacc cgtthttata 1380
thtactctc ccattccctt ctcttcaat ccttctctcc tctctctccc thtctctct 1440
tcccctctth cattthccag ccaactacaaa cthttctatc tctactthtt thctctcga 1500
thtcaggta cthttgagac cthttgtht gattthtgaa cacaccccc aattacgtht 1560
gattthtgat ccgcacatga thtcaattc tccgthctg agthtcttht ggatctgggt 1620
gtcttgagct aatctthtgc atctgthgt taccgattth actcatgcgt atgttcatta 1680
caccattgtt tathgttht atcaacaaa agactcatgt thttcaaatg thtttaatat 1740
aaththtctg attgaaatth ataaththt catgattctg gatccagaat atctctctc 1800
thcttcatt thgtctgta thgattgtht thtgaaaaag gattgthctt tgtatctgta 1860

```

-continued

```

ttggtgaaaa aggattgta tttgttgata aaaatttgat ctttaaaca tgtttggtt 1920
tgcataaagg tagaagacc 1939

<210> SEQ ID NO 25
<211> LENGTH: 1935
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 25
taaatgacat cagatacacg cttgtgaacc atctttaag tattgatgga ctcttcacta 60
tgaaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaattaat 120
tagaggtcca aattcaaaaa aatgtcgtat tgaatcattc cactactaaa ttggttcaat 180
gtcagattta aacagcctag ggataattta gtgagatag agattctact ttcaacatat 240
actaatccta aatctctagc aactttttat ataagctata aatatcatga aaatgtattt 300
taatcgtttc ataatttatg cagtcacact aatggaaaa aggccaatta ttattatttt 360
cttcagacta taaatgaaaa cataaattaa aatgcagatt agtttaaaat ttaataaagt 420
aagtaaaatg cttatagcct tatacaaaat catatttggg agtttctaac attggtgcaa 480
ttgtttatca caaatcacag taatatttgt aagcttaggc ctcacgttg aagatgcctc 540
tgccgacagt ggtcccaaa atggaccccc acccacgagg agcatcgtgg aaaaagaaga 600
cgttccaacc acgtcttcaa agcaagtggg ttgatgtgat atctccactg acgtaaggga 660
tgacgcacaa tcccactatc cttcgaggcc tcatcgttga agatgcctct gccgacagtg 720
gtcccaaaaga tggaccccc cccacgagga gcacgttggg aaaagaagac gttccaacca 780
cgtcttcaaa gcaagtggat tgatgttata tctccactga cgttaaggat gacgcacaat 840
cccactatcc ttcgaagctt actaattagt aattacaact atacacaaat ttaaatgggt 900
aatcatatat ttgtgtccag tggattgaa caaatatgct ggccccatgc gaagtaatgc 960
caattttggg tgagtaaaag ccatgogaaa ttttcacata agaaatgcat gctttttggt 1020
ttcaacgaca tgagttgcat gctttttatc attgcttata tagttgcaag ttgcaactc 1080
cttgatattt tttttatgta gacactacta ccacaaaaa cttttggtct gcttattctt 1140
gtttactatg taaaaaaaa aatgaattg tttatttact ccgatttgat ggagtctggt 1200
ttatgaggtt ttatagcctt tacagaaat tgatagttac aaaaatattt tcaaaaata 1260
aaagggtaaa accgtcattt caagtgtta ttgttttggg ggactggatt tgaatgaaa 1320
tatagaaccg gaaaacaagg tgagccgaag tcgaagcctt tggaccggt tttatattta 1380
ctcctcccat tcccttctcc ttcaatcctt ccttctctct cctccctctt tcttctccc 1440
ctctttcatt ttccagccac taaaaacttt tctatcteta ctttttttcc tctcgatttc 1500
aggtactttt tgagaccctt tgtgttgatt ttogaacaca caccccaatt acgtttgatt 1560
ttgatcccc catcgatttc aatcatccg tttctgagtt tcttttggat ctgggtgtct 1620
tgagctaate ttttcgatct gttgtttatc gattttactc atgcgatgt tcattacacc 1680
atgtgttatt tgtttaatca accaaaagac tcatgttttt caaatgtctt taatataatt 1740
ttctgattg aattttataa tatttacatg attctggatc cagaatatcc ttcttctct 1800
tccattttgt cctgtattga tttgtcttg aaaaaggatt gttctttgta tctgtattgg 1860
tgaaaaagga ttgttatttg ttgataaaaa tttgatcttt aaacaatggt tggttttgca 1920
taaaggtaga agacc 1935

```

-continued

```

<210> SEQ ID NO 26
<211> LENGTH: 1935
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 26
taaatgacat cagatacacg cttgtgaacc atctttaaag tattgatgga ctcttcacta    60
tgaaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaatat    120
tagaggcca aattcaaaaa aatgtcgtat tgaatcattc cactactaaa ttggttcaat    180
gtcagattta aacagcctag ggataattta gtgagatag agattctact ttcaacatat    240
actaatccta aatctctagc aactttttat ataagctata aatatcatga aaatgtattt    300
taatcgtttc ataatttatg cagtcacact aatggaaaaa aggccaatta ttattat    360
cttcagacta taaatgaaaa cataaattaa aatgcagatt agtttaaaat ttaataagt    420
aagtaaaatg cttatagcct tatacaaaat catatttggg agtttctaac attgttgcaa    480
ttgttatca caaatcacag taatatttgt aagcttcgaa ggatagtgga attgtgcgctc    540
atcccttacg tcagtggaga tataacatca atccacttgc tttgaagacg tggttggaac    600
gtctcttttt tccacgatgc tctcctgtggg tgggggtcca tctttgggac cactgtcggc    660
agaggcatct tcaacgatga ggcctcgaag gatagtgga ttgtgcgca tcccttacgt    720
cagtggagat atcacatcaa tccacttgc tgaagacgt ggttggaacg tcttcttttt    780
ccacgatgct cctcgtgggt ggggggtccat ctttgggacc actgtcggca gaggcattct    840
caacgatgag gcctaagcct actaattagt aattacaact atacacaaat ttaaatgggt    900
aatcatatat ttgtgtccag tggattgaac aaatatgctc ggcccatgcg gaagtaatgc    960
caattttggg tgagtaaagc ccatgogaaa ttttcacata agaaatgcat gctttttggt    1020
ttcaacgaca tgagttgcat gctttttatc attgcttata tagttgcaag ttgcaactc    1080
cttgatattt tttttatgta gacactacta ccaccaaaaa cttttggtct gcttattctt    1140
gtttactatg taaaaaaat aatgaattg tttatttact ccgatttgat ggagtctggt    1200
ttatgaggtt ttatagcctt tacagaaaat tgatagttac aaaaatattt tcaaaaata    1260
aaagggtaaa accgtcattt caagtgtta ttgttttggg ggactggatt tgaaatgaaa    1320
tatagaaccg gaaaacaagg tgagccgaag tcgaagcctt tggaccggt tttatattta    1380
ctctcccat tcccttctcc ttcaatcctt ccttctcct cctccctct tcttctccc    1440
ctcttctcatt ttccagccac taaaaacttt tctatcteta ctttttttcc tctcgatttc    1500
aggctacttt tgagaccctt tgttgtgatt ttogaacaca caccocaatt acgtttgatt    1560
tttgatcccg catcgatttc aatcatccg tttctgagtt tcttttggat ctgggtgtct    1620
tgagctaate ttttcgatct gttgtttatc gattttactc atgcgtatgt tcattacacc    1680
atttgttatt tgtttaatca accaaaagac tcatgttttt caaatgtctt taatataatt    1740
tttctgattg aattttataa tatttacatg attctggatc cagaatatcc ttcttctct    1800
tccattttgt cctgatttga tttgtctttg aaaaaggatt gttctttgta tctgatttg    1860
tgaaaaagga ttgttatttg ttgataaaaa tttgatcttt aaacaatggt tggttttgca    1920
taaaggtaga agacc    1935

```

<210> SEQ ID NO 27

-continued

```

<211> LENGTH: 1963
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 27
ggcgcctgc aggaagctgt accccccaag cttaaatgac atcagataca cgcttgatgaa    60
ccatctttaa agtattgatg gactcttcac tatgaaagct ctctttaaaa ttaattttct    120
ttgtacatgt ctctaagcaa tgtcaaatta attagaggtc caaattcaaa aaaatgtcgt    180
attgaaatcat tccattacta aattgggtca atgtcagatt taaacagcct agggataaatt    240
tagtgagata tgagattcta ctttcaacat atactaatcc taaatctcta gcaacttttt    300
atataagcta taaatatcat gaaaatgat tttaatcgtt tcataattta tgcagtcaca    360
ctaatggaaa aaaggccaat tattattatt ttcttcagac tataaatgaa aacataaatt    420
aaaaatgcaga ttagtttaaa attttaataa gtaagtaaaa tgcttatagc cttatacaaa    480
atcatatttg gaagtttcta acattgttgc aatttgttat cacaaatcac agtaaatatt    540
gtataactaat tagtaattac aactatacac aaatttaaat gggtaatcat atatttgtgt    600
ccagtgattg gaacaaatat gctcggccca tgcggaagta atgccaattt tgggtgagta    660
aagccatgac gaaattttca cataagaaaa gcttaggcct catcgttgaa gatgcctctg    720
ccgacagtgg tcccaaatgat ggacccccac ccacgaggag catcgttgaa aaagaagacg    780
ttccaaccac gtcttcaaa caagtggatt gatgtgatat ctccactgac gtaagggatg    840
acgcacaatc ccactatcct tcgaggcctc atcgttgaag atgcctctgc cgacagtggg    900
cccaaatgat gacccccacc cacgaggagc atcgttgaaa aagaagacgt tccaaccacc    960
tttcaaaage aagtggattg atgttatatc tccactgacg taagggatga cgcacaatcc   1020
cactatcctt cgaagcttgc tttttgtttt caacgacatg agttgcatgc tttttatcat   1080
tgcttatata gttgcaagtt tgcaactcct tgatattttt tttatgtaga cactactacc   1140
accaaaaact tttggtctgc ttattcttgt ttactatgta aaaaaataa atgaattggt   1200
tatttactcc gatttgatgg agtctggttt atgaggtttt atagccttta cagaaaattg   1260
atagttacaa aatatTTTTT caaaaataaa agggtaaaac cgtcatttca agttgttatt   1320
gttttggggg actggatttg aatgaaata tagaacggga aaacaagggtg agccgaagtc   1380
gaagcctttg gaccctttt tatatttact cctccattc ccttctcctt caatccttcc   1440
ttctctctcc tccctctctc ttcttccct ctttcatttt ccagccacta caaacttttc   1500
tatctctact tttttctctc tcgatttcag gtactttttg agaccctttg ttgtgatttt   1560
cgaacacaca cccaattac gtttgatttt tgatcccga tcgatttcaa ttcacccgtt   1620
tctgagtttc ttttgatct ggggtctctg agctaactct ttcgatctgt tgtttatcga   1680
ttttactcat gcgtatgttc attacacat ttgttatttg tttaatcaac caaaagactc   1740
atgtttttca aatgtcttta atataatTTT tctgattgaa ttttataata tttacatgat   1800
tctggatcca gaatatcctt cttctctctc cttttgtcc tgtattgatt tgtctttgaa   1860
aaaggattgt tctttgtatc tgatttggtg aaaaaggatt gttatttggg gataaaaaatt   1920
tgatctttaa acaatgtttg gttttgcata aaggtagaag acc                        1963

```

```

<210> SEQ ID NO 28
<211> LENGTH: 1963
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 28

```

ggcgcctgc aggaagctgt accccccaag cttaaatgac atcagataca cgcttgtaa    60
ccatctttaa agtattgatg gactcttcac tatgaaagct ctctttaaaa ttaattttct    120
ttgtacatgt ctctaagcaa tgtcaaatta attagaggtc caaattcaaa aaaatgtcgt    180
attgaatcat tccattacta aattggttca atgtcagatt taaacagcct agggataatt    240
tagtgagata tgagattcta ctttcaacat atactaatcc taaatctcta gcaacttttt    300
atataagcta taaatatcat gaaaatgat tttaatcgtt tcataattta tgcagtcaca    360
ctaattggaaa aaaggccaat tattattatt ttcttcagac tataaatgaa aacataaatt    420
aaaaatgcaga ttagtttaaa attttaataa gtaagtaaaa tgcttatagc cttatacaaa    480
atcatatttg gaagtttcta acattgttgc aatttgttat cacaaatcac agtaaatatt    540
gtataactaat tagtaattac aactatacac aaatttaaat gggtaatcat atatttggtg    600
ccagtggtatt gaacaaatat gctcggccca tgcggaagta atgccaattt tgggtgagta    660
aagcccatgc gaaattttca cataagaaaa gcttcgaagg atagtgggat tgtgcgtcat    720
cccttacgtc agtgagata taacatcaat ccacttgctt tgaagacgtg gttggaacgt    780
cttcttttcc cactgatgct ctcgtgggtg ggggtccatc tttgggacca ctgctggcag    840
aggcatcttc aacgatgagg cctcgaagga tagtgggatt gtgcgtcatc ccttacgtca    900
gtggagatat cacatcaatc cacttgcttt gaagacgtgg ttggaacgtc ttctttttcc    960
acgatgctcc tcgtgggtgg ggggtccatc ttgggaccac tgtcggcaga ggcactctca   1020
acgatgaggc ctaagcttgc tttttgtttt caacgacatg agttgcatgc tttttatcat   1080
tgcttatata gttgcaagtt tgcaactcct tgatattttt tttatgtaga cactactacc   1140
accaaaaact tttggtctgc ttattcttgt ttactatgta aaaaaataa atgaattggt   1200
tatttactcc gatttgatgg agtctgggtt atgaggtttt atagccttta cagaaaattg   1260
atagttacaa aatatTTTTT caaaaataaa agggtaaaac cgtcatttca agttgttatt   1320
gttttggggg actggatttg aatgaaata tagaacggga aaacaagggt agccgaagtc   1380
gaagcctttg gaccctttt tatatttact cctccattc ccttctcctt caatccttcc   1440
ttctctctcc tccctctctc ttcttccctt ctttcatttt ccagccacta caaacttttc   1500
tatctctact tttttctctc tcgatttcag gtactttttg agaccctttg ttgtgatttt   1560
cgaacacaca cccaattac gtttgatttt tgatcccga tcgatttcaa ttcacccgtt   1620
tctgagtttc ttttgatctc ggggtgcttg agctaactct ttcgatctgt tgtttatcga   1680
ttttactcat gcgatgttcc attacacatc ttgttatttg tttaatcaac caaaagactc   1740
atgtttttca aatgtcttta atataatttt tctgattgaa ttttataata tttacatgat   1800
tctggatcca gaatccctt cttctctctc cattttgtcc tgtattgatt tgtctttgaa   1860
aaaggattgt tctttgtatc tgtattgggt aaaaaggatt gttatttggg gataaaaatt   1920
tgatctttaa acaatgtttg gttttgcata aaggtagaag acc                          1963

```

<210> SEQ ID NO 29

<211> LENGTH: 1971

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

-continued

<400> SEQUENCE: 29

```

ggcgcctgc aggaagctgt acccccacag cttaaatgac atcagataca cgcttgtaa 60
ccatctttaa agtattgatg gactcttcac tatgaaagct ctctttaaaa ttaattttct 120
ttgtacatgt ctctaagcaa tgtcaaatta attagaggtc caaattcaaa aaaatgtcgt 180
attgaatcat tccattacta aattgggtca atgtcagatt taaacagcct agggataatt 240
tagtgagata tgagattcta ctttcaacat atactaatcc taaatctcta gcaacttttt 300
atataagcta taaatatcat gaaaatgat tttaatcgtt tcataattta tgcagtcaca 360
ctaattgaaa aaaggccaat tattattatt ttcttcagac tataaatgaa aacataaatt 420
aaaaatgcaga ttagtttaaa attttaataa gtaagtaaaa tgcttatagc cttatacaaa 480
atcatatttg gaagtttcta acattgttgc aatttggtat cacaaatcac agtaaatatt 540
gtataactaat tagtaattac aactatacac aaatttaaat gggtaatcat atatttgtgt 600
ccagtggatt gaacaaatat gctcggccca tgcggaagta atgccaattt tgggtgagta 660
aagcccatgc gaaattttca cataagaaat gcatgctttt tgttttcaac gacatgagtt 720
gcatgctttt tatcattgct tatatagttg caagtttgca actccttgat atttttttta 780
tgtagacact actaccacca aaaacttttg gtctgcttat tcttgtttac tatgtaaaaa 840
aaataaatga attgtttatt tactcogatt tgatggagtc tggtttatga ggttttatag 900
cctttacaga aaattgatag ttacaaaaat atttttcaaa aataaaaggg taaaaccgtc 960
atttcaagtt gttattgttt tgggggactg gatttgaat gaagcctagg cctcatcgtt 1020
gaagatgcct ctgccgacag tgggtccaaa gatggacccc caccacgag gagcatcgtg 1080
gaaaaagaag acgttccaac cacgtcttca aagcaagtgg attgatgtga tatctccact 1140
gacgtaaggg atgacgcaca atcccactat ccttcgaggc ctcacgttg aagatgcctc 1200
tgccgacagt ggtcccaag atggaccccc acccagagg agcatcgtgg aaaaagaaga 1260
cgttccaacc acgtcttcaa agcaagtgga ttgatgttat atctccactg acgtaaggga 1320
tgacgcacaa tcccactatc cttcgaagct atgaaatata gaaccggaaa acaagggtgag 1380
ccgaagtga agcctttgga cccgttttta tatttactcc tcccattccc ttctccttca 1440
atcttcctt cctcctcctc ccttcttctt ctccctctct ttcattttcc agccactaca 1500
aacttttcta tctctacttt ttttctctct gatttcaggt actttttgag accctttggt 1560
gtgattttcg aacacacacc ccaattacgt ttgatttttg atcccgcatc gatttcaatt 1620
catccgtttc tgagtttctt ttggatctgg gtgtcttgag ctaatctttt cgatctgttg 1680
tttatcgatt ttactcatgc gtatgttcat tacaccattt gttatttgtt taatcaacca 1740
aaagactcat gtttttcaaa tgtctttaat ataatttttc tgattgaatt ttataatatt 1800
tacatgattc tggatccaga atatccttct tcttcttcca tttgtctctg tattgatttg 1860
tctttgaaaa aggattgttc tttgtatctg tattgggtgaa aaaggattgt tatttgttga 1920
taaaaatttg atctttaaac aatgtttggt tttgcataaa ggtagaagac c 1971

```

<210> SEQ ID NO 30

<211> LENGTH: 1816

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 30

```

taaatgacat cagatacacg cttgtgaacc atctttaaag tattgatgga ctcttcaact 60

```

-continued

```

tgaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaattaat 120
tagagggtcca aattcaaaaa aatgtcgtat tgaatcattc cactactaaa ttgggttcaat 180
gtcagattta aacagcctag agcttctgac gtaagggatg acgcacctga cgtaagggat 240
gacgcacctg acgtaagggg tgacgcacct gacgtaaggg atgacgcact cgagatcccc 300
atctccactg acgtaagggg tgacgcacaa toccactatc cttcgcaaga cccttctctc 360
atataaggaa gttcattttca tttggagagg acacgctgac aagctagctt ggctgcaggt 420
agatcctagg gataatttag tgagatagta gattctactt tcaacatata ctaatcctaa 480
atctctagca actttttata taagctataa atatcatgaa aatgtatfff aatcgtttca 540
taatttatgc agtcacacta atggaaaaaa ggccaattat tattatfff ctcagactat 600
aaatgaaaac ataaatfata atgcagatta gtttaaaatt ttaataagta agtaaaatgc 660
ttatagcctt atacaaaatc atatttgaa gtttctaaca ttgttgcaat ttgttatcac 720
aaatcacagt aatatttgta tactaattag taattacaac tatacacaaa tttaaatggg 780
taatcatata tttgtgtcca gtggattgaa caaatatgct cggcccatgc ggaagtaatg 840
ccaatfffgg gtgagfataag cccatgcaaa atfffcatc aagaatgca tgctfffgt 900
ttcaacgac atgagttgca tgctfffat cattgcttat atagttgcaa gtttgcaact 960
ccttgatatt tfffatgt agacactact accacaaaa actfffggc tgettattc 1020
tgfffactat gtaaaaaaa faaatgaatt gfffatfacc tccgattga tggagtctgg 1080
fffatgaggt fffatagcct ffacagaaaa ttgatagfca caaaatatt fffcaaaat 1140
aaaagggfca aaccgctatt fcaagttgtt atgtfffgg gggactggat ttgaaatgaa 1200
atatagaacc ggaafcaag gtgagccgaa gfcgaagcct ttggaccctg fffatattt 1260
actcctccca ffccctctc cffcaatcct fcctcctcc fcctcctcc fctctctcc 1320
cctctffcat fffccagcca cfacaaactt fctctctct actfffctc cctcagatt 1380
caggtactff ttgagaccct ttgttgatg fffcgaaac acaccccaat facgtttgat 1440
fffatgaccc gcatcgatt caatfcctc gfffctgag fctfffggg fctgggtgfc 1500
ttgagctaat cfffctgac tgttfffat cgaatffact catgctgatg ffcattacac 1560
catttffat ttgtffaat aacfaaaaga fcctgffff fcaaatgct ffaatataat 1620
fffctgatt gaatffata atatffcat gattctggat ccagaatac cttctctcc 1680
ffccatfff fcctgtatg atffgtctt gaaaaaggat tgtctfffgt atctgtatg 1740
gtgaaaaagg atgtffatf gffgataaaa atffgatcct faaacatgt ttggtfffgc 1800
ataaaggtag aagacc 1816

```

<210> SEQ ID NO 31

<211> LENGTH: 1816

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 31

```

taaatgacat cagatacacg cttgtgaacc atctfataag tattgatgga ccttcacta 60
tgaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaattaat 120
tagagggtcca aattcaaaaa aatgtcgtat tgaatcattc cactactaaa ttgggttcaat 180
gtcagattta aacagcctag gatctacctg cagccaagct agcttgtcag cgtgctctc 240

```

-continued

```

ccaatgaaa tgaacttctt tatatagagg aagggtcttg cgaaggatag tgggattgtg 300
cgtcacccct tacgtcagtg gagatgggga tctcgagtgc gtcacccctt acgtcagggtg 360
cgtcacccct tacgtcaggt gcgcatccc ttacgtcagg tgcgcatccc cttacgtcag 420
aagctctagg gataatttag tgagatatga gattctactt tcaacatata ctaatcctaa 480
atctctagca actttttata taagtataa atatcatgaa aatgtathtt aatcgtttca 540
taatttatgc agtcacacta atggaaaaaa ggccaattat tattattttc ttcagactat 600
aaatgaaaac ataaattaaa atgcagatta gttttaaatt ttaataagta agtaaaatgc 660
ttatagcctt atacaaaatc atatttgaa gtttctaaca ttgttgcaat ttgttatcac 720
aaatcacagt aatatttgta tactaattag taattacaac tatacaciaa tttaaatggg 780
taatcatata tttgtgtcca gtggattgaa caaatatgct cggcccatgc ggaagtaatg 840
ccaattttgg gtgagtaaag cccatgcgaa attttcacat aagaaatgca tgctttttgt 900
tttcaacgac atgagttgca tgctttttat cattgcttat atagttgcaa gtttgcaact 960
ccttgatatt ttttttatgt agacactact accacaaaa acttttggtc tgcttattct 1020
tgtttactat gtaaaaaaaaa taaatgaatt gtttatttac tccgatttga tggagtctgg 1080
tttatgaggt tttatagcct ttacagaaaa ttgatagtta caaaaatatt tttcaaaaat 1140
aaaagggtaa aaccgtcatt tcaagttggt attgttttgg gggactggat ttgaaatgaa 1200
atatagaacc ggaaaacaag gtgagccgaa gtcgaagcct ttggaccctt ttttatattt 1260
actcctccca ttcccttctc ctccaatcct tccttctctc tcctcccttc ttcttcttcc 1320
cctctttcat tttccagcca ctacaaactt ttctatctct actttttttc ctctcgattt 1380
caggtacttt ttgagaccct ttgttgatgatt tttcgaacac acacccaat tacgtttgat 1440
ttttgatccc gcatcgattt caattcatcc gtttctgagt ttcttttggga tctgggtgtc 1500
ttgagctaat cttttcgatc tgttgtttat cgattttact catgcgtatg ttcattacac 1560
catttgttat ttgtttaatc aacccaaaaga ctcatgtttt tcaaatgtct ttaataataat 1620
ttttctgatt gaattttata atatttcat gattctggat ccagaatata cttcttcttc 1680
ttccattttg tcctgtattg atttgtcttt gaaaaggat tgttctttgt atctgtattg 1740
gtgaaaaagg attgttattt gttgataaaa atttgatctt taaacaatgt ttggttttgc 1800
ataaaggtag aagacc 1816

```

```

<210> SEQ ID NO 32
<211> LENGTH: 1812
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 32

```

```

taaatgacat cagatacacg cttgtgaacc atctttaaag tattgatgga ctcttcacta 60
tgaaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaattaat 120
tagagggtcca aattcaaaaa aatgctgat tgaatcattc cactactaaa ttggttcaat 180
gtcagattta aacagcctag ggataattta gtgagatag agattctact ttcaacatat 240
actaatccta aatctctagc aactttttat ataagctata aatatcatga aaatgtatht 300
taatcgtttc ataatttatg cagtcacact aatggaaaa aggccaatta ttattatttt 360
cttcagacta taaatgaaaa cataaattaa aatgcagatt agtttaaaat ttaataaagt 420
aagtaaatg cttatagcct tatacaaaat catatttggga agtttctaac attgttgcaa 480

```

-continued

```

ttgttatca caaatcacag taatatttgt aagcttctga cgtaagggat gacgcacctg 540
acgtaagga tgacgcacct gacgtaagg atgacgcacc tgacgtaagg gatgacgcac 600
tcgagatccc catctccact gacgtaagg atgacgcaca atcccactat ccttcgcaag 660
acccttcctc tatataagga agttcatttc atttgagag gacacgctga caagctagct 720
tggtgcagg tagatctact aattagtaat tacaactata cacaaattta aatgggtaat 780
catatatttg tgtccagtgg attgaacaaa tatgctcggc ccatgcgga gtaatgccaa 840
ttttgggtga gtaaagccca tgcgaaattt tcacataaga aatgcatgct ttttgttttc 900
aacgacatga gttgcatgct ttttatcatt gottatatag ttgcaagttt gcaactcctt 960
gatatttttt ttatgtagac actactacca ccaaaaactt ttggtctgct tattcttgtt 1020
tactatgtaa aaaaaataaa tgaattgttt atttactcgg atttgatgga gtctggttta 1080
tgaggtttta tagcctttac agaaaattga tagttacaaa aatatttttc aaaaaataaa 1140
gggtaaaacc gtcatttcaa gttgttattg ttttggggga ctggatttga aatgaaatat 1200
agaaccggaa aacaagggtga gccgaagtgc aagcctttgg acccgttttt atatttactc 1260
ctcccattcc ctctccttc aatccttctc tcctcctcct cccttcttct tcttccctc 1320
tttcattttc cagccactac aaacttttct atctctactt ttttctctct cgatttcagg 1380
tactttttga gaccctttgt tgtgattttc gaacacacac cccaattacg tttgattttt 1440
gatcccgcac cgatttcaat tcatcogttt ctgagtttct tttggatctg ggtgtcttga 1500
gctaactctt tcgatctggt gtttatcgat tttactcatg cgtatgttca ttacaccatt 1560
tgttatttgt ttaatcaacc aaaagactca tgtttttcaa atgtctttaa tataattttt 1620
ctgattgaat tttataatat ttacatgatt ctggatccag aatacccttc ttcttcttcc 1680
atttgtctct gtattgattt gtctttgaaa aaggattggt ctttgtatct gtattggtga 1740
aaaaggattg ttatttgttg ataaaaattt gatctttaa caatgtttgg ttttgcataa 1800
aggtagaaga cc 1812

```

<210> SEQ ID NO 33

<211> LENGTH: 1812

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 33

```

taaatgacat cagatacacg cttgtgaacc atctttaaag tattgatgga ctcttcaacta 60
tgaaagctct ctttaaaatt aattttcttt gtacatgtct ctaagcaatg tcaaattaat 120
tagagggtcca aattcaaaaa aatgtcgtat tgaatcattc cactactaaa ttggttcaat 180
gtcagattta aacagcctag ggataattta gtgagatag agattctact ttcaacatat 240
actaatccta aatctctagc aactttttat ataagctata aatatcatga aatgtattt 300
taatcgtttc ataatttatg cagtcacact aatggaaaa aggccaatta ttattatttt 360
cttcagacta taaatgaaaa cataaattaa aatgcagatt agtttaaaat ttaataaagt 420
aagtaaaatg cttatagcct tacaacaaa catatttggg agtttctaac attggtgcaa 480
ttgttatca caaatcacag taatatttgt agatctacct gcagccaagc tagcttgtca 540
gcgtgtcctc tccaaatgaa atgaacttcc ttatatagag gaagggtcct gcgaaggata 600
gtgggattgt gcgtcatccc ttacgtcagt ggagatgggg atctcagtg cgctcatcct 660

```

-continued

```

tacgtcaggt gcgtcatccc ttacgtcagg tgcgtcatcc cttacgtcag gtgcgtcacc 720
ccttacgtca gaagcttact aattagtaat tacaactata cacaaattta aatgggtaat 780
catatatattg tgtccagtggt attgaaacaaa tatgctcggc ccatgcggaa gtaatgccaa 840
ttttgggtga gtaaagccca tgcgaaattt tcacataaga aatgcatgct ttttgttttc 900
aacgacatga gttgcatgct ttttatcatt gcttatatag ttgcaagttt gcaactcctt 960
gatatttttt ttatgtagac actactacca ccaaaaactt ttggtctgct tattcttggt 1020
tactatgtaa aaaaaataaa tgaattgttt atttactccg atttgatgga gtctggttta 1080
tgaggtttta tagcctttac agaaaattga tagttacaaa aatatttttc aaaaaataaa 1140
gggtaaaacc gtcatttcaa gttgttattg ttttggggga ctggatttga aatgaaatat 1200
agaaccggaa aacaaggatga gccgaagtcg aagcctttgg acccgttttt atatttactc 1260
ctccattcc cttctccttc aatccttctc tctcctcctc cccttcttct tcttccctc 1320
tttcattttc cagccactac aaacttttct atctctactt ttttctctct cgatttcagg 1380
tactttttga gaccctttgt tgtgattttc gaacacacac cccaattacg tttgattttt 1440
gatcccgcac cgatttcaat tcatcogttt ctgagtttct tttggatctg ggtgtcttga 1500
gctaactctt tcgatctggt gtttatcgat tttactcatg cgtatgttca ttacaccatt 1560
tgttatttgt ttaatcaacc aaaagactca tgtttttcaa atgtctttaa tataattttt 1620
ctgattgaat tttataatat ttacatgatt ctggatccag aatacccttc ttcttcttcc 1680
atttgtctct gtattgattt gcttttgaaa aaggattggt ctttgtatct gtattggtga 1740
aaaaggattg ttatttgttg ataaaaattt gatctttaa caatgtttgg tttgcataa 1800
aggtagaaga cc 1812

```

<210> SEQ ID NO 34

<211> LENGTH: 1840

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 34

```

ggccgcctgc aggaagctgt acccccacag cttaaatgac atcagataca cgcttgtgaa 60
ccatctttaa agtattgatg gactcttcac tatgaaagct ctctttaaaa ttaattttct 120
ttgtacatgt ctctaagcaa tgtcaaatta attagaggtc caaattcaaa aaaatgctgt 180
attgaaatcat tccattacta aattgggtca atgtcagatt taaacagcct agggataatt 240
tagtgagata tgagatttca ctttcaacat atactaatcc taaatctcta gcaacttttt 300
atataagcta taaatatcat gaaaatgtat tttaatcgtt tcataattta tgcagtcaca 360
ctaattggaaa aaaggccaat tattattatt ttcttcagac tataaatgaa aacataaatt 420
aaaaatgcaga ttagttttaa attttaataa gtaagtaaaa tgcttatagc cttatacaaa 480
atcatatttg gaagtttcta acattgttgc aatttggtat cacaaatcac agtaaatatt 540
gtataactaat tagtaattac aactatacac aaatttaaat gggtaatcat atatttgtgt 600
ccagtggtatt gaacaaatat gctcggccca tgcggaagta atgccaattt tgggtgagta 660
aagcccatgc gaaattttca cataagaaaa gcttctgacg taagggatga cgcacctgac 720
gtaagggatg acgcacctga cgtaaggat gacgcacctg acgtaaggga tgacgcactc 780
gagatcccca tctccactga cgtaaggat gacgcacaat cccactatcc ttcgcaagac 840
ccttctctca tataaggaag ttcatttcat ttggagagga cacgctgaca agctagcttg 900

```

-continued

```

gctgcaggta gatctgcttt ttgttttcaa cgacatgagt tgcattgcttt ttatcattgc 960
ttatatagtt gcaagtttgc aactccttga ttttttttt atgtagacac tactaccacc 1020
aaaaactttt ggtctgctta ttcttggtta ctatgtaaaa aaaataaatg aattgtttat 1080
ttactccgat ttgatggagt ctggtttatg aggttttata gcctttacag aaaattgata 1140
gttacaaaaa tttttttcaa aaataaaagg gtaaaaccgt catttcaagt tgttattggt 1200
ttgggggact ggatttgaaa tgaaatatag aaccggaaaa caaggtagc cgaagtcgaa 1260
gcctttggac ccgtttttat atttactcct cccattccct tctcctcaa tcttccctc 1320
ctcctcctcc cttcttcttc ttcccctctt tcattttcca gccactacaa acttttctat 1380
ctctactttt tttcctctcg atttcaggta ctttttgaga ccctttgttg tgattttcga 1440
acacacaccc caattacggt tgatttttga tcccgcatcg atttcaattc atccgtttct 1500
gagtttcttt tggatctggg tgtcttgagc taatcttttc gatctgttgt ttatcgattt 1560
tactcatgcy tatgttcatt acaccatttg ttatttgttt aatcaaccaa aagactcatg 1620
tttttcaaat gtctttaata taatttttct gattgaattt tataaatattt acatgattct 1680
ggatccagaa tacccttctt cttcttccat tttgtcctgt attgatttgt cttgaaaaa 1740
ggattgttct ttgtatctgt attggtgaaa aaggattggt atttgttgat aaaaatttga 1800
tctttaaaca atgtttggtt ttgcataaag gtagaagacc 1840

```

<210> SEQ ID NO 35

<211> LENGTH: 1848

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic promoter sequence

<400> SEQUENCE: 35

```

ggcgcctgc aggaagctgt acccccacaag cttaaagtac atcagataca cgcttggtgaa 60
ccatctttaa agtattgatg gactcttcac tatgaaagct ctctttaaaa ttaattttct 120
ttgtacatgt ctctaagcaa tgtcaaatga attagaggtc caaattcaaa aaaatgtcgt 180
attgaaatcat tccattacta aattgggttca atgtcagatt taaacagcct agggataatt 240
tagtgagata tgagattcta ctttcaacat atactaatcc taaatctcta gcaacttttt 300
atataagcta taaatatcat gaaaatgtat tttaatcggt tcataattta tgcagtcaca 360
ctaatggaaa aaaggccaat tattattatt ttcttcagac tataaatgaa aacataaatt 420
aaaatgcaga ttagtttaaa attttaataa gtaagtaaaa tgcttatagc cttatacaaa 480
atcatatttg gaagtttcta acattgttgc aatttggtat cacaaatcac agtaaatatt 540
gtataactat tagtaattac aactatacac aaatttaaat gggtaatcat atatttgtgt 600
ccagtggatt gaacaaatat gctcggccca tgcggaagta atgccaattt tgggtgagta 660
aagcccatgc gaaattttca cataagaaat gcatgctttt tgttttcaac gacatgagtt 720
gcatgctttt tatcattgct tatatagttg caagtttgca actccttgat atttttttta 780
tgtagacact actaccacca aaaacttttg gtctgcttat tcttgtttac tatgtaaaaa 840
aaataaatga attgtttatt tactccgatt tgatggagtc tggtttatga ggttttatag 900
cctttacaga aaattgatag ttacaaaaat atttttcaaa aataaaaggg taaaaccgtc 960
atttcaagtt gttattgttt tgggggactg gatttgaat gaagcttctg acgtaagggg 1020
tgacgcacct gacgtaaggg atgacgcacc tgacgtaagg gatgacgcac ctgacgtaag 1080

```

-continued

ggatgacgca ctcgagatcc ccatctccac tgacgtaagg gatgacgcac aatcccacta	1140
tccttcgcaa gacccttcct ctatataagg aagttcattt catttgagaga ggacacgctg	1200
acaagctagc ttggctgcag gtagatcatg aaatatagaa cggaaaaaca aggtgagccg	1260
aagtgaage ctttggaacc gtttttatat ttaactcctcc cattcccttc tccttcaatc	1320
cttccttcct cctcctccct tctttctctt cccctcttcc attttccagc cactacaaac	1380
ttttctatct ctactttttt tctctctgat ttcaggtact ttttgagacc ctttggtgtg	1440
attttcgaac acacacccca attacgtttg atttttgatc ccgcacgat tcaattcat	1500
ccgtttctga gtttcttttg gatctgggtg tcttgagcta atcttttcca tctggtgttt	1560
atcgatttta ctcatgcgta tggtcattac accatttgtt atttgttta tcaacaaaa	1620
gactcatggt tttcaaatgt ctttaataata atttttctga tgaatttta taatatttac	1680
atgattctgg atccagaata tcttctctct tcttccattt tgcctctgat tgattgtct	1740
ttgaaaaagg attggtcttt gtatctgtat tggtgaaaaa ggattgttat ttggtgataa	1800
aaatttgatc tttaaacaat gtttggtttt gcataaaggc agaagacc	1848

We claim:

1. A chimeric promoter comprising a Cauliflower mosaic virus 35S promoter enhancer fused with a rice actin gene promoter to form said chimeric promoter, wherein said chimeric promoter enhancer is at least 99% identical to the 2x A1-B3 enhancer sequence of SEQ ID NO:5, and wherein the promoter enhancer is fused to the rice Act1 promoter at the -848 nt position of SEQ ID NO:7.
2. A construct comprising the chimeric promoter of claim 1, wherein said chimeric promoter is operably linked to a

- 25 transcribable polynucleotide molecule operably linked to a 3' transcription termination polynucleotide molecule.
3. The construct of claim 2, wherein said transcribable polynucleotide molecule is a gene of agronomic interest.
4. The construct of claim 3, wherein said gene of agronomic interest is a yield gene.
5. A transgenic plant stably transformed with the construct of claim 2.
6. A seed of the transgenic plant of claim 5, which comprises the construct of claim 2.

* * * * *