



US011198887B2

(12) **United States Patent**  
**Anderson et al.**

(10) **Patent No.:** **US 11,198,887 B2**

(45) **Date of Patent:** **Dec. 14, 2021**

(54) **CORN TRANSGENIC EVENT MON 95379 AND METHODS FOR DETECTION AND USES THEREOF**

(71) Applicant: **Monsanto Technology LLC**, St. Louis, MO (US)

(72) Inventors: **Heather M. Anderson**, Wildwood, MO (US); **Sarah L. Brown**, St. Louis, MO (US); **Renato A. Carvalho**, St. Louis, MO (US); **Ancideriton A. Castro**, St. Louis, MO (US); **Katherine M. Dunkmann**, St. Louis, MO (US); **Adam J. Evans**, St. Louis, MO (US); **Stanislaw Flasiniski**, Ballwin, MO (US); **Cara L. Griffith**, Catawissa, MO (US); **Tianxiang Shen**, St. Louis, MO (US); **Todd R. Smith**, St. Louis, MO (US); **Heidi M. Windler**, St. Louis, MO (US)

(73) Assignee: **Monsanto Technology LLC**, St. Louis, MO (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/525,278**

(22) Filed: **Jul. 29, 2019**

(65) **Prior Publication Data**  
US 2020/0032289 A1 Jan. 30, 2020

**Related U.S. Application Data**  
(60) Provisional application No. 62/711,810, filed on Jul. 30, 2018.

(51) **Int. Cl.**  
**C12N 15/82** (2006.01)  
**C07K 14/325** (2006.01)  
**C12N 15/10** (2006.01)

(52) **U.S. Cl.**  
CPC ..... **C12N 15/8286** (2013.01); **C07K 14/325** (2013.01); **C12N 15/102** (2013.01); **C12N 15/8274** (2013.01); **Y02A 40/146** (2018.01)

(58) **Field of Classification Search**  
None  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

6,593,273 B2	7/2003	Asrar et al.	
9,469,880 B2	10/2016	Adams et al.	
2016/0108426 A1*	4/2016	Baum .....	A01N 63/10 800/279
2016/0108428 A1*	4/2016	Baum .....	A01N 63/10 800/302

**OTHER PUBLICATIONS**

Invitation to Pay Additional Fees regarding PCT Application No. PCT/US2019/043666, dated Oct. 11, 2019, 3 pages.  
Genbank submission KG815221 [database online], Dec. 23, 2014, [retrieved on Nov. 12, 2019]. Retrieved from: <https://www.ncbi.nlm.nih.gov/nuccore/KG815221> full document.  
Genbank submission CP015659 [database online], May 23, 2016, [retrieved on Nov. 5, 2019]. Retrieved from: <https://www.ncbi.nlm.nih.gov/nuccore/LR584442> full document.  
Genbank submission F1982754 [database online], May 22, 2010, [retrieved on Nov. 12, 2019]. Retrieved from: <https://www.ncbi.nlm.nih.gov/nuccore/F1982754> full document.  
Genbank submission BZ318215 [database online], Nov. 6, 2002, [retrieved on Nov. 12, 2019]. Retrieved from: <https://www.ncbi.nlm.nih.gov/nuccore/BZ318215> full document.  
Genbank submission LN503569 [database online], Sep. 15, 2014, [retrieved on Nov. 12, 2019]. Retrieved from: <https://www.ncbi.nlm.nih.gov/nuccore/LN503569> full document.  
Genbank submission AC230033 [database online], Sep. 21, 2013, [retrieved on Nov. 12, 2019]. Retrieved from: <https://www.ncbi.nlm.nih.gov/nuccore/AC230033> full document.  
International Search Report and Written Opinion regarding PCT Application No. PCT/US2019/043666, dated Jan. 7, 2020, 12 pages.

\* cited by examiner

*Primary Examiner* — Anne Kubelik  
(74) *Attorney, Agent, or Firm* — Dentons US LLP;  
Timothy K. Ball, Esq.

(57) **ABSTRACT**

The invention provides a transgenic corn event MON 95379, plants, plant cells, seeds, plant parts, progeny plants, and commodity products comprising event MON 95379. The invention also provides polynucleotides specific for event MON 95379 and methods for using and detecting event MON 95379 as well as plants, plant cells, seeds, plant parts, progeny plants, and commodity products comprising event MON 95379.

**17 Claims, 3 Drawing Sheets**  
**Specification includes a Sequence Listing.**




Fig. 1




Fig. 2




Fig. 3

Continue self crosses as well as trait integration into other germplasms

1

## CORN TRANSGENIC EVENT MON 95379 AND METHODS FOR DETECTION AND USES THEREOF

### REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. provisional application No. 62/711,810, filed Jul. 30, 2018, which is herein incorporated by reference in its entirety.

### INCORPORATION OF SEQUENCE LISTING

The sequence listing contained in the file named MONS463US\_ST25 is 74.7 kilobytes (measured in Microsoft Windows®), was created on Jul. 25, 2019, is filed herewith by electronic submission, and is incorporated by reference.

### FIELD OF THE INVENTION

The invention relates to recombinant DNA molecules present in and/or isolated from corn event MON 95379. The invention also relates to transgenic corn plants, plant parts, and seeds, cells, and agricultural products containing corn event MON 95379, as well as methods of using the same and detecting the presence of corn event MON 95379. Transgenic corn plants, parts, seeds and cells containing corn event MON 95379 DNA exhibit resistance to insect infestations in the family Lepidoptera.

### BACKGROUND OF THE INVENTION

Corn (*Zea mays*) is an important crop and is a primary food source in many areas of the world. The methods of biotechnology have been applied to corn for improvement of the agronomic traits and quality of the product. One such agronomic trait is insect resistance, which is accomplished through the expression of heterologous insect toxins, also known as transgenes, inserted into the genome of the corn plant.

The expression of such transgenes in a transgenic plant, plant part, seed or cell may be influenced by many different factors, including the elements used in the cassettes driving transgene expression and the interaction of those elements within a cassette. This is complicated further for a transgenic insertion containing two or more expression cassettes, with each expression cassette having a transgene conferring a separate trait, also known as a multi-gene transgenic event. A commercially useful multi-gene transgenic event requires that each of the transgenes in the transgenic insertion express in the manner necessary for each trait. To achieve this, individual expression cassettes first are designed and tested in plants, and the expression cassettes that show the best insect activity, while not resulting in negative phenotypes due to expression, are selected for each trait. Next, the selected expression cassettes for one trait are combined with the selected expression cassettes for the other trait into a single construct. Multiple constructs are designed using different orientations to provide the best resistance and prevent the occurrence of negative phenotypes or negative agronomics, such as lower yield. The constructs are tested to ensure all the expression cassettes function well together and each transgene is properly expressed. Then, the selected combination and orientation of expression cassettes is used as a single transgenic insert to produce hundreds transgenic events, each event being the result of the random insertion of the construct in a different genomic location.

2

Each transgenic event is unique in its molecular profile and chromosomal insertion point. Because of the variability involved in event creation, each unique event must be analyzed through multiple generations of plants—in each step assessing the molecular characterization, protein expression efficacy, and agronomics—to select a superior event for commercial use. The performance of an event in a transgenic plant, plant part, seed or cell, and therefore its effectiveness, may be influenced by the genomic location of the transgene insertion. Specifically, the effectiveness of the event can be impacted by cis and/or trans factors relative to the integration site or chromatin structure. Events can have the same transgenic insertion and nonetheless have different transgene expression levels and performance across tissues and developmental stages, in various germplasm, or under specific growth conditions. There may also be undesirable phenotypic or agronomic differences between some events. Therefore, it is necessary to produce and analyze a large number of individual plant transformation events in order to select an event having superior properties relative to the desirable trait, and the optimal phenotypic and agricultural characteristics necessary to make it suitable for commercial purposes. Further, the creation of a multi-gene event for commercial use requires rigorous molecular characterization, greenhouse testing, and field trials over multiple years, in multiple locations, and under a variety of conditions so extensive agronomic, phenotypic, and molecular data may be collected. The resulting data must then be analyzed by scientists and agronomists to select an event that is useful for commercial purposes. Once selected, such an event may then be introgressed using plant breeding methods as a single locus having multiple insect resistance traits into new germplasm suitably adapted to specific local growing conditions, and stacked/combined by breeding with other different events conferring traits different from the traits conferred by the event of the present invention.

### SUMMARY OF THE INVENTION

The invention provides a novel transgenic corn event—MON 95379—that provides insecticidal control over Lepidopteran pests of corn. The invention also provides transgenic plants, plant cells, seeds, plant parts, and commodity products comprising event MON 95379. In another embodiment, the invention provides polynucleotides specific for event MON 95379 and plants, plant cells, seeds, plant parts, progeny plants, and commodity products comprising event MON 95379. In yet another embodiment, methods related to event MON 95379 are provided.

Thus, in one aspect the invention provides a recombinant DNA molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, and a complete complement thereof.

In one embodiment, the recombinant DNA molecule is derived from corn event MON 95379 in a sample of seed containing said event having been deposited as ATCC Accession No. PTA-125027.

Another aspect of the invention provides a DNA molecule comprising a polynucleotide segment of sufficient length to function as a DNA probe that hybridizes specifically under stringent hybridization conditions with corn event MON 95379 DNA in a sample, wherein detecting hybridization of said DNA molecule under said stringent hybridization conditions is diagnostic for the presence of corn event MON 95379 DNA in said sample. In certain embodiments, the

3

sample comprises a corn plant, corn plant cell, corn seed, corn plant part, corn progeny plant, processed corn seed, animal feed comprising corn, corn oil, corn meal, corn flour, corn flakes, corn bran, pasta made with corn, corn biomass, and fuel products produced using corn and corn parts.

Yet another aspect of the invention provides a pair of DNA molecules, comprising a first DNA molecule and a second DNA molecule different from the first DNA molecule, that function as DNA primers when used together in an amplification reaction with a sample containing corn event MON 95379 template DNA to produce an amplicon diagnostic for the presence of said corn event MON 95379 DNA in said sample, wherein said amplicon comprises the nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10.

Another embodiment of the invention is a method of detecting the presence of a DNA segment diagnostic for corn event MON 95379 DNA in a sample, said method comprising: a) contacting the sample with a DNA molecule that functions as a probe and hybridizes specifically under stringent conditions with corn event MON 95379; b) subjecting said sample and said DNA molecule to stringent hybridization conditions; and c) detecting hybridization of said DNA molecule to said DNA in said sample, wherein said detection is diagnostic for the presence of said corn event MON 95379 DNA in said sample.

Yet another embodiment of the invention is a method of detecting the presence of a DNA segment diagnostic for corn event MON 95379 DNA in a sample, the method comprising: a) contacting said sample with the pair of DNA molecules of the invention; b) performing an amplification reaction sufficient to produce a DNA amplicon; and c) detecting the presence of said DNA amplicon in said reaction, wherein said DNA amplicon comprises the nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10.

Another embodiment of the invention is a corn plant, corn plant part, corn cell, or part thereof comprising a recombinant polynucleotide molecule comprising the nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. This corn plant, corn plant part, corn cell, or part thereof is insecticidal when provided in the diet of a Lepidopteran insect pest. Lepidopteran pests can include Fall Armyworm (*Spodoptera frugiperda*), Corn Earworm (*Helicoverpa zea*), Southwestern Corn Borer (*Diatraea grandiosella*), Surgarcane Borer (*Diatraea saccharalis*), and Lesser Cornstalk Borer (*Elasmopalpus lignosellus*). In addition, the corn plant can be further defined as progeny of any generation of a corn plant comprising the corn event MON 95379.

Yet another embodiment of the invention is a method for protecting a corn plant from insect infestation, wherein said method comprises providing in the diet of a Lepidopteran insect pest an insecticidally effective amount of cells or tissue of a corn plant comprising corn event MON 95379. Again, contemplated Lepidopteran insect pests include Fall Armyworm (*Spodoptera frugiperda*), Corn Earworm (*Helicoverpa zea*), Southwestern Corn Borer (*Diatraea grandiosella*), Surgarcane Borer (*Diatraea saccharalis*), and Lesser Cornstalk Borer (*Elasmopalpus lignosellus*).

4

Another embodiment of the invention is a method of producing an insect resistant corn plant comprising: a) sexually crossing two different corn plants with at least one of the two different corn plants comprising transgenic corn event MON 95379 DNA; b) sampling seed or tissue from the progeny of step (a); c) detecting in said sample from step (b) progeny comprising corn event MON 95379 DNA; and d) selecting said progeny comprising corn event MON 95379 DNA.

A further embodiment of the invention is a corn seed, nonliving corn plant material, or a microorganism comprising a detectable amount of the nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10, or complete complements thereof.

Yet another embodiment of the invention is a commodity product comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10, or complete complements thereof. Contemplated commodity products include whole or processed corn seed, animal feed comprising corn, corn oil, corn meal, corn flour, corn flakes, corn bran, corn biomass, and fuel products produced using corn and corn parts.

Another embodiment of the invention is a corn plant, corn plant part, or corn seed thereof comprising DNA functional as a template when tested in DNA amplification method producing an amplicon diagnostic for the presence of event MON 95379 DNA.

Yet another embodiment of the invention is a method of determining the zygosity of a corn plant or corn seed comprising event MON 95379 comprising: a) contacting a sample comprising corn DNA with a primer pair that is capable of producing an amplicon of one of the toxin coding sequences encoding Cry1B.868 or Cry1Da<sub>7</sub>; b) contacting said sample comprising corn DNA with a primer pair that is capable of producing an amplicon of an internal standard known to be single-copy and homozygous in the corn plant; c) contacting the DNA sample with a probe set which contains at least a first probe that specifically hybridizes to one of the toxin coding sequences encoding Cry1B.868 or Cry1Da<sub>7</sub>, and a second probe that specifically hybridizes to the internal standard genomic DNA known to be single-copy and homozygous in the corn plant; d) performing a DNA amplification reaction using real-time PCR and determining the cycle thresholds (Ct values) of the amplicon corresponding to the toxin coding sequence and the single-copy, homozygous internal standard; e) calculating the difference ( $\Delta$ Ct) between the Ct value of the single-copy, homozygous internal standard amplicon and the Ct value of the toxin coding sequence amplicon; and f) determining zygosity, wherein a  $\Delta$ Ct of about zero (0) indicates homozygosity of the inserted T-DNA of event MON 95379 and a  $\Delta$ Ct of about one (1) indicates heterozygosity of the inserted T-DNA of event MON 95379. In certain embodiments of this method, the primer pairs are selected from the group consisting of SEQ ID NO:18 combined with SEQ ID NO:19, and SEQ ID NO:21 combined with SEQ ID NO:22; and the probes are SEQ ID NO:20 and SEQ ID NO:23. In another embodiment, the primer pairs are selected from the group consisting of SEQ ID NO:18 combined with SEQ ID NO:19, and SEQ ID NO:24 combined with SEQ ID NO:25; and the probes are SEQ ID NO:20 and SEQ ID NO:26. In yet another embodiment of this invention the  $\Delta$ Ct of about one (1) indicating

heterozygosity of the inserted T-DNA of event MON 95379 is in the range of 0.75 to 1.25.

A further embodiment of the invention is a method of determining the zygosity of a corn plant or corn seed comprising event MON 95379 comprising: a) contacting a sample comprising corn DNA with a set of primer pairs comprising at least two different primer pairs capable of producing a first amplicon diagnostic for event MON 95379 and a second amplicon diagnostic for native corn genomic DNA not comprising event MON 95379; i) performing a nucleic acid amplification reaction with the sample and the set of primer pairs; ii) detecting in the nucleic acid amplification reaction the first amplicon diagnostic for event MON 95379, or the second amplicon diagnostic for native corn genomic DNA not comprising event MON 95379, wherein the presence of only the first amplicon is diagnostic of a corn plant or corn seed homozygous for event MON 95379, and the presence of both the first amplicon and the second amplicon is diagnostic of a corn plant or corn seed heterozygous for event MON 95379; orb) contacting a sample comprising corn DNA with a probe set which contains at least a first probe that specifically hybridizes to event MON 95379 DNA and at least a second probe that specifically hybridizes to corn genomic DNA that was disrupted by insertion of the heterologous DNA of event MON 95379 and does not hybridize to event MON 95379 DNA; i) hybridizing the probe set with the sample under stringent hybridization conditions, wherein detecting hybridization of only the first probe under the hybridization conditions is diagnostic for a corn plant or corn seed homozygous for event MON 95379, and wherein detecting hybridization of both the first probe and the second probe under the hybridization conditions is diagnostic for a corn plant or a corn seed heterozygous for event MON 95379. In one embodiment of this method, the set of primer pairs comprises SEQ ID NO:15 combined with SEQ ID NO:16, and SEQ ID NO:15 combined with SEQ ID NO:27. In another embodiment of this method, the probe set comprises SEQ ID NO:17 and SEQ ID NO:28.

The foregoing and other aspects of the invention will become more apparent from the following detailed description.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 represents the sequence of corn event MON 95379. Horizontal lines and boxes correspond to the positions of SEQ ID NO:1 ([1]), SEQ ID NO:2 ([2]), SEQ ID NO:3 ([3]), SEQ ID NO:4 ([4]), SEQ ID NO:5 ([5]), SEQ ID NO:6 ([6]), SEQ ID NO:7 ([7]), SEQ ID NO:8 ([8]), SEQ ID NO:9 ([9]), SEQ ID NO:11 ([11]), and SEQ ID NO:12 ([12]) relative to SEQ ID NO:10 ([10]). The horizontal arrows labeled SQ51219 (SEQ ID NO:15) ([15]), SQ21524 (SEQ ID NO:16) ([16]), SQ50998 (SEQ ID NO:21) ([21]), SQ50997 (SEQ ID NO:22) ([22]), SQ50485 (SEQ ID NO:24) ([24]) and SQ50484 (SEQ ID NO:25) ([25]) represent the approximate position of subsets of primers that can be used to detect corn event MON 95379. The horizontal arrows labeled PB10269 (SEQ ID NO:17) ([17]), PB50340 (SEQ ID NO:23) ([23]), PB50138 (SEQ ID NO:26) ([26]) represent the approximate position of a DNA probe that can be used to detect corn event MON 95379. “E” represents an enhancer element, “P” represents a promoter element, “L” represents a leader (5' UTR) element, “I” represents an intron element, “T” represents a 3' UTR, “Cry1B.868” represents the Cry1B.868 coding sequence element, “Cry1Da\_7” represents the Cry1Da\_7 coding sequence ele-

ment, “LoxP” represents the site at which Cre-recombinase marker excision occurred, leaving behind one of the two LoxP sites after marker excision, and “LB” represents the left T-DNA border.

FIG. 2 is a diagrammatic representation of the T-DNA cassettes before integration to form event MON 95379, after integration, and after Cre-excision. The top horizontal box represents the T-DNA cassette in the plasmid vector used to transform event MON 95379, presented as SEQ ID NO:13 ([13]) (“T-DNA Before Integration”). The horizontal arrows below [13] represent the individual genetic elements comprised within the two transgene cassettes. “LB” represents a T-DNA left border element, “E” represents an enhancer element, “P” represents a promoter element, “L” represents a leader (5' UTR) element, “I” represents an intron element, “T” represents a 3' UTR, “Cry1B.868” represents the Cry1B.868 coding sequence element, “Cry1Da\_7” represents the Cry1Da\_7 coding sequence element, “CP4” represents the CP4 selectable marker, “TS” represented a targeting sequence, “LoxP” represents the site at which Cre-recombinase marker excision occurs, and “RB” represents a T-DNA right border element. The middle horizontal box, “Inserted T-DNA After Integration,” represents the T-DNA cassette integrated into the corn genome after transformation wherein the right T-DNA border (RB) was lost during integration. The bottom horizontal box, “Inserted T-DNA After Cre-Excision,” represents the integrated T-DNA cassette after the CP4 selectable marker cassette was excised, leaving behind one of the two LoxP sites and the LB region.

FIG. 3 is a diagrammatic representation of the breeding process to produce the marker-free corn event MON 95379. R<sub>0</sub> generation events (“transformants”) are those that are derived from the initial transformation with the binary transformation vector used to generate corn event MON 95379. Subsequent “R” generations (R<sub>1</sub>, and R<sub>2</sub>) represent successive generations produced through self-pollination of plants derived from the initial R<sub>0</sub> transformant that resulted in the corn event MON 95379. The R<sub>2</sub> transformants which are homozygous for the T-DNA insertion are cross-pollinated with an elite transgenic corn line comprising a transgene cassette for the expression of Cre-recombinase, resulting in an F<sub>1</sub> generation, wherein many of the progeny have lost the CP4 selectable marker cassette due to Cre-recombinase excision. Hemizygous T-DNA positive, CP4 negative plants are selected and self-pollinated, resulting in an F<sub>2</sub> generation. F<sub>2</sub> plants homozygous for the inserted T-DNA allele without the CP4 marker and lacking the Cre-recombinase transgene cassette are selected and self-pollinated giving rise to an F<sub>3</sub> generation. The F<sub>3</sub> generation plants are self-pollinated giving rise to a pure line of F<sub>4</sub> Gold Standard Seed.

#### BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO:1 is a 50 nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:1 is found within SEQ ID NO:10 at nucleotide positions 838-887.

SEQ ID NO:2 is a 50 nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA. SEQ ID NO:2 is found within SEQ ID NO:10 at nucleotide positions 14,156-14,205.

SEQ ID NO:3 is a 100 nucleotide sequence representing the 5' junction region of corn genomic DNA and the inte-

grated transgenic expression cassette. SEQ ID NO:3 is found within SEQ ID NO:10 at nucleotide positions 813-912.

SEQ ID NO:4 is a 100 nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA. SEQ ID NO:4 is found within SEQ ID NO:10 at nucleotide positions 14,131-14,230.

SEQ ID NO:5 is a 200 nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:5 is found within SEQ ID NO:10 at nucleotide positions 763-962.

SEQ ID NO:6 is a 200 nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA. SEQ ID NO:6 is found within SEQ ID NO:10 at nucleotide positions 14,081-14,280.

SEQ ID NO:7 is a 1,160 nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:7 is found within SEQ ID NO:10 at nucleotide positions 1-1,160.

SEQ ID NO:8 is a 1,178 nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA. SEQ ID NO:8 is found within SEQ ID NO:10 at nucleotide positions 14,039-15,216.

SEQ ID NO:9 is a 13,318 nucleotide sequence corresponding to the transgenic inserted T-DNA of corn event MON 95379.

SEQ ID NO:10 is a 15,216 nucleotide sequence corresponding to the contig nucleotide sequence of the 5' genomic flanking DNA nucleotide sequence, the inserted T-DNA nucleotide sequence in event MON 95379, and the 3' genomic flanking DNA nucleotide sequence; and includes SEQ ID NO:11 (nucleotides 1-862), SEQ ID NO:9 (nucleotides 863-14,180), and SEQ ID NO:12 (nucleotides 14,181-15,216).

SEQ ID NO:11 is an 862 nucleotide sequence representing the 5' flanking corn genomic DNA up to the inserted T-DNA. SEQ ID NO:11 is found within SEQ ID NO:10 at nucleotide positions 1-862.

SEQ ID NO:12 is a 1,036 nucleotide sequence representing the 3' flanking corn genomic DNA after the inserted T-DNA. SEQ ID NO:12 is found within SEQ ID NO:10 at nucleotide positions 14,181-15,216.

SEQ ID NO:13 is a 18,376 nucleotide sequence representing the transgene cassette comprised within the binary plasmid transformation vector used to transform corn to produce corn event MON 95379.

SEQ ID NO:14 is a 35 nucleotide sequence representing the LoxP sites used for Cre-mediated excision and recombination. A remaining LoxP site after marker excision can be found within SEQ ID NO:10 at nucleotide positions 1,080-1,114.

SEQ ID NO:15 is a 20 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ51219 used to identify corn event MON 95379 DNA in a sample, and is identical to the nucleotide sequence corresponding to positions 833-852 of SEQ ID NO:10.

SEQ ID NO:16 is a 30 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ21524 used to identify corn event MON 95379 DNA in a sample, and is identical to the reverse complement of the nucleotide sequence corresponding to positions 905-934 of SEQ ID NO:10.

SEQ ID NO:17 is a 16 nucleotide sequence corresponding to a probe referred to as PB10269 used to identify corn event MON 95379 DNA in a sample, and is identical to the reverse complement of the nucleotide sequence corresponding to positions 886-901 of SEQ ID NO:10.

SEQ ID NO:18 is a 24 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ20222 used as an internal control for the event and zygosity assay for MON 95379 and hybridizes to a region of the corn genome.

SEQ ID NO:19 is a 28 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ20221 used as an internal control for the event and zygosity assay for MON 95379 and hybridizes to a region of the corn genome.

SEQ ID NO:20 is a 29 nucleotide sequence corresponding to a probe referred to as PB50237 used as an internal control for the event and zygosity assay for MON 95379 and hybridizes to a region of the corn genome.

SEQ ID NO:21 is a 20 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50998 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1B.868 within SEQ ID NO:10; and is identical to the nucleotide sequence corresponding to positions 2,809-2,828 of SEQ ID NO:10.

SEQ ID NO:22 is a 20 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50997 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1B.868 within SEQ ID NO:10; and is identical to the reverse complement of the nucleotide sequence corresponding to positions 2,852-2,871 of SEQ ID NO:10.

SEQ ID NO:23 is an 18 nucleotide sequence corresponding to a probe referred to as PB50340 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1B.868 within SEQ ID NO:10; and is identical to the reverse complement of the nucleotide sequence corresponding to positions 2,833-2,850 of SEQ ID NO:10.

SEQ ID NO:24 is a 19 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50485 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1Da\_7 within SEQ ID NO:10; and is identical to the nucleotide sequence corresponding to positions 12,820-12,838 of SEQ ID NO:10.

SEQ ID NO:25 is an 18 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50484 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1Da\_7 within SEQ ID NO:10; and is identical to the reverse complement of the nucleotide sequence corresponding to positions 12,855-12,872 of SEQ ID NO:10.

SEQ ID NO:26 is a 14 nucleotide sequence corresponding to a probe referred to as PB50138 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1Da\_7 within SEQ ID NO:10; and is identical to the reverse complement of the nucleotide sequence corresponding to positions 12,840-12,853 of SEQ ID NO:10.

SEQ ID NO:27 is a 21 nucleotide sequence corresponding to a thermal amplification primer referred to as PNEG DNA used in the zygosity assay for event MON 95379 and hybridizes to a region of corn genomic DNA which was deleted when the T-DNA used to produce event MON 95379 inserted into the corn genome. An amplicon derived from the

combination of primers SQ51219 and PNEG DNA is diagnostic for the wild-type allele lacking the event MON 95379 inserted T-DNA.

SEQ ID NO:28 is a 14 nucleotide sequence corresponding to a probe referred to as PRBNEG DNA used in the zygosity assay for event MON 95379 and hybridizes to a region of corn genomic DNA which was deleted when the T-DNA used to produce event MON 95379 inserted into the corn genome.

#### DETAILED DESCRIPTION

The present invention provides a transgenic corn event—MON 95379—that achieves insecticidal control over Lepidopteran pests of corn by expression of Cry1B.868 and Cry1Da\_7. Specifically, expression of the Cry1B.868 and Cry1Da\_7 insect inhibitory proteins in corn event MON 95379 provides resistance to the Lepidopteran insect pests Fall Armyworm (*Spodoptera frugiperda*), Corn Earworm (*Helicoverpa zea*), Southwestern Corn Borer (*Diatraea grandiosella*), Surgarcane Borer (*Diatraea saccharalis*), and Lesser Cornstalk Borer (*Elasmopalpus lignosellus*). Event MON 95379 will meet a great need for control of these insects in the corn market, as chemical insecticides often do not provide adequate control of these insects, or require multiple applications over the growing season, increasing the input of chemical pesticides in the environment and adding cost to the production of corn.

It should be understood that reference to event MON 95379 is equivalent to reference to event MON95379; they are interchangeable and represent the same transgenic corn event.

Plant transformation techniques are used to insert foreign DNA (also known as transgenic DNA) randomly into a chromosome of the genome of a cell to produce a genetically engineered cell, also referred to as a “transgenic” or “recombinant” cell. Using this technique, many individual cells are transformed, each resulting in a unique “transgenic event” or “event” due to the random insertion of the foreign DNA into the genome. A transgenic plant is then regenerated from each individual transgenic cell. This results in every cell of the transgenic plant containing the uniquely inserted transgenic event as a stable part of its genome. This transgenic plant can then be used to produce progeny plants, each containing the unique transgenic event.

Corn event MON 95379 was produced by an *Agrobacterium*-mediated transformation process of corn immature embryos with a single T-DNA binary system. In this system, an *Agrobacterium* strain employing one binary plasmid vector with a single T-DNA was utilized. The T-DNA construct comprised two transgene cassettes for the expression of the insect toxin coding sequences encoding Cry1B 0.868 and Cry1Da\_7, and a transgene cassette used for the selection of transformed corn cells using glyphosate selection (CP4). The T-DNA construct is SEQ ID NO:13, and illustrated in FIG. 2 (“T-DNA Before Integration”). During integration, the right T-DNA border was lost as shown in FIG. 2 (“Inserted T-DNA After Integration”). The glyphosate selection cassette was flanked on both sides with LoxP recognition sites which are recognized by Cre-recombinase, derived from Enterobacteria phage P1 (Larry Gilbertson (2003) *Cre-lox recombination: Cre-active tools for plant biotechnology*. *TRENDS in Biotechnology*, 21:12, 550-555).

As specifically described herein, corn event MON 95379 was produced by a complex research and development process in which: (1) hundreds of plasmid vector constructs—which varied with respect to the coding sequences

for the insecticidal proteins, the coding sequences for the transcriptional regulatory elements, and number and orientation of the cassettes within the constructs—were developed and transformed into corn cells to create thousands of events that were tested and analyzed, resulting in the selection of the construct used to generate event MON 95379; (2) hundreds of corn cells were transformed with the construct used to generate event MON 95379, creating a population of transgenic plants in which each plant contained a unique transgenic event that was regenerated and tested; (3) the final event MON 95379 was selected after a rigorous multi-year event selection process involving the testing and analysis of molecular characteristics, efficacy, protein expression, and agronomic properties in a variety of genetic backgrounds; and (4) the glyphosate selection cassette in corn event MON 95379 was removed through *in vivo* Cre-excision to create a “marker-free” final event MON 95379. Corn event MON 95379 was thus produced and selected as a uniquely superior event useful for broad-scale agronomic purposes.

The plasmid DNA inserted into the genome of corn event MON 95379 was characterized by detailed molecular analysis. This analysis included: the insert number (number of integration sites within the corn genome), the genomic insert location (the specific site in the corn genome where the insertion occurred), the copy number (the number of copies of the T-DNA within one locus), and the integrity of the transgenic inserted DNA. The detailed molecular analysis demonstrated that the integrated T-DNA containing the Cry1B.868 and Cry1Da\_7 expression cassettes remained intact after integration and Cre-excision of the glyphosate (CP4) selection cassette. As used herein, an “expression cassette” or “cassette” is a recombinant DNA molecule comprising a combination of distinct elements that are to be expressed by a transformed cell. Table 1 provides a list of the elements contained in SEQ ID NO:10, the DNA sequence that corresponds to corn event MON 95379.

TABLE 1

Description of corn event MON 95379		
Element	Position in SEQ ID NO: 10	Description
5' Flanking DNA	1-862	DNA sequence flanking the 5' end of the transgenic insert.
Left Border Region	863-1044	DNA region from <i>Agrobacterium tumefaciens</i> containing the left border sequence.
LoxP	1080-1114	A recognition sequence for a site-specific recombinase from <i>Enterobacteria phage P1</i> .
T-Os.LTP:1	1225-1524	The 3' untranslated region for a Lipid Transfer Protein-like gene (LTP) from <i>Oryza sativa</i> (rice).
Cry1B.868	1534-5133	Coding sequence of a chimeric insect toxin comprised of domains 1 and 2 of Cry1Be2, domain 3 of Cry1Ca, and the protoxin domain of Cry1Ab3.
I-Zm.UbqM1-1:1:16	5160-6212	An intron derived from an Ubiquitin 1 gene of <i>Zea mays Mexicana</i> .
L-Zm.UbqM1-1:1:4	6213-6290	A 5' UTR derived from an Ubiquitin 1 gene of <i>Zea mays Mexicana</i> .
P-Zm.UbqM1-1:1:5	6291-7167	A promoter derived from an Ubiquitin 1 gene of <i>Zea mays Mexicana</i> .
E-FMV.35S-1:1:2	7195-7731	The enhancer of the 35S gene from Figwort Mosaic Virus (FMV).
P-SETit.Tip-1:1:1	7743-8659	A promoter of a tonoplast membrane integral protein gene from <i>Setaria italica</i> (foxtail millet).

TABLE 1-continued

Description of corn event MON 95379		
Element	Position in SEQ ID NO: 10	Description
L-SETit.Tip-1:1:1	8660-8723	A 5' UTR of a tonoplast membrane integral protein gene from <i>Setaria italica</i> (foxtail millet).
I-Os.Act15-1:1:1	8732-10024	The first intron from the Actin 15 (Act 15) gene from <i>Oryza sativa</i> (rice).
Cry1Da_7	10043-13543	Coding sequence of a Cry1Da insect toxin with amino acid modifications to improve efficacy.
T-Os.GOS2-1:1:1	13560-14027	The 3' untranslated region from the GOS2 gene encoding a translation initiation factor from <i>Oryza sativa</i> (rice).
3' Flanking Sequence	14181-15216	DNA sequence flanking the 3' end of the transgenic insert.

Corn event MON 95379 is characterized as an insertion into a single locus in the corn genome, resulting in two new loci or junction sequences (e.g., sequences set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8) spanning a portion of the inserted DNA and the corn genomic DNA that are not known to appear naturally in the corn genome or other transgenic corn events—they are unique to event MON 95379. These junction sequences are useful in detecting the presence of the event MON 95379 in corn cells, corn tissue, corn seed, and corn plants or corn plant products, such as corn commodity products. DNA molecular probes and primer pairs are described herein that have been developed for use in identifying the presence of these various junction segments in biological samples containing or suspected of containing corn cells, corn seed, corn plant parts, or corn plant tissue that contain the event MON 95379.

A sample is intended to refer to a composition that is either substantially pure corn DNA or a composition that contains corn DNA. In either case, the sample is a biological sample, i.e., it contains biological materials, including but not limited to DNA obtained or derived from, either directly or indirectly, the genome of corn event MON 95379. “Directly” refers to the ability of the skilled artisan to directly obtain DNA from the corn genome by fracturing corn cells (or by obtaining samples of corn that contain fractured corn cells) and exposing the genomic DNA for the purposes of detection. “Indirectly” refers to the ability of the skilled artisan to obtain the target or specific reference DNA, i.e., a novel and unique junction segment described herein as being diagnostic for the presence of the event MON 95379 in a particular sample, by means other than by obtaining directly via fracturing of corn cells or obtaining a sample of corn that contains fractured corn cells. Such indirect means include, but are not limited to, amplification of a DNA segment that contains the DNA sequence targeted by a particular probe designed to bind with specificity to the target sequence, or amplification of a DNA segment that can be measured and characterized, i.e., measured by separation from other segments of DNA through some efficient matrix such as an agarose or acrylamide gel or the like, or characterized by direct sequence analysis of the amplicons, or cloning of the amplicon into a vector and direct sequencing of the inserted amplicon present within such vector.

Detailed molecular analysis demonstrated that event MON 95379 contains a single T-DNA insertion with one

copy of each of the Cry1B.868 and Cry1Da\_7 expression cassettes. No additional elements from the transformation construct other than portions of the *Agrobacterium tumefaciens* left border region used for transgenic DNA transfer from the plant transformation plasmid to the corn genome were identified in event MON 95379. Finally, thermal amplification producing specific amplicons diagnostic for the presence of event MON 95379 in a sample and DNA sequence analyses were performed to determine the arbitrarily assigned 5' and 3' insert-to-plant genome junctions, confirm the organization of the elements within the insert, and determine the complete DNA sequence of the inserted transgenic DNA (SEQ ID NO:9). SEQ ID NO:11 is a sequence representing the eight hundred sixty-two (862) base-pair (bp) 5' LH244 corn genomic DNA sequence flanking the inserted T-DNA sequence presented as SEQ ID NO:9. SEQ ID NO:12 is a sequence representing the one thousand thirty-six (1,036) bp 3' LH244 corn genomic DNA sequence flanking the inserted T-DNA sequence presented as SEQ ID NO:9. SEQ ID NO:7 is a sequence representing the eight hundred sixty-two (862) base-pair (bp) 5' LH244 corn genomic DNA sequence flanking the inserted T-DNA sequence combined with two hundred ninety-eight (298) bp of inserted T-DNA sequence presented as SEQ ID NO:9. SEQ ID NO:8 is a sequence representing one hundred forty-two (142) bp of inserted T-DNA sequence with the one thousand thirty-six (1,036) bp 3' LH244 corn genomic DNA sequence flanking the inserted T-DNA sequence presented as SEQ ID NO:9. SEQ ID NO:10 corresponds to corn event MON 95379 and contains a contiguous sequence (contig) comprising the 5' LH244 flanking sequence, the transgene insert of event MON 95379, and the 3' LH244 flanking sequence, and thus contains the insert-to-plant genome junction sequences.

Unless otherwise noted herein, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may be found in Rieger et al., *Glossary of Genetics: Classical and Molecular*, 5<sup>th</sup> edition, Springer-Verlag: New York, 1991; and Lewin, *Genes V*, Oxford University Press: New York, 1994, along with other sources known to those of ordinary skill in the art. As used herein, the term “corn” means species belong to the genus *Zea*, preferably *Zea mays* and includes all plant varieties that can be bred with corn plants containing event MON 95379, including wild corn species as well as those plants belonging to the genus *Zea* that permit breeding between species.

The present invention provides for transgenic plants which have been transformed with a DNA construct that contains expression cassettes expressing toxic amounts of the insecticidal proteins Cry1B.868 and Cry1Da\_7. What is meant by toxic amount is an efficacious amount, an insecticidal amount, an insecticidally-effective amount, a target insect suppressive amount, an efficacious pesticidal amount, an amount in the diet of insects in the order of Lepidoptera that is insecticidal, and other similar terms to be understood according to conventional usage by those of ordinary skill in the relevant art. Corn plants transformed according to the methods and with the DNA construct disclosed herein are resistant to Lepidopteran insect pests.

A transgenic “plant” is produced by transformation of a plant cell with heterologous DNA, i.e., a polynucleic acid construct that includes a number of efficacious features of interest, regeneration of a plant resulting from the insertion of the transgene into the genome of the plant cell, and selection of a particular plant characterized by insertion into a particular genome location and the number of efficacious

features of the regenerated transgenic plant. The term “event” refers to DNA from the original transformant comprising the inserted DNA, and flanking genomic sequences immediately adjacent to the inserted DNA. Such DNA is unique and would be expected to be transferred to a progeny that receives the inserted DNA, including the transgene of interest, as the result of a sexual cross of a parental line that includes the inserted DNA (e.g., the original transformant and progeny resulting from selfing) and a parental line that does not contain the inserted DNA. The present invention also provides the original transformant plant and progeny of the transformant that include the heterologous DNA. Such progeny may be produced by a sexual outcross between plants comprising the event and another plant wherein the progeny includes the heterologous DNA. Even after repeated back-crossing to a recurrent parent, the event is present in the progeny of the cross at the same chromosomal location.

As used herein, the term “recombinant” refers to a non-natural DNA, protein, or organism that would not normally be found in nature and was created by human intervention. A “recombinant DNA molecule” is a DNA molecule comprising a combination of DNA molecules that would not naturally occur together and is the result of human intervention. For example, a DNA molecule that is comprised of a combination of at least two DNA molecules heterologous to each other, such as a DNA molecule that comprises a transgene and the plant genomic DNA adjacent to the transgene, is a recombinant DNA molecule.

The terms “DNA” and “DNA molecule” referred to herein refer to a deoxyribonucleic acid (DNA) molecule. A DNA molecule may be of genomic or synthetic origin, and is by convention from the 5' (upstream) end to the 3' (downstream) end. As used herein, the term “DNA sequence” refers to the nucleotide sequence of the DNA molecule. By convention, the DNA sequences of the invention and fragments thereof are disclosed with reference to only one strand of the two strand complementary DNA sequence strands. By implication and intent, the complementary sequences of the sequences provided here (the sequences of the complementary strand), also referred to in the art as the reverse complementary sequences, are within the scope of the invention and are expressly intended to be within the scope of the subject matter claimed.

As used herein, the term “fragment” refers to a smaller piece of the whole. For example, fragments of SEQ ID NO:10 would include sequences that are at least about 12 consecutive nucleotides, at least about 13 consecutive nucleotides, at least about 14 consecutive nucleotides, at least about 15 consecutive nucleotides, at least about 16 consecutive nucleotides, at least about 17 consecutive nucleotides, at least about 18 consecutive nucleotides, at least about 19 consecutive nucleotides, at least about 20 consecutive nucleotides, at least about 25 consecutive nucleotides, at least about 30 consecutive nucleotides, at least about 35 consecutive nucleotides, at least about 40 consecutive nucleotides, at least about 45 consecutive nucleotides, at least about 50 consecutive nucleotides, at least about 60 consecutive nucleotides, at least about 70 consecutive nucleotides, at least about 80 consecutive nucleotides, at least about 90 consecutive nucleotides, or at least about 100 consecutive nucleotides of the complete sequence of SEQ ID NO:10.

Reference in this application to an “isolated DNA molecule” or an equivalent term or phrase is intended to mean that the DNA molecule is one that is present alone or in combination with other compositions, but not within its

natural environment. For example, nucleic acid elements such as a coding sequence, intron sequence, untranslated leader sequence, promoter sequence, transcriptional termination sequence, and the like, that are naturally found within the DNA of the genome of an organism are not considered to be “isolated” so long as the element is within the genome of the organism and at the location within the genome in which it is naturally found. However, each of these elements, and subparts of these elements, would be “isolated” within the scope of this disclosure so long as the element is not within the genome of the organism and at the location within the genome in which it is naturally found. Similarly, a nucleotide sequence encoding an insecticidal protein or any naturally occurring insecticidal variant of that protein would be an isolated nucleotide sequence so long as the nucleotide sequence was not within the DNA of the bacterium from which the sequence encoding the protein is naturally found. A synthetic nucleotide sequence encoding the amino acid sequence of the naturally occurring insecticidal protein would be considered to be isolated for the purposes of this disclosure. For the purposes of this disclosure, any transgenic nucleotide sequence, i.e., the nucleotide sequence of the DNA inserted into the genome of the cells of a plant or bacterium, or present in an extrachromosomal vector, would be considered to be an isolated nucleotide sequence whether it is present within the plasmid or similar structure used to transform the cells, within the genome of the plant or bacterium, or present in detectable amounts in tissues, progeny, biological samples or commodity products derived from the plant or bacterium. In any circumstance, the isolated DNA molecule is a chemical molecule, regardless of whether it is referred to as a nucleic acid, a nucleic acid sequence, a polynucleotide sequence, and the like. It is a novel, inventive molecule that exhibits industrial applicability both when present in a plant cell or in a plant genome, and when present outside of a plant cell, and therefore, exhibits and is intended to exhibit such utility regardless of where the molecule is located.

The DNA sequence of the region spanning the connection by phosphodiester bond linkage of one end of the transgenic insert to the flanking corn genomic DNA is referred to as a “junction.” A junction is the connection point of the transgenic insert and flanking DNA as one contiguous molecule. One junction is found at the 5' end of the transgenic insert and the other is found at the 3' end of the transgenic insert, referred to herein as the 5' and 3' junction, respectively. A “junction sequence” refers to a DNA sequence of any length that spans the 5' or 3' junction of an event. Junction sequences of corn event MON 95379 are apparent to one of skill in the art using SEQ ID NO:10. Examples of junction sequences of event MON 95379 are provided as SEQ ID NOs:1-8. FIG. 1 illustrates the physical arrangement of the junction sequences, arranged from 5' to 3', relative to SEQ ID NO:10. The junction sequences of event MON 95379 may be present as part of the genome of a plant, seed, or cell containing event MON 95379. The identification of any one or more of the junction sequences in a sample from a plant, plant part, seed, or cell indicates that the DNA was obtained from corn containing event MON 95379, and is diagnostic for the presence of event MON 95379.

The junction sequences for event MON 95379 may be represented by a sequence from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:10. For example, the junction sequences may be arbitrarily represented by the nucleotide sequences provided as SEQ ID NO:1 and SEQ ID NO:2. Alternatively, the

junction sequences may be arbitrarily represented by the nucleotide sequences provided as SEQ ID NO:3 and SEQ ID NO:4. Alternatively, the junction sequences may be arbitrarily represented by the nucleotide sequences provided as SEQ ID NO:5 and SEQ ID NO:6. Alternatively, the junction sequences may be arbitrarily represented by the nucleotide sequences provided as SEQ ID NO:7 and SEQ ID NO:8. These nucleotides are connected by phosphodiester linkage, and in corn event MON 95379 are present as part of the recombinant plant cell genome.

The identification of one or more of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:10 in a sample derived from a corn plant, corn seed, or corn plant part is diagnostic that the DNA was obtained from corn event MON 95379. The invention thus provides a DNA molecule that contains at least one of the nucleotide sequences provided as SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. Any segment of DNA derived from transgenic corn event MON 95379 that is sufficient to include at least one of the sequences provided as SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10 is within the scope of the invention. In addition, any polynucleotide comprising a sequence complementary to any of the sequences described within this paragraph is within the scope of the invention.

The invention provides exemplary DNA molecules that can be used either as primers or probes for detecting the presence of DNA derived from a corn plant comprising event MON 95379 DNA in a sample. Such primers or probes are specific for a target nucleic acid sequence and, as such, are useful for the identification of corn event MON 95379 nucleic acid sequence by the methods of the invention described herein.

A "probe" is a nucleic acid molecule that is complementary to a strand of target nucleic acid and is useful in hybridization methods. A probe may be attached a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid and, in the case of the present invention, to a strand of DNA from event MON 95379 whether from an event MON 95379 containing plant or from a sample that includes event MON 95379 DNA. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids, but also polyamides and other probe materials that bind specifically to a target DNA sequence and can be used to detect the presence of that target DNA sequence. Exemplary DNA sequences useful as a probe for detecting corn event MON 95379 are provided as: SEQ ID NO:17 (PB10269), SEQ ID NO:23 (PB50340); SEQ ID NO:26 (PB50138).

A "primer" is typically a DNA molecule that is designed for use in specific annealing or hybridization methods that involve thermal amplification. A pair of primers may be used with template DNA (such as a sample of corn genomic DNA) in a thermal amplification (such as polymerase chain reaction (PCR)) to produce an amplicon, where the amplicon produced from such reaction would have a DNA sequence corresponding to sequence of the template DNA located between the two sites where the primers hybridized to the template.

A primer is typically designed to hybridize to a complementary target DNA strand to form a hybrid between the primer and target DNA strand, and the presence of the

primer is a point of recognition by a polymerase to begin extension of the primer (i.e., polymerization of additional nucleotides into a lengthening nucleotide molecule) using as a template the target DNA strand. Primer pairs refer to use of two primers binding opposite strands of a double stranded nucleotide segment for the purpose of amplifying linearly the polynucleotide segment between the positions targeted for binding by the individual members of the primer pair, typically in a thermal amplification reaction or other conventional nucleic-acid amplification methods. Exemplary DNA molecules useful as primers are provided as SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25 and SEQ ID NO:27.

The primer pair SEQ ID NO:15 and SEQ ID NO:16 are useful as a first DNA molecule and a second DNA molecule that is different from the first DNA molecule, and both are each of sufficient length of contiguous nucleotides of SEQ ID NO:10 to function as DNA primers that, when used together in a thermal amplification reaction with template DNA derived from corn event MON 95379, to produce an amplicon diagnostic for corn event MON 95379 DNA in a sample. The primer pair SEQ ID NO:21 and SEQ ID NO:22 are useful as a first DNA molecule and a second DNA molecule that is different from the first DNA molecule, and both are each of sufficient length of contiguous nucleotides of SEQ ID NO:10 to function as DNA primers that, when used together in a thermal amplification reaction with template DNA derived from corn event MON 95379, to produce an amplicon diagnostic for the zygosity of corn event MON 95379 DNA in a sample. The primer pair SEQ ID NO:24 and SEQ ID NO:25 are useful as a first DNA molecule and a second DNA molecule that is different from the first DNA molecule, and both are each of sufficient length of contiguous nucleotides of SEQ ID NO:10 to function as DNA primers that, when used together in a thermal amplification reaction with template DNA derived from corn event MON 95379, to produce an amplicon diagnostic for the zygosity of corn event MON 95379 DNA in a sample. The primer pair SEQ ID NO:18 and SEQ ID NO:19 are useful as a first DNA molecule and a second DNA molecule that is different from the first DNA molecule, and both are each of sufficient length of contiguous nucleotides of a locus within the corn genome to function as DNA primers that, when used together in a thermal amplification reaction with template DNA derived from corn event MON 95379, to produce an amplicon that serves as an internal control for both the diagnosis of corn event MON 95379, as well as the zygosity of corn event MON 95379 DNA in a sample.

DNA probes and DNA primers are generally eleven (11) polynucleotides or more in length, often eighteen (18) polynucleotides or more, twenty-four (24) polynucleotides or more, or thirty (30) polynucleotides or more. Such probes and primers are selected to be of sufficient length to hybridize specifically to a target sequence under high stringency hybridization conditions. Preferably, probes and primers according to the present invention have complete sequence similarity with the target sequence, although probes differing from the target sequence that retain the ability to hybridize to target sequences may be designed by conventional methods.

The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA molecule. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic plant in a sample. Polynucleic acid molecules also referred to as nucleic acid segments or

fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances.

As used herein, two polynucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be "minimally complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional "low-stringency" conditions. Similarly, the molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions. Conventional stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, D.C. (1985). Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

As used herein, a substantially homologous sequence is a nucleic acid sequence that will specifically hybridize to the complement of the nucleic acid sequence to which it is being compared under high stringency conditions. Appropriate stringency conditions that promote DNA hybridization, for example, 6.0× sodium chloride/sodium citrate (SSC) at about 45° C., followed by a wash of 2.0×SSC at 50° C., are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0×SSC at 50° C. to a high stringency of about 0.2×SSC at 50° C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22° C., to high stringency conditions at about 65° C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed. In a preferred embodiment, a polynucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, or SEQ ID NO:10, or complements thereof or fragments thereof under moderately stringent conditions, for example at about 2.0×SSC and about 65° C. In a particularly preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, or SEQ ID NO:10, or complements or fragments thereof under high stringency conditions. In one aspect of the present invention, a preferred marker nucleic acid molecule of the present invention has the nucleic acid sequence set forth in SEQ ID NO:1, or SEQ ID NO:2, or SEQ ID NO:3, or SEQ ID NO:4, or SEQ ID NO:5, or SEQ ID NO:6, or SEQ ID NO:7, or SEQ ID NO:8, or SEQ ID

NO:9, or SEQ ID NO:10, or complements thereof, or fragments thereof. The hybridization of the probe to the target DNA molecule can be detected by any number of methods known to those skilled in the art, these can include, but are not limited to, fluorescent tags, radioactive tags, antibody based tags, and chemiluminescent tags.

Regarding the amplification of a target nucleic acid sequence (e.g., by PCR) using a particular amplification primer pair, "stringent conditions" are conditions that permit the primer pair to hybridize only to the target nucleic acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA thermal amplification reaction.

The term "specific for (a target sequence)" indicates that a probe or primer hybridizes under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

As used herein, "amplified DNA" or "amplicon" refers to the product of polynucleic acid amplification method directed to a target polynucleic acid molecule that is part of a polynucleic acid template. For example, to determine whether a corn plant resulting from a sexual cross contains transgenic plant genomic DNA from a corn plant comprising event MON 95379 of the present invention, DNA that is extracted from a corn plant tissue sample may be subjected to a polynucleic acid amplification method using a primer pair that includes a first primer derived from a genomic DNA sequence in the region flanking the heterologous inserted DNA of event MON 95379 and is elongated by polymerase 5' to 3' in the direction of the inserted DNA. The second primer is derived from the heterologous inserted DNA molecule is elongated by the polymerase 5' to 3' in the direction of the flanking genomic DNA from which the first primer is derived. The amplicon may range in length from the combined length of the primer pair plus one nucleotide base pair, or plus about fifty nucleotide base pairs, or plus about two hundred-fifty nucleotide base pairs, or plus about four hundred-fifty nucleotide base pairs or more. Alternatively, a primer pair can be derived from genomic sequence on both sides of the inserted heterologous DNA so as to produce an amplicon that includes the entire insert polynucleotide sequence (e.g., a forward primer isolated from the genomic portion on the 5' end of SEQ ID NO:10 and a reverse primer isolated from the genomic portion on the 3' end of SEQ ID NO:10 that amplifies a DNA molecule comprising the inserted DNA sequence (SEQ ID NO:9) identified herein in the event MON 95379 genome). A member of a primer pair derived from the plant genomic sequence adjacent to the inserted transgenic DNA is located a distance from the inserted DNA sequence, this distance can range from one nucleotide base pair up to about twenty thousand nucleotide base pairs. The use of the term "amplicon" specifically excludes primer dimers that may be formed in the DNA thermal amplification reaction.

For practical purposes, one should design primers which produce amplicons of a limited size range, for example, between 100 to 1000 bases. Smaller (shorter polynucleotide length) sized amplicons in general are more reliably produced in thermal amplification reactions, allow for shorter cycle times, and can be easily separated and visualized on agarose gels or adapted for use in endpoint TAQMAN®-like assays. Smaller amplicons can be produced and detected by methods known in the art of DNA amplicon detection. In addition, amplicons produced using the primer pairs can be cloned into vectors, propagated, isolated, and sequenced or can be sequenced directly with methods well established in

the art. Any primer pair derived from the combination of SEQ ID NO:11 and SEQ ID NO:9 or the combination of SEQ ID NO:12 and SEQ ID NO:9 that are useful in a DNA amplification method to produce an amplicon diagnostic for event MON 95379 or progeny thereof is an aspect of the invention. Any single isolated DNA polynucleotide primer molecule comprising at least 15 contiguous nucleotides of SEQ ID NO:11, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for event MON 95379 or progeny thereof is an aspect of the invention. Any single isolated DNA polynucleotide primer molecule comprising at least 15 contiguous nucleotides of SEQ ID NO:12, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for plants comprising event MON 95379 or progeny thereof is an aspect of the invention. Any single isolated DNA polynucleotide primer molecule comprising at least 15 contiguous nucleotides of SEQ ID NO:9, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for event MON 95379 or progeny thereof is an aspect of the invention.

Polynucleic acid amplification can be accomplished by any of the various polynucleic acid amplification methods known in the art, including the polymerase chain reaction (PCR). Amplification methods are known in the art and are described, inter alia, in U.S. Pat. Nos. 4,683,195 and 4,683,202 and in *PCR Protocols: A Guide to Methods and Applications*, ed. Innis et al., Academic Press, San Diego, 1990. PCR amplification methods have been developed to amplify up to 22 kb (kilobase) of genomic DNA and up to 42 kb of bacteriophage DNA (Cheng et al., *Proc. Natl. Acad. Sci. USA* 91:5695-5699, 1994). These methods as well as other methods known in the art of DNA amplification may be used in the practice of the present invention. The sequence of the heterologous DNA insert or flanking genomic DNA sequence from corn event MON 95379 can be verified (and corrected if necessary) by amplifying such DNA molecules from corn seed containing event MON 95379 DNA or corn plants grown from the corn seed containing event MON 95379 DNA deposited with the ATCC having accession No. PTA-125027, using primers derived from the sequences provided herein, followed by standard DNA sequencing of the PCR amplicon or cloned DNA fragments thereof.

The diagnostic amplicon produced by these methods may be detected by a plurality of techniques. One such method is Genetic Bit Analysis (Nikiforov, et al. *Nucleic Acid Res.* 22:4167-4175, 1994) where a DNA oligonucleotide is designed that overlaps both the adjacent flanking genomic DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microtiter plate. Following PCR of the region of interest (using one primer in the inserted sequence and one in the adjacent flanking genomic sequence), a single-stranded PCR product can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labeled dideoxynucleotide triphosphates (ddNTPs) specific for the expected next base. Readout may be fluorescent or ELISA-based. A signal indicates presence of the transgene/genomic sequence due to successful amplification, hybridization, and single base extension.

Another method is the Pyrosequencing technique as described by Winge (Innov. Pharma. Tech. 00:18-24, 2000). In this method, an oligonucleotide is designed that overlaps the adjacent genomic DNA and insert DNA junction. The oligonucleotide is hybridized to single-stranded PCR product from the region of interest (one primer in the inserted sequence and one in the flanking genomic sequence) and

incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. DNTPs are added individually and the incorporation results in a light signal that is measured. A light signal indicates the presence of the transgene/genomic sequence due to successful amplification, hybridization, and single or multi-base extension.

Fluorescence Polarization as described by Chen, et al., (*Genome Res.* 9:492-498, 1999) is a method that can be used to detect the amplicon of the present invention. Using this method an oligonucleotide is designed that overlaps the genomic flanking and inserted DNA junction. The oligonucleotide is hybridized to single-stranded PCR product from the region of interest (one primer in the inserted DNA and one in the flanking genomic DNA sequence) and incubated in the presence of a DNA polymerase and a fluorescent-labeled ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorometer. A change in polarization indicates the presence of the transgene/genomic sequence due to successful amplification, hybridization, and single base extension.

Real-time Polymerase Chain Reaction (PCR) is the ability to monitor the progress of the PCR as it occurs (i.e., in real time). Data is collected throughout the PCR process, rather than at the end of the PCR. In real-time PCR, reactions are characterized by the point in time during cycling when amplification of a target is first detected rather than the amount of target accumulated after a fixed number of cycles. In a real-time PCR assay, a positive reaction is detected by accumulation of a fluorescent signal. The higher the starting copy number of the nucleic acid target, the sooner a significant increase in fluorescence is observed. The cycle threshold (Ct value) is defined as the number of cycles required for the fluorescent signal to cross the threshold (i.e., exceeds background level). Ct levels are inversely proportional to the amount of target nucleic acid in the sample (i.e., the lower the Ct value, the greater the amount of target nucleic acid in the sample).

Taqman® (PE Applied Biosystems, Foster City, Calif.) is described as a method of detecting and quantifying the presence of a DNA sequence using real-time PCR and is fully understood in the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed that overlaps the genomic flanking and insert DNA junction. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermalstable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the transgene/genomic sequence due to successful amplification and hybridization.

Molecular Beacons have been described for use in sequence detection as described in Tyangi, et al. (*Nature Biotech.* 14:303-308, 1996). Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking genomic and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermalstable polymerase and dNTPs. Following successful PCR amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties. A fluores-

cent signal results. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

DNA detection kits that are based on DNA amplification methods contain DNA primer molecules that hybridize specifically to a target DNA and amplify a diagnostic amplicon under the appropriate reaction conditions. The kit may provide an agarose gel based detection method or any number of methods of detecting the diagnostic amplicon that are known in the art. DNA detection kits can be developed using the compositions disclosed herein and are useful for identification of corn event MON 95379 DNA in a sample and can be applied to methods for breeding corn plants containing event MON 95379 DNA. A kit that contains DNA primers that are homologous or complementary to any portion of the corn genomic region as set forth in SEQ ID NO:10 and to any portion of the inserted transgenic DNA as set forth in SEQ ID NO:10 is an object of the invention. The DNA molecules can be used in DNA amplification methods (PCR) or as probes in polynucleic acid hybridization methods, i.e., southern analysis, northern analysis. Kits of the invention may optionally also comprise reagents or instructions for performing the detection or diagnostic reactions described herein.

Probes and primers according to the invention may have complete sequence identity with the target sequence, although primers and probes differing from the target sequence that retain the ability to hybridize preferentially to target sequences may be designed by conventional methods. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of transgenic DNA from corn event MON 95379 in a sample. Probes and primers are generally at least about 11 nucleotides, at least about 18 nucleotides, at least about 24 nucleotides, or at least about 30 nucleotides or more in length. Such probes and primers hybridize specifically to a target DNA sequence under stringent hybridization conditions. Conventional stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, D.C. (1985).

Any number of methods well known to those skilled in the art can be used to isolate and manipulate a DNA molecule, or fragment thereof, disclosed in the invention, including thermal amplification methods. DNA molecules, or fragments thereof, can also be obtained by other techniques such as by directly synthesizing the fragment by chemical means, as is commonly practiced by using an automated oligonucleotide synthesizer.

The DNA molecules and corresponding nucleotide sequences provided herein are therefore useful for, among other things, identifying corn event MON 95379, detecting the presence of DNA derived from the transgenic corn event MON 95379 in a sample, and monitoring samples for the presence and/or absence of corn event MON 95379 or plant parts derived from corn plants comprising event MON 95379.

The invention provides corn plants, corn plant cells, corn seeds, corn plant parts (such as pollen, ovule, silk, spike, anther, cob, root tissue, stalk tissue, leaf tissue), corn progeny plants, and corn commodity products. These corn plants, corn plant cells, corn seeds, corn plant parts, corn progeny plants, and corn commodity products contain a detectable

amount of a polynucleotide of the invention, e.g., such as a polynucleotide having at least one of the sequences provided as SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. Corn plants, plant cells, seeds, plant parts, and progeny plants of the invention may also contain one or more additional transgenes. Such additional transgene may be any nucleotide sequence encoding a protein or RNA molecule conferring a desirable trait including but not limited to increased insect resistance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, and/or increased herbicide tolerance.

The invention provides corn plants, corn plant cells, corn seeds, corn plant parts (such as pollen, ovule, silk, spike, anther, cob, root tissue, stalk tissue, leaf tissue), corn progeny plants derived from a transgenic corn plant containing event MON 95379 DNA. A representative sample of corn seed containing event MON 95379 DNA has been deposited according to the Budapest Treaty with the American Type Culture Collection (ATCC®). The ATCC repository has assigned the Patent Deposit Designation PTA-125027 to the seed containing event MON 95379 DNA.

The invention provides a microorganism comprising a DNA molecule having at least one sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10 present in its genome. An example of such a microorganism is a transgenic plant cell. Microorganisms, such as a plant cell of the invention, are useful in many industrial applications, including but not limited to: (i) use as research tool for scientific inquiry or industrial research; (ii) use in culture for producing endogenous or recombinant carbohydrate, lipid, nucleic acid, or protein products or small molecules that may be used for subsequent scientific research or as industrial products; and (iii) use with modern plant tissue culture techniques to produce transgenic plants or plant tissue cultures that may then be used for agricultural research or production. The production and use of microorganisms such as transgenic plant cells utilizes modern microbiological techniques and human intervention to produce a man-made, unique microorganism. In this process, recombinant DNA is inserted into a plant cell's genome to create a transgenic plant cell that is separate and unique from naturally occurring plant cells. This transgenic plant cell can then be cultured much like bacteria and yeast cells using modern microbiology techniques and may exist in an undifferentiated, unicellular state. The transgenic plant cell's new genetic composition and phenotype is a technical effect created by the integration of the heterologous DNA into the genome of the cell. Another aspect of the invention is a method of using a microorganism of the invention. Methods of using microorganisms of the invention, such as transgenic plant cells, include (i) methods of producing transgenic cells by integrating recombinant DNA into the genome of the cell and then using this cell to derive additional cells possessing the same heterologous DNA; (ii) methods of culturing cells that contain recombinant DNA using modern microbiology techniques; (iii) methods of producing and purifying endogenous or recombinant carbohydrate, lipid, nucleic acid, or protein products from cultured cells; and (iv) methods of using modern plant tissue culture techniques with transgenic plant cells to produce transgenic plants or transgenic plant tissue cultures.

Plants of the invention may pass along the event MON 95379 DNA, including transgene inserted in corn event MON 95379, to progeny. As used herein, "progeny"

includes any plant, plant cell, seed, and/or regenerable plant part containing the event MON 95379 DNA derived from an ancestor plant and/or comprising a DNA molecule having at least one sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. Plants, progeny, and seeds may be homozygous or heterozygous for the transgene of event MON 95379. Progeny may be grown from seeds produced by a corn event MON 95379 containing plant and/or from seeds produced by a plant fertilized with pollen from a corn event MON 95379 containing plant.

Progeny plants may be self-pollinated (also known as “selfing”) to generate a true breeding line of plants, i.e., plants homozygous for the transgene. Selfing of appropriate progeny can produce plants that are homozygous for both added exogenous genes.

Alternatively, progeny plants may be out-crossed, e.g., bred with another unrelated plant, to produce a varietal or a hybrid seed or plant. The other unrelated plant may be transgenic or non-transgenic. A varietal or hybrid seed or plant of the invention may thus be derived by sexually crossing a first parent that lacks the specific and unique DNA of the corn event MON 95379 with a second parent comprising corn event MON 95379, resulting in a hybrid comprising the specific and unique DNA of the corn event MON 95379. Each parent can be a hybrid or an inbred/varietal, so long as the cross or breeding results in a plant or seed of the invention, i.e., a seed having at least one allele containing the DNA of corn event MON 95379 and/or a DNA molecule having at least one sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. Two different transgenic plants may thus be crossed to produce hybrid offspring that contain two independently segregating, added, exogenous genes. For example, event MON 95379 containing Cry1B.868 and Cry1Da\_7 conferring insect resistance to corn can be crossed with other transgenic corn plants to produce a plant having the characteristics of both transgenic parents. One example of this would be a cross of event MON 95379 containing Cry1B.868 and Cry1Da\_7 conferring Lepidopteran resistance to corn with a plant having one or more additional traits such as herbicide tolerance, insect resistance, or drought tolerance, resulting in a progeny plant or seed that has resistance to Lepidopteran insect pests and has at least one or more additional traits. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison Wis. (1987).

Plants, progeny, seeds, cells and plant parts of the invention may also contain one or more additional corn trait(s) or transgenic events, particularly those introduced by crossing a corn plant containing corn event MON 95379 with another corn plant containing the additional trait(s) or transgenic events. Such trait(s) or transgenic events include, but are not limited to, increased insect resistance, herbicide tolerance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, improved nutritional quality, hybrid seed production, or disease or fungal resistance. Corn transgenic events are known to those of skill in the art. For example, a list of such traits is provided by the United States Department of Agri-

culture’s (USDA) Animal and Plant Health Inspection Service (APHIS) and can be found on their website: [www.aphis.usda.gov](http://www.aphis.usda.gov). Two or more transgenic events may thus be combined in a progeny seed or plant by crossing two parent plants each comprising one or more transgenic events, collecting the progeny seed, and selecting for progeny seed or plants that contain the two or more transgenic events. These steps may be repeated until the desired combination of transgenic events in a progeny is achieved. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, and is vegetative propagation.

The invention provides a plant part that is derived from corn plants comprising event MON 95379. As used herein, a “plant part” refers to any part of a plant which is comprised of material derived from a corn plant comprising event MON 95379. Plant parts include but are not limited to pollen, ovule, silk, spike, anther, cob, root tissue, stalk tissue, and leaf tissue. Plant parts may be viable, nonviable, regenerable, and/or nonregenerable.

The invention provides a commodity product that is derived from corn plants comprising event MON 95379 and that contains a detectable amount of a nucleic acid specific for event MON 95379. As used herein, a “commodity product” refers to any composition or product which is comprised of material derived from a corn plant, whole or processed corn seed, or one or more plant cells and/or plant parts containing the corn event MON 95379 DNA. Nonviable commodity products include but are not limited to nonviable seeds, whole or processed seeds, seed parts, and plant parts; animal feed comprising corn, corn oil, corn meal, corn flour, corn flakes, corn bran, pasta made with corn, corn biomass, and fuel products produced using corn and corn parts. Viable commodity products include but are not limited to seeds, plants, and plant cells. The corn plants comprising event MON 95379 can thus be used to manufacture any commodity product typically acquired from corn. Any such commodity product that is derived from corn plants comprising event MON 95379 may contain at least a detectable amount of the specific and unique DNA corresponding to corn event MON 95379, and specifically may contain a detectable amount of a polynucleotide comprising a DNA molecule having at least one sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. Any standard method of detection for nucleotide molecules may be used, including methods of detection disclosed herein. A commodity product is within the scope of the invention if there is any detectable amount of a DNA molecule having at least one sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10 in the commodity product.

The corn plants, corn plant cells, corn seeds, corn plant parts (such as pollen, ovule, silk, spike, anther, cob, root tissue, stalk tissue, leaf tissue), corn progeny plants, and commodity products of the invention are therefore, useful for, among other things, growing plants for the purpose of producing seed and/or plant parts comprising corn event MON 95379 for agricultural purposes, producing progeny comprising corn event MON 95379 for plant breeding and research purposes, use with microbiological techniques for industrial and research applications, and sale to consumers.

Methods for producing an insect resistant corn plant comprising the DNA sequences specific and unique to event MON 95379 of the invention are provided. Transgenic plants used in these methods may be homozygous or het-

erozygous for the transgene. Progeny plants produced by these methods may be varietal or hybrid plants; may be grown from seeds produced by corn event MON 95379 containing plant and/or from seeds produced by a plant fertilized with pollen from a corn event MON 95379 containing plant; and may be homozygous or heterozygous for the transgene. Progeny plants may be subsequently self-pollinated to generate a true breeding line of plants, i.e., plants homozygous for the transgene, or alternatively may be out-crossed, e.g., bred with another unrelated plant, to produce a varietal or a hybrid seed or plant.

Methods of detecting the presence of DNA derived from a corn cell, corn tissue, corn seed, or corn plant comprising corn event MON 95379 in a sample are provided. One method consists of (i) extracting a DNA sample from at least one corn cell, corn tissue, corn seed, or corn plant; (ii) contacting the DNA sample with at least one primer that is capable of producing DNA sequence specific to event MON 95379 DNA under conditions appropriate for DNA sequencing; (iii) performing a DNA sequencing reaction; and then (iv) confirming that the nucleotide sequence comprises a nucleotide sequence specific for event MON 95379, of the construct comprised therein, such as one selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. Another method consists of (i) extracting a DNA sample from at least one corn cell, corn tissue, corn seed, or corn plant; (ii) contacting the DNA sample with a primer pair that is capable of producing an amplicon from event MON 95379 DNA under conditions appropriate for DNA amplification; (iii) performing a DNA amplification reaction; and then (iv) detecting the amplicon molecule and/or confirming that the nucleotide sequence of the amplicon comprises a nucleotide sequence specific for event MON 95379, such as one selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. The amplicon should be one that is specific for event MON 95379, such as an amplicon that comprises SEQ ID NO:1, or SEQ ID NO:2, or SEQ ID NO:3, or SEQ ID NO:4, or SEQ ID NO:5, or SEQ ID NO:6, or SEQ ID NO:7, or SEQ ID NO:8, or SEQ ID NO:9, or SEQ ID NO:10. The detection of a nucleotide sequence specific for event MON 95379 in the amplicon is determinative and/or diagnostic for the presence of the corn event MON 95379 specific DNA in the sample. An example of a primer pair that is capable of producing an amplicon from event MON 95379 DNA under conditions appropriate for DNA amplification is provided as SEQ ID NO:15 and SEQ ID NO:16. Other primer pairs may be readily designed by one of skill in the art and would produce an amplicon comprising SEQ ID NO:1, or SEQ ID NO:2, or SEQ ID NO:3, or SEQ ID NO:4, or SEQ ID NO:5, or SEQ ID NO:6, or SEQ ID NO:7, or SEQ ID NO:8 wherein such a primer pair comprises at least one primer within the genomic region flanking the insert and a second primer within the insert. Another method of detecting the presence of DNA derived from a corn cell, corn tissue, corn seed, or corn plant comprising corn event MON 95379 in a sample consists of (i) extracting a DNA sample from at least one corn cell, corn tissue, corn seed, or corn plant; (ii) contacting the DNA sample with a DNA probe specific for event MON 95379 DNA; (iii) allowing the probe and the DNA sample to hybridize under stringent hybridization conditions; and then (iv) detecting hybridization between the probe and the target DNA sample. An example of the sequence of a DNA probe

that is specific for event MON 95379 is provided as SEQ ID NO:17. Other probes may be readily designed by one of skill in the art and would comprise at least one fragment of genomic DNA flanking the insert and at least one fragment of insert DNA such as the sequence provided in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:10. Detection of probe hybridization to the DNA sample is diagnostic for the presence of corn event MON 95379 specific DNA in the sample. Absence of hybridization is alternatively diagnostic of the absence of corn event MON 95379 specific DNA in the sample.

DNA detection kits are provided that are useful for the identification of corn event MON 95379 DNA in a sample and can also be applied to methods for breeding corn plants containing the appropriate event DNA. Such kits contain DNA primers and/or probes comprising fragments of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. One example of such a kit comprises at least one DNA molecule of sufficient length of continuous nucleotides of SEQ ID NO:10 to function as a DNA probe useful for detecting the presence and/or absence of DNA derived from transgenic corn plants comprising event MON 95379 in a sample. The DNA derived from transgenic corn plants comprising event MON 95379 would comprise a DNA molecule having at least one sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. A DNA molecule sufficient for use as a DNA probe is provided that is useful for determining, detecting, or diagnosing the presence and/or absence of corn event MON 95379 DNA in a sample is provided as SEQ ID NO: 17. Other probes may be readily designed by one of skill in the art and should comprise at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, at least 30, at least 31, at least 32, at least 33, at least 34, at least 35, at least 36, at least 37, at least 38, at least 39, or at least 40 contiguous nucleotides of SEQ ID NO:10 and be sufficiently unique to corn event MON 95379 DNA in order to identify DNA derived from the event.

Another type of kit comprises a primer pair useful for producing an amplicon useful for detecting the presence and/or absence of DNA derived from transgenic corn event MON 95379 in a sample. Such a kit would employ a method comprising contacting a target DNA sample with a primer pair as described herein, then performing a nucleic acid amplification reaction sufficient to produce an amplicon comprising a DNA molecule having at least one sequence selected from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10 and then detecting the presence and/or absence of the amplicon. Such a method may also include sequencing the amplicon or a fragment thereof, which would be determinative of, i.e., diagnostic for, the presence of the corn event MON 95379 specific DNA in the target DNA sample. Other primer pairs may be readily designed by one of skill in the art and should comprise at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, or at least 30 contiguous nucleotides of sequences provided in, but not limited to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10,

and be sufficiently unique to corn event MON 95379 DNA in order to identify DNA derived from the event.

The kits and detection methods of the invention are useful for, among other things, identifying corn event MON 95379, selecting plant varieties or hybrids comprising corn event MON 95379, detecting the presence of DNA derived from the transgenic corn plant comprising event MON 95379 in a sample, and monitoring samples for the presence and/or absence of corn plants comprising event MON 95379, or plant parts derived from corn plants comprising event MON 95379.

The sequences of the heterologous DNA insert, junction sequences, or flanking sequence from corn event MON 95379 can be verified (and corrected if necessary) by amplifying such sequences from the event using primers derived from the sequences provided herein followed by standard DNA sequencing of the amplicon or of the cloned DNA.

Methods of detecting the zygosity of the transgene allele of DNA derived from a corn cell, corn tissue, corn seed, or corn plant comprising corn event MON 95379 in a sample are provided. One method consists of (i) extracting a DNA sample from at least one corn cell, corn tissue, corn seed, or corn plant; (ii) contacting the DNA sample with a primer pair that is capable of producing a first amplicon diagnostic for event MON 95379; (iii) contacting the DNA sample with a primer pair that is capable of producing a second amplicon diagnostic for native corn genomic DNA not comprising event MON 95379; (iv) performing a DNA amplification reaction; and then (v) detecting the amplicons, wherein the presence of only the first amplicon is diagnostic of a homozygous event MON 95379 DNA in the sample, and the presence of both the first amplicon and the second amplicon is diagnostic of a corn plant heterozygous for event MON 95379 allele. An exemplary set of primers pairs are presented as SEQ ID NO:15 and SEQ ID NO:16 which produce an amplicon diagnostic for event MON 95379; and SEQ ID NO:15 and SEQ ID NO:27 which produces an amplicon diagnostic for non-inserted wild-type corn genomic DNA not comprising event MON 95379. A set of probes can also be incorporated into such an amplification method to be used in a real-time PCR format using the primer pair sets described above. An exemplary set of probes are presented as SEQ ID NO:17 (diagnostic for the amplicon for the event MON 95379) and SEQ ID NO:28 (diagnostic for the amplicon for wild-type corn genomic DNA not comprising event MON 95379).

Another method for determining zygosity consists of (i) extracting a DNA sample from at least one corn cell, corn tissue, corn seed, or corn plant; (ii) contacting the DNA sample with a probe set which contains at least a first probe that specifically hybridizes to event MON 95379 DNA and at least a second probe that specifically hybridizes to corn genomic DNA that was disrupted by insertion of the heterologous DNA of event MON 95379 and does not hybridize to event MON 95379 DNA; (iii) hybridizing the probe set with the sample under stringent hybridization conditions, wherein detecting hybridization of only the first probe under the hybridization conditions is diagnostic for a homozygous allele of event MON 95379 DNA in the sample; and wherein detecting hybridization of both the first probe and the second probe under the hybridization conditions is diagnostic for a heterozygous allele of event MON 95379 in a DNA sample.

Yet another method for determining zygosity consists of (i) extracting a DNA sample from at least one corn cell, corn tissue, corn seed, or corn plant; (ii) contacting the DNA sample with a primer pair that is capable of producing an amplicon of one of the toxin coding sequences encoding

Cry1B.868 or Cry1Da<sub>7</sub>; (iii) contacting the DNA sample with a primer pair that is capable of producing an amplicon of an internal standard known to be single-copy and homozygous in the corn plant; (iv) contacting the DNA sample with a probe set which contains at least a first probe that specifically hybridizes to one of the toxin coding sequences encoding Cry1B.868 or Cry1Da<sub>7</sub>, and at least a second probe that specifically hybridizes to the internal standard genomic DNA known to be single-copy and homozygous in the corn plant; (v) performing a DNA amplification reaction using real-time PCR and determining the cycle thresholds (Ct values) of the amplicon corresponding to the toxin coding sequence and the single-copy, homozygous internal standard; (vi) calculating the difference ( $\Delta$ Ct) between the Ct value of the single-copy, homozygous internal standard amplicon and the Ct value of the toxin coding sequence amplicon; and (vii) determining zygosity, wherein a  $\Delta$ Ct of around zero (0) indicates homozygosity of the inserted T-DNA and a  $\Delta$ Ct of around one (1) indicates heterozygosity of the inserted T-DNA. Heterozygous and homozygous events are differentiated by a  $\Delta$ Ct value unit of approximately one (1). Given the normal variability observed in real-time PCR due to multiple factors such as amplification efficiency and ideal annealing temperatures, the range of "about one (1)" is defined as a  $\Delta$ Ct of 0.75 to 1.25. Primer pairs and probes for the above method for determining zygosity can either amplify and detect amplicons from the Cry1B.868 coding sequence and internal standard, or amplify and detect amplicons from the Cry1Da<sub>7</sub> coding sequence and internal standard. Exemplary primer pairs for the detection of the amplicons corresponding to the Cry1B.868 coding sequence and internal standard are presented as SEQ ID NO:18 combined with SEQ ID NO:19 (internal standard) and SEQ ID NO:21 combined with SEQ ID NO:22 (Cry1B.868). The accompanying exemplary probes are presented as SEQ ID NO:20 (internal standard) and SEQ ID NO:23 (Cry1B.868). Exemplary primer pairs for the detection of the amplicons corresponding to the Cry1Da<sub>7</sub> coding sequence and internal standard are presented as SEQ ID NO:18 combined with SEQ ID NO:19 (internal standard) and SEQ ID NO:24 combined with SEQ ID NO:25 (Cry1Da<sub>7</sub>). The accompanying exemplary probes are presented as SEQ ID NO:20 (internal standard) and SEQ ID NO:26 (Cry1Da<sub>7</sub>).

#### Deposit Information

A deposit of a representative sample of corn seed containing event MON 95379 was made on Apr. 20, 2018 according to the Budapest Treaty with the American Type Culture Collection (ATCC) having an address at 10801 University Boulevard, Manassas, Va. USA, Zip Code 20110, and assigned ATCC Accession No. PTA-125027. Access to the deposits will be available during the pendency of the application to the Commissioner of Patents and Trademarks and persons determined by the Commissioner to be entitled thereto upon request. Upon issuance of the patent, all restrictions upon availability to the public will be irrevocably removed. The deposit has been accepted under the Budapest Treaty and will be maintained in the depository for a period of thirty (30) years, or five (5) years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced as necessary during that period.

#### EXAMPLES

The following Examples are included to more fully describe the invention. Summarized are the construction and

testing of one hundred and twenty-five (125) constructs, the production of about ten thousand seven hundred and eighty-five (10,785) events (both proof of concept and commercial), and the analysis of hundreds of thousands of individual plants over six (6) years through the rigorous molecular, agronomic, and field testing required for the creation and selection of corn event MON 95379.

The Examples demonstrate certain preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the Examples that follow represent approaches the inventors have found function well in the practice of the invention, and thus can be considered to constitute examples of preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

#### Example 1

##### Expression Cassette Testing, Construct Design, Plant Testing and Construct Selection

Transgene expression in plants is influenced by numerous different factors. The right combination of insecticidal proteins and different expression elements driving expression in plants, while not resulting in off-phenotypes, must be found. Further, beyond the expression elements themselves and their combination and orientation in a cassette, the expression of transgenes in plants is known to be influenced by chromosomal insertion position, perhaps due to chromatin structure (e.g., heterochromatin) or the proximity of transcriptional regulation elements (e.g., enhancers) close to the integration site (Kurt Weising et al., (1988) *Foreign genes in plants: transfer, structure, expression and applications. Annu. Rev. Genet.* 22: 421-77). For example, it has been observed in plants and in other organisms that there may be wide variation in the levels of expression of an introduced gene from the same construct among events with different chromosomal insertion positions. Different chromosomal insertion positions may also produce differences in spatial or temporal patterns of expression, that may not correspond to the patterns expected from transcriptional regulatory elements present in the introduced gene construct.

For these reasons, it is often necessary to create and screen a large number of constructs and transformation events in order to identify a construct, and then an event, which demonstrates optimal expression of the introduced genes of interest, while also not producing agronomic or phenotypic off-types.

For these reasons, the development of a transgenic corn plant comprising insecticidal proteins that were active against Lepidopterans without any negative effects on agronomics, yield, or stacking viability required extensive research, development, and analysis. Specifically, over a six (6) year period, approximately ten thousand, seven hundred eight-five (10,785) proof of concept and commercial transgenic events derived from one hundred twenty-five (125) different plasmid vector constructs were developed, tested, and analyzed.

This Example describes the design and testing in corn plants of one hundred and twenty five (125) different constructs to identify the preferred construct for event creation. Each construct varied with respect to the coding sequences for the insecticidal proteins and the transcriptional regulatory elements. Testing was done to select the best construct

for use in expressing the insecticidal proteins in plants. Each construct had a unique configuration, varying by expression cassette composition (both insecticidal proteins and expression elements), orientation, and whether or not proteins were targeted to the chloroplast.

In an initial proof of concept and developmental stage, one hundred seventeen (117) constructs comprising different combinations of twenty-six (26) distinct promoters, twenty-six (26) distinct introns, and ten (10) distinct insect toxin coding sequences, were used to generate approximately six thousand (6,000) transformed events. After initial molecular characterization for the presence of the transgene(s), five thousand fifty-two (5,052) single and double-copy transformed corn events were selected for further characterization and efficacy testing. These events were evaluated for phenotypic or agronomic off-types, the level of expression of the insect toxin proteins, and efficacy against selected Lepidopteran insect pest species. The resulting efficacy and protein expression data, along with any information regarding phenotypic and agronomic off-types was used to eliminate inefficacious proteins, expression elements and combinations, and was used to design a smaller number of binary commercial transformation plasmid constructs to be used in the next phase of development.

In the next phase of development, eight (8) new constructs were created. These constructs comprised combinations of two (2) to four (4) insect toxin transgene expression cassettes in different orientations (convergent or divergent). These eight (8) constructs were used to generate a total of five thousand seven hundred thirty-three (5,733) transformed events (also referred to as "transformants"). After shoot formation in culture, a subset of the transformed events were selected based upon visual characteristics and early molecular analysis. A total of eight hundred twenty-three (823) transformed events were selected and transplanted to pots and grown for further study.

The resulting  $R_0$  generation transformed events were analyzed for efficacy against selected Lepidopteran species, toxin protein expression, plant health, seed return, and phenotypic and agronomic off-types. The  $R_0$  generation events were also characterized molecularly to ensure cassette intactness and proper insertion in the corn genome. Many of the events were dropped from testing due to failure to pass the agronomic analysis and molecular characterization testing. In addition, one (1) of the eight (8) constructs was dropped from further study at this  $R_0$  stage because it produced events with off-phenotypes. In addition to these agronomic problems, later mode of action ("MOA") studies conducted demonstrated that an insect toxin protein contained in this construct demonstrated an overlapping MOA to a commercially-available protein.

Mode of Action studies were conducted on one of the insect proteins common to four (4) of the eight (8) constructs. These studies demonstrated that this insect protein had an overlapping MOA to a commercially-available protein. Proteins that demonstrate an analogous or overlapping MOA to a currently utilized commercial insecticidal protein are not desirable because of resistance development, which could render a protein with a similar MOA ineffective against insect populations. As such, these four (4) constructs, and the events arising therefrom, were dropped. As noted previously, one (1) of the four (4) dropped constructs also produced events with off-phenotypes at the  $R_0$  stage.

In the next stage of development, one hundred fifty (150) events derived from the remaining four (4) constructs were further evaluated at the  $F_1$  (heterozygous hybrid)/ $R_1$  (homozygous inbred) and  $R_2$  generation for efficacy, seed return

and segregation, phenotypic and agronomic off-types, and further molecular characterization. Two (2) constructs from the remaining four (4) constructs were dropped from further study in this stage for failure to meet one or more of the criteria for advancement, leaving events derived from (2) constructs for further evaluation.

Seventy-seven (77) events, forty-one (41) derived from the construct used to generate event MON 95379 (“Construct MON95”) and thirty-six (36) events derived from the other construct (“Construct 1”), were evaluated as R<sub>2</sub> inbreds and F<sub>1</sub> hybrids for efficacy, seed return and segregation, phenotypic and agronomic off-types and further molecular characterization. Based upon the results of these evaluations, events associated with Construct 1 were de-prioritized, shelved and stored.

Thus, numerous rounds of testing and comparison of various constructs revealed that the transgene cassette provided as SEQ ID NO:13, Construct MON95, was the best option for efficacy against the Lepidopteran pest species Fall Armyworm (FAW, *Spodoptera frugiperda*), Corn Earworm (CEW, *Helicoverpa zea*), Southwestern Corn Borer (SWCB, *Diatraea grandiosella*), Surgarcane Borer (SCB, *Diatraea saccharalis*), and Lesser Cornstalk Borer (LSCB, *Elasmopalpus lignosellus*), with the best molecular characterization and agronomic performance.

Table 2 illustrates the number of transformed events derived (“Plugged”), the number of transformed events selected for growth as R<sub>0</sub> events (“Transplanted”), and the points at which each respective construct was dropped in the evaluation, research and development process that led to the selection of Construct MON95.

TABLE 2

Event construct selection.			
Construct	Plugged	Transplanted	Dropped
Construct MON 95	1202	210	
Construct 1	799	173	Shelved based on R <sub>2</sub> and F <sub>1</sub> data
Construct 2	679	113	R <sub>2</sub> /F <sub>1</sub> /R <sub>1</sub>
Construct 3	1344	20	MOA/R <sub>0</sub>
Construct 4	232	41	R <sub>2</sub> /F <sub>1</sub> /R <sub>1</sub>
Construct 5	544	88	MOA
Construct 6	584	95	MOA
Construct 7	349	83	MOA

Example 2

Field Trials, Molecular Testing and Event Selection

This Example describes the molecular characterization, analysis, and testing in field trials of events created with Construct MON95 in multiple locations over several years, which lead to the selection of the final event, MON 95379.

Table 3 illustrates the process used to select the final event, MON 95379. At the commercial transformation R<sub>0</sub> screen, two hundred ten (210) R<sub>0</sub> transformed events from Construct MON95 were derived and selected for growth. Of the initial two hundred ten (210) selected R<sub>0</sub> transformed events, one hundred forty-seven events (147) were dropped due to concerns regarding efficacy, protein expression, seed return and plant health, or molecular characterization. This left sixty-three (63) events for assay and testing in the next stage of development, the F<sub>1</sub> Screen and the R<sub>1</sub> Nursery stage. In this stage, eleven (11) events were dropped due to

efficacy concerns in the greenhouse testing. Another three (3) events were dropped because of insufficient return of seed from the nursery and/or segregation analysis of the resulting seed. Finally, another five (5) events were removed due to issues discovered in molecular characterization and three (3) events were removed due to issues discovered in molecular southern analysis, leaving forty-one (41) events for assay in the next generation. At the R<sub>2</sub>/F<sub>1</sub> stage of testing, two (2) of the remaining forty-one (41) events were dropped due to issues discovered in further molecular southern characterization, leaving thirty-nine (39) events.

The remaining thirty-nine (39) events were advanced in two different concurrent parallel testing stages: 1) further field trials; and 2) Cre-excision of the selection cassette and the production of gold standard seed. Events were dropped in each of these concurrent parallel testing stages.

During Cre-excision, eleven (11) events were dropped due to issues discovered in molecular characterization after cre-excision of the glyphosate selection cassette. Further, another six (6) events were dropped due to issues discovered in molecular characterization during gold standard seed production.

During the concurrent field testing, based on data collected from the 2016 U.S. Field Trails, another four (4) events were dropped due to efficacy concerns and another twelve (12) events were dropped due to agronomic concerns. Then, based on data collected from the Brazil Field Trials, another event was dropped due to efficacy concerns. Next, bioinformatic analysis conducted during the 2017 U.S. Field Trials resulted in the removal of another three (3) events from further testing, leaving two events: Event 1 and MON 95379. After further analysis of the agronomics of the events from multiple field trials in the U.S., Brazil, Argentina and Puerto Rico, event MON 95379 was selected as the event for commercialization because it ranked higher than Event 1 when all the characteristics of molecular characterization, protein expression, efficacy and agronomics of each event were compared.

TABLE 3

MON 95379 event selection.			
Stage	Assay	Events Removed	Events Remaining 210
Comm. TFN R <sub>0</sub> Screen	Efficacy	21	63
	Expression	6	
R <sub>1</sub> /F <sub>1</sub>	Seed Return/plant health	51	41
	Molecular characterization	69	
	Greenhouse Efficacy	11	
	Molecular characterization	5	
	Nursery return/segregation	3	
R <sub>2</sub> /F <sub>1</sub>	Molecular southern	3	39
	Molecular southern	2	
Cre Excision	Molecular characterization after CP4 excision	11	28
	Molecular characterization	6	
Gold Standard Seed			
U.S. Field 2016	Efficacy	4	6
	Agronomics	12	
Brazil Field	Efficacy	1	5
	Bioinformatic molecular analysis	3	
U.S. Field 2017			2
Commercial Selection	Further analysis of molecular characterization, protein expression, efficacy and agronomics from multiple field trials	1	MON 95379

## Example 3

Cre-Excision of the Glyphosate Selection Cassette  
in Corn Event MON 95379

This Example describes the removal of the glyphosate selection cassette from corn event MON 95379 through in vivo Cre-excision. The glyphosate selection cassette was used to select transformed events. By removal of the selection cassette, a “marker-free” event was created wherein only the insecticidal protein expression cassettes remained in the final event.

FIG. 3 illustrates the breeding process used to generate the marker-free event MON 95379 corn event. Corn variety LH244 immature embryos were transformed using an *Agrobacterium*-mediated transformation process with Construct MON95 (presented as SEQ ID NO:13, and illustrated in FIG. 2). Construct MON95 comprises three (3) expression cassettes: two (2) expression cassettes for the expression of the insecticidal proteins Cry1B.868 and Cry1Da<sub>7</sub>, and a single cassette used for the selection of transformed plant cells using glyphosate selection. The selection cassette was flanked on both sides with LoxP Cre-recombinase recognition sites.

After transformation, the R<sub>0</sub> transformants were self-pollinated for two (2) generations, during which time many events were removed based upon various assays such as efficacy, protein expression, seed return and plant health, and molecular characterization. By the R<sub>2</sub> generation, thirty-nine (39) events remained from the initial two hundred ten (210) events. The thirty-nine (39) homozygous R<sub>2</sub> generation events were bred with an elite line of transformed corn plants expressing Cre-recombinase enzyme, derived from *Enterobacteria* phage P1.

This stage in which R<sub>2</sub> generation events were bred with plants expressing Cre-recombinase is identified as “Cre Cross”. Specifically in this stage, de-tasseled (female) R<sub>2</sub> generation plants homozygous for SEQ ID NO:13 were cross-pollinated with transgenic corn plants (male) homozygous for a transgene cassette used for expression of Cre-recombinase enzyme. The Cre-recombinase expressing male donor pollen germinates after landing on the silk tissue of the female plant comprising SEQ ID NO:13. Once the pollen tube enters the embryo sac, the pollen tube ruptures, setting free the two sperms of the Cre-recombinase expressing male donor. The nucleus of one sperm fuses with the egg nucleus, forming the zygote. The other sperm nucleus fuses with one of the two polar nuclei which in turn fuses with the other polar nucleus, thereby establishing the primary endosperm nucleus.

Thus, in using the Cre-recombinase expressing plant as the male pollen donor, both the embryo and endosperm of the resulting cross will express Cre-recombinase as the cells divide and develop and become a corn kernel (i.e., seed). The Cre-recombinase binds to inverted repeats in the LoxP site and catalyzes a crossover in an eight-base pair spacer region of the two LoxP sites that flank the expression cassette, resulting in the excision of the marker cassette with one LoxP site remaining in the integrated T-DNA due to recombination (see FIG. 2, “Inserted T-DNA After Cre-Excision”).

The F<sub>1</sub> progeny resulting from the Cre Cross were selected for the absence of the CP4 selection cassette and allowed to self-pollinate. Through this process, the two

alleles—the Cre-recombinase allele and the allele for the T-DNA used to generate event MON 95379—segregate in the resulting F<sub>2</sub> population, resulting in progeny homozygous or heterozygous for one or both alleles.

The F<sub>2</sub> progeny which demonstrated the absence of the Cre-recombinase allele and homozygosity for SEQ ID NO:9, the transgenic inserted T-DNA after Cre-excision, were selected. These selected F<sub>2</sub> progeny were self-pollinated, giving rise to an F<sub>3</sub> generation homozygous for SEQ ID NO:9.

A further self-pollination resulted in F<sub>3</sub> progeny seed (F<sub>4</sub> seed) which were assayed for purity, and were designated as “Gold Standard Seed.” F<sub>4</sub> was the first generation of gold standard seed.

Excision of the glyphosate selection marker cassette did not affect the expression of Cry1B.868 and Cry1Da<sub>7</sub>. Removing the glyphosate selection cassette from corn event MON 95379 through Cre-excision provided a transgenic corn event which is resistant to Lepidopteran pests without adding tolerance to glyphosate in the final event. This “marker-free” event assures flexibility when building corn breeding stacks with other corn transgenic events to provide a multiplicity of products incorporating event MON 95379 and allowing multiple options for providing additional traits in the final breeding stacks.

## Example 4

Corn Event MON 95379 Demonstrates Resistance  
to the Lepidopteran Insect Pests Fall Armyworm,  
Corn Earworm, Southwestern Corn Borer,  
Sugarcane Borer

This Example describes the activity of the MON 95379 event against Lepidopteran insect pests. The insect toxin proteins Cry1B.868 and Cry1Da<sub>7</sub>, when expressed together in corn event MON 95379, provide resistant to Fall Armyworm (*Spodoptera frugiperda*), Corn Earworm (*Helicoverpa zea*), Southwestern Corn Borer (*Diatraea grandiosella*), and Sugarcane Borer (*Diatraea saccharalis*).

After transformation and insertion of Construct MON95, forty-one (41) R<sub>0</sub> events were selected for bioassay using leaf discs. Bioassays using plant leaf disks were performed analogous to those described in U.S. Pat. No. 8,344,207. A non-transformed LH244 corn plant was used to obtain tissue to be used as a negative control. Plates comprising wells with one insect per leaf disc in each well were incubated for three (3) days. After three (3) days, the plates were examined. If at least fifty percent (50%) of the leaf disc in the negative controls was consumed, measurements were taken of the transgenic event leaf discs. If less than fifty percent (50%) of the leaf discs in the negative controls had not yet been consumed, the insects were allowed to continue feeding until the fifty percent (50%) target was achieved. Measurements of leaf damage (“leaf damage ratings” or “LDR”) and mortality were taken for each well. An average of each measure was determined. The leaf damage ratings ranged from one (1) to eleven (11) and reflect a percentage of the consumed leaf disc. Table 4 shows the leaf damage rating scale used for the R<sub>0</sub> leaf disc assays. On this rating scale, the negative controls will always have an LDR of at least 10.

TABLE 4

Leaf Damage Ratings (LDR) scale for R <sub>0</sub> leaf disc assays.	
Leaf Damage Rating (LDR)	Amount of feeding damage
1	≤5%
2	≤10%
3	≤15%
4	≤20%
5	≤25%
6	≤30%
7	≤35%
8	≤40%
9	≤45%
10	≤50%
11	>50%

Table 5 shows the mean leaf damage ratings for the forty-one (41) events transformed with Construct MON95, including the MON 95379 event. As can be seen in Table 5, expression of the two insecticidal proteins, Cry1B.868 and Cry1Da\_7, provided resistance to Fall Armyworm (FAW), Corn Earworm (CEW), and Southwestern Corn Borer (SWCB). The LDRs for the negative controls were between 10 and 11. The FAW and SWCB consumed only approximately five percent (5%) of the event MON 95379 leaf disc in comparison to the negative controls which consumed at least fifty percent (50%) of the leaf disc. With respect to CEW, only approximately 6.25% of the leaf discs were consumed in comparison to the negative controls which consumed at least fifty percent (50%) of the leaf disc. In addition, one hundred percent (100%) of the FAW and CEW were killed after consuming the event MON 95379 containing leaf discs.

TABLE 5

Mean Leaf Damage Rating (LDR) scores and Mean Mortality for R <sub>0</sub> plants expressing Cry1B.868 and Cry1Da_7.			
Event	FAW Mean LDR	CEW Mean LDR	SWCB Mean LDR
MON 95379	1.00	1.25	1.00
Event 2	1.00	2.00	1.00
Event 3	1.00	1.00	1.00
Event 4	1.00	1.25	1.00
Event 5	1.00	1.50	1.00
Event 6	1.00	3.25	1.00
Event 7	1.00	2.25	1.00
Event 8	1.00	1.50	1.00
Event 9	1.00	1.75	1.00
Event 10	1.00	2.25	1.25
Event 11	1.00	1.75	1.00
Event 12	1.00	1.25	1.00
Event 13	1.00	1.25	1.00
Event 14	1.00	3.75	1.00
Event 15	1.00	1.00	1.00
Event 16	1.00	1.25	1.25
Event 17	1.00	3.00	1.00
Event 18	1.00	1.25	1.00
Event 19	1.00	1.75	1.25
Event 20	1.00	1.00	1.25
Event 21	1.00	5.00	1.00
Event 22	1.00	1.33	1.00
Event 23	1.00	1.00	1.00
Event 24	1.00	1.50	1.00
Event 25	1.00	1.25	1.00
Event 26	1.00	2.33	1.25
Event 27	1.00	1.25	1.00
Event 28	1.00	1.50	1.00
Event 29	1.00	1.00	1.00

TABLE 5-continued

Mean Leaf Damage Rating (LDR) scores and Mean Mortality for R <sub>0</sub> plants expressing Cry1B.868 and Cry1Da_7.			
Event	FAW Mean LDR	CEW Mean LDR	SWCB Mean LDR
5			
Event 30	1.00	2.00	1.00
Event 31	1.00	2.50	1.50
10			
Event 32	1.00	1.00	1.00
Event 33	1.00	1.67	1.00
Event 34	1.00	3.00	1.00
Event 35	1.00	1.25	1.00
Event 36	1.00	1.50	1.00
Event 37	1.00	2.50	1.00
15			
Event 38	1.00	3.50	1.25
Event 39	1.00	1.00	1.25
Event 40	1.00	1.00	1.00
Event 41	1.00	1.25	1.00

The forty-one (41) events were crossed with non-transgenic 93IDI3 variety plants. F<sub>1</sub> heterozygous progeny plants were selected that comprised Construct MON95. Around five (5) F<sub>1</sub> plants for each event were artificially infested in a greenhouse for each insect pest species. With respect to FAW, approximately forty (40) neonates were used to infest each F<sub>1</sub> plant in the V6 to V8 stage whorl. With respect to SWCB, approximately thirty (30) neonates were used to infest the F<sub>1</sub> plant in the V6 to V10 stage whorl. Measures of leaf damage for FAW and SWCB were taken approximately fourteen (14) days after infestation. Tables 6 and 7 show the damage rating scales used to assess the leaf damage.

TABLE 6

Leaf damage rating scale for corn plants infested with FAW.		
Leaf Damage Rating (LDR)	Description	
0	No visible damage	
1	Only pinhole lesions present on whorl leaves	
40	2	Pinholes and small, circular lesions present on whorl leaves
3	Small, circular lesions and 1-3 small, elongated lesions present on whorl and furl leaves	
4	4-6 small to mid-sized, elongated lesions present on 1-3 whorl and furl leaves	
45	5	4-6 large, elongated lesions present on 1-3 whorl and furl leaves and/or 1-3 small or mid-sized, uniform to irregular lesions, or both eaten from whorl and/or furl leaves
6	4-6 large, elongated lesions present on 4-6 whorl and furl leaves and/or 4-6 large uniform to irregular shaped holes eaten from whorl and furl leaves	
50	7	7+ elongated lesions of all sizes present on several whorl and furl leaves plus 4-6 large uniform to irregular shaped holes eaten from the whorl and furl leaves
8	7+ elongated lesions of all sizes on most whorl and furl leaves plus 7+ mid- to large-sized uniform to irregular-shaped holes eaten from the whorl and furl leaves	
55	9	Whorl and furl leaves almost totally destroyed as well as the plant showing signs of stunting

TABLE 7

Leaf damage rating scale for corn plants infested with SWCB.		
Leaf Damage Rating (LDR)	Description	
60		
65	0	No lesions
	1	Small amount of pin or fine holes on leaves

TABLE 7-continued

Leaf damage rating scale for corn plants infested with SWCB.	
Leaf Damage Rating (LDR)	Description
2	Small amount of shot-hole injury on a few present
3	Shot-hole injury common on several leaves
4	Several leaves with shot-hole and elongated lesions
5	Several leaves with elongated lesions
6	Several leaves with elongated lesions-2.5 cm. Long
7	Long lesions common on ~50% of leaves
8	Long lesions common on ~70% of leaves
9	Most or all leaves with long lesions

The SWCB infested F<sub>1</sub> plants were also assessed for the length of stalk boring caused by SWCB. To determine the length of stalk boring, corn stalks of the corn plants were broken at approximately eye level and the top portion was used to inspect for boring damage. The stalks were split using a double handled knife and the length of the tunnel bored out by SWCB was measured in centimeters (cm). In these experiments, the tunnel length was capped at ten centimeters (10 cm).

In addition, five (5) F<sub>1</sub> plants for each event were also infested with CEW to measure the amount of damage caused by CEW to the corn ear. Approximately forty (40) CEW nymphs were used to infest each plant and were placed on the green silks of R<sub>1</sub> stage plants. Twenty-one (21) days after infestation, the developing ears were examined, and the damage was recorded as cm<sup>2</sup> ear damage.

Table 8 shows the mean leaf damage ratings for the F<sub>1</sub> events infested with FAW and SWCB, the stalk boring lengths caused by SWCB, and the ear damage caused by CEW, wherein "NT" indicates not tested.

TABLE 8

Mean leaf damage ratings of F <sub>1</sub> transgenic corn plants infested with FAW and SWCB, stalk boring lengths caused by SWCB, and ear damage caused by CEW.				
Event	FAW LDR	SWCB LDR	SWCB tunnel length (cm)	CEW ear damage (cm <sup>2</sup> damage)
MON 95379	0.30	1.00	0.00	3.25
Event 2	0.50	1.00	0.00	3.25
Event 3	0.30	1.00	0.00	0.00
Event 4	0.30	1.30	0.00	2.30
Event 5	1.00	1.00	0.00	1.00
Event 6	0.30	1.00	0.00	0.00
Event 7	0.80	0.50	0.00	3.00
Event 8	0.50	1.00	0.00	0.00
Event 9	1.30	0.80	0.00	0.25
Event 10	NT	NT	NT	NT
Event 11	NT	NT	NT	NT
Event 12	0.30	1.00	0.00	3.50
Event 13	NT	NT	NT	NT
Event 14	NT	NT	NT	NT
Event 15	NT	NT	NT	NT
Event 16	NT	NT	NT	NT
Event 17	0.50	NT	NT	NT
Event 18	0.00	1.30	0.00	2.50
Event 19	0.80	1.30	0.00	0.00
Event 20	NT	NT	NT	NT
Event 21	NT	NT	NT	NT
Event 22	NT	NT	NT	NT
Event 23	NT	NT	NT	NT

TABLE 8-continued

Mean leaf damage ratings of F <sub>1</sub> transgenic corn plants infested with FAW and SWCB, stalk boring lengths caused by SWCB, and ear damage caused by CEW.					
Event	FAW LDR	SWCB LDR	SWCB tunnel length (cm)	CEW ear damage (cm <sup>2</sup> damage)	
10	Event 24	0.50	0.50	2.50	2.75
	Event 25	0.00	1.50	2.25	0.00
	Event 26	0.30	1.30	0.00	3.00
	Event 27	0.50	1.50	2.50	1.25
	Event 28	0.30	1.00	1.00	1.75
	Event 29	0.30	1.00	1.00	0.25
15	Event 30	NT	NT	NT	NT
	Event 31	0.30	1.00	0.00	2.50
	Event 32	0.50	1.00	0.00	0.25
	Event 33	0.50	1.00	0.00	2.25
	Event 34	0.30	1.00	0.00	1.50
	Event 35	0.30	1.30	1.25	0.00
20	Event 36	NT	NT	NT	NT
	Event 37	0.50	1.00	0.00	1.50
	Event 38	1.30	1.00	0.00	0.50
	Event 39	0.00	1.00	1.75	2.00
	Event 40	0.30	0.50	1.25	1.75
	Event 41	0.00	0.80	0.00	2.25
25	Negative Control	7.80	9.00	10.00	13.25

As can be seen in Table 8, leaf damage to corn event MON 95379 was minimal for both FAW and SWCB when compared to the negative controls. Essentially, once the insects started to feed on the event MON 95379 F<sub>1</sub> leaf, expression of the Cry1B.868 and Cry1Da\_7 insecticidal proteins in the corn leaves containing event MON 95379 caused the insect to cease consuming the leaf. SWCB tunneling was not observed in event MON 95379 while the negative controls showed extensive tunneling. With respect to CEW ear damage, the damage to the ear was much less compared to the negative control, and was comparable to the ear damage observed in several commercially-available transgenic corn events. Infestation of the magnitude used in the F<sub>1</sub> assays was much higher than what is usually seen in nature. The F<sub>1</sub> assays demonstrated that corn event MON 95379 provides superior control of FAW, SWCB, and CEW.

In the summer of 2016, the F<sub>1</sub> progeny from the remaining thirty-nine (39) events after R<sub>2</sub>/F<sub>1</sub> described in Example 2/Table 3 were assayed for resistance to FAW, CEW, and SWCB in field experiments using artificial infestation. Multiple locations were used to assay resistance.

FAW resistance was assayed in three (3) locations: Jerseyville, Ill.; Thomasboro, Ill.; and Union City, Tenn. In each location, each event was assayed in three (3) field plots using one (1) row per plot and thirty (30) seeds per row. Forty (40) FAW neonates were used to infest each plant twice, at the early and mid-whorl stage (V4 and V7 vegetative stage). Leaf feeding damage ratings were assessed using the scale as provided in Table 6.

SWCB resistance was assayed in three (3) locations: one (1) in Jonesboro, Ark. and two (2) in Union City, Tenn. In each location, each event was assayed in three (3) field plots using one (1) row per plot and thirty (30) seeds per row. Thirty (30) SWCB neonates were used to infest each plant at the mid-whorl stage (V7-V8). At the time of fifty percent (50%) pollen shed, the plants were infested again with thirty (30) SWCB neonates per plant. Stalk tunneling damage was assessed as previously described.

CEW resistance was assayed in five (5) locations: Jerseyville, Ill., Jonesboro, Ark., Monmouth, Ill., Thomasboro, Ill.,

and Union City, Tenn. In each location, each event was assayed in three (3) field plots using one (1) row per plot and thirty (30) seeds per row. Plants were infested when the silks were fresh and green, and some ear formation had started (R1 through R3 stage). CEW egg strips were used for infestation. Each strip contained approximately forty (40) eggs. One (1) strip was placed between the ear and stalk of each plant, with the eggs facing ear and close to the silks. Evaluation of ear damage was determined twenty-one (21) to twenty-eight (28) days after infestations. By this time, the insect has progressed from larval to pupal stage. Damage to the ears was measured as previously described.

For FAW and SWCB, data from all three (3) locations was used. For CEW, due to various field conditions, only data from Jonesboro, Ark. could be used. Table 9 shows the mean FAW leaf damage ratings, the SWCB tunnel lengths, and the CEW ear damage measurements for each of the tested events and the negative control.

TABLE 9

Mean FAW leaf damage ratings, SWCB tunnel length, and CEW ear damage for 2016 field efficacy trials.			
Event	FAW LDR	SWCB Tunnel Length (cm)	CEW Ear Damage (cm <sup>2</sup> )
MON 95379	1.56	0.37	2.86
Event 2	1.28	1.84	4.41
Event 3	1.33	0.14	3.28
Event 4	2.01	0.11	3.00
Event 5	1.83	0.89	4.09
Event 6	1.46	1.09	2.89
Event 7	1.67	0.39	3.40
Event 8	1.44	0.72	3.19
Event 9	1.59	0.00	3.98
Event 10	2.85	0.35	2.96
Event 11	1.50	2.59	2.84
Event 12	1.39	0.42	3.04
Event 13	1.74	0.00	3.34
Event 14	2.52	1.22	3.01
Event 15	1.53	0.22	3.09
Event 16	1.50	1.61	3.24
Event 17	2.31	1.72	2.80
Event 18	1.70	0.13	4.02
Event 19	1.28	1.00	3.96
Event 20	1.28	0.52	2.79
Event 21	1.39	0.39	3.55
Event 22	1.72	0.94	4.34
Event 23	1.86	0.22	3.04
Event 24	1.93	0.00	3.60
Event 25	1.57	0.06	2.28
Event 26	1.65	0.00	2.28
Event 27	1.65	0.28	2.73
Event 28	1.63	2.72	3.60
Event 29	2.62	0.78	3.42
Event 30	4.78	12.59	5.25
Event 31	1.78	1.81	3.79
Event 32	1.63	0.00	3.08
Event 33	1.37	0.00	3.40
Event 34	1.80	0.19	3.20
Event 35	2.20	0.84	3.99
Event 36	1.96	0.22	3.78
Event 37	1.35	0.06	3.07
Event 38	1.22	0.27	3.94
Event 39	1.46	0.14	2.87
Negative Control	7.15	7.21	32.25

As demonstrated in Table 9, corn event MON 95379 provided excellent control of FAW, SWCB, and CEW when compared to the negative control. The level of infestation in these assays was much higher than what would normally be encountered in the field under natural conditions, demonstrating the superior performance of event MON 95379 under high insect pressure.

During concurrent field trials and Cre-excision of the selection cassette and the production of Gold Standard Seed, further characterization of the events was performed. As a result of extensive molecular characterization, efficacy, expression, and agronomic studies, events were dropped from testing, leaving two (2) events: Event 1 and MON 95379. Event 1 was de-prioritized based on observed yield drag in agronomic studies, leaving event MON 95379 for advancement.

During the 2016 to 2017 growing season in Argentina, event MON 95379 was assayed for resistance to FAW, CEW, and SCB in temperate and subtropical regions under natural infestation conditions. FAW leaf damage ratings were determined for event MON 95379 grown in the subtropical region of Argentina using the scale provided in Table 6. SCB tunneling data was obtained for event MON 95379 from two (2) locations in the temperate region of Argentina. CEW ear damage data was obtained for event MON 95379 from two (2) locations in the temperate region and three (3) locations in the subtropical regions of Argentina. Table 11 shows the mean FAW leaf damage ratings, SCB tunnel length, and CEW ear damage under natural infestation conditions for event MON 95379 and a negative control during the 2016-2017 Argentina growing seasons.

TABLE 10

Mean FAW leaf damage ratings, SCB tunnel length, and CEW ear damage for 2016-2017 Argentina field efficacy trials.			
Event	FAW (LDR)	SCB Tunnel Length (cm)	CEW Ear Damage (cm <sup>2</sup> )
MON 95379	1.27	0.00	1.32
Negative Control	7.48	4.43	5.83

As can be seen in Table 10, event MON 95379 provided resistance to FAW, SCB, and CEW when compared to the negative control under natural infestation conditions in Argentina.

Event MON 95379 was also evaluated for resistance against FAW resistant to a commercially-available corn event (MON89034, which expresses Cry1a.105 and Cry2Ab2) over three (3) growing seasons in Puerto Rico (January 2016, July 2016, and January 2017). Table 11 shows the mean leaf damage ratings based upon the scale presented in Table 6 for each of the three (3) growing seasons compared with event MON89034 and the negative control.

TABLE 11

Mean leaf damage ratings for event MON 95379 and event MON89034 naturally-infested with event MON89034-resistant FAW.			
Event	January 2016	July 2016	January 2017
MON 95379	2.30	2.22	1.40
MON89034	5.68	4.54	7.36
Negative Control	8.73	6.84	9.00

As can be seen in Table 11, corn event MON 95379 demonstrated resistance to event MON89034-resistant FAW under high natural pressure relative to the negative control.

In the summer of 2017, event MON 95379 was evaluated for resistance against FAW, SWCB, and CEW in the United States using methods similar to that described for the summer of 2016. FAW resistance was assayed at three (3)

locations: Jerseyville, Ill.; Thomasboro, Ill.; and Monmouth, Ill. In each location, each event was assayed in three (3) field plots using one (1) row per plot and thirty (30) seeds per row. Forty (40) FAW neonates were used to infest each plant two times. The first infestation occurred around V5 stage. The second infestation for plants in Monmouth, Ill. and Jerseyville, Ill. occurred around V8 stage. Due to a low hatch rate and poor weather, a second infestation was not possible in Thomasboro, Ill. FAW leaf feeding damage ratings were assessed using the scale as provided in Table 6.

SWCB resistance was assayed at three (3) locations, one (1) in Jonesboro, Ark. and two (2) in Union City, Ill. In each location, each event was assayed in three (3) field plots using one (1) row per plot and thirty (30) seeds per row. Thirty (30) SWCB neonates were used to infest each plant two times. Under normal conditions, the first infest is performed at the mid-whorl stage (V7-V8) in half of the row, but infestation was delayed about a week. Regardless, strong insect pressure was established. At the time of fifty percent (50%) pollen shed the second half of the row of plants were infested with thirty (30) SWCB neonates per plant. Stalk tunneling damage was assessed as previously described.

CEW resistance was assayed at six (6) locations: Jerseyville, Ill., Jonesboro, Ark., Paragould, Ark., Monmouth, Ill., and two locations in Union City, Tenn. In each location, each event was assayed in three (3) field plots using one (1) row per plot and thirty (30) seeds per row. Due to a shortage of insects, infestations in Monmouth, Ill. and Jerseyville, Ill. were infested two (2) to three (3) weeks later than when silks are fresh and green. In Monmouth, approximately twenty-two (22) neonates were used to infest each plant. In Jerseyville, Ill., twenty-three (23) to twenty-four (24) neonates were used to infest partially opened corn ears. In Jonesboro, Ark., one (1) of the three (3) rows received approximately thirty (30) neonates per plant, and the other two (2) rows received sixteen (16) to (18) neonates per plant. In Paragould, Ark., all three (3) rows received approximately thirty (30) neonates per plant. Infestation was delayed in the two locations of Union City, Tenn. due to insect availability. Both locations received eighteen (18) neonates per plant. Evaluation of ear damage was determined twenty-one (21) to twenty-eight (28) days after infestations. Damage to the ears was expressed as previously described. Artificial infestations were conducted on both marker and marker-free event MON 95379 plants. In addition, assays were also conducted using the natural insect pressure at the locations for the marker-containing event MON 95379 plants. Tables 12 and 13 show the FAW leaf damage ratings, SWCB tunnel lengths, and the CEW ear damage for marker-containing and marker-free event MON 95379 plants.

TABLE 12

Mean FAW leaf damage ratings, SWCB tunnel length, and CEW ear damage for event MON 95379 plants before Cre-excision of the selection marker under conditions of artificial and natural infestation.					
Before Cre-excision of CP4 Marker					
Event	FAW Artificial Infestation (LDR)	FAW Natural Infestation (LDR)	CEW Artificial Infestation (cm <sup>2</sup> )	CEW Natural Infestation (cm <sup>2</sup> )	SWCB Artificial Infestation Tunnel Length (cm)
MON 95379	1.17	1.15	4.93	5.81	0.00
Negative Control	7.08	8.15	8.49	14.57	14.34

TABLE 13

Mean FAW leaf damage ratings, SWCB tunnel length, and CEW ear damage for marker-free event MON 95379 plants under artificial infestation.			
After Cre-excision of CP4 Marker			
Event	FAW Artificial Infestation (EDR)	CEW Artificial Infestation (cm <sup>2</sup> )	SWCB Artificial Infestation Tunnel Length (cm)
MON 95379	1.20	5.39	0.27
Negative Control	7.08	8.49	14.34

As can be seen in Tables 12 and 13, event MON 95379 provided resistance against FAW, SWCB, and CEW under artificial (marker and marker-free) and natural (marker-free) infestation conditions.

In 2018, a hybrid cross of event MON 95379 with event MON89034 was assayed for resistance to FAW in a Brazil field trial under natural infestation conditions. The field trial was conducted in Santa Helena de Goiás, State of Goiás. In this location there are FAW populations resistant to the transgenic corn event MON89034. Transgenic corn plants corresponding to the cross of events MON 95379×MON89034, event MON89034, and a conventional corn plant (negative control) were planted. At V6 stage, leaf damage rating scores were determined for sixty (60) plants corresponding to the cross of events MON 95379×MON89034, thirty (30) plants corresponding to event MON89034, and thirty (30) negative controls using the scale presented in Table 6. In addition, the number of FAW neonates, larvae greater than two millimeters (2 mm) and less than or equal to 1.5 centimeters, and larvae greater than 1.5 centimeters were recorded for each plant. Table 14 shows the mean leaf damage ratings for the cross of events MON 95379×MON89034, event MON89034, and the negative control, along with the numbers of neonates and larvae observed on the corn plants.

TABLE 14

Mean FAW leaf damage rating and number of neonates and larvae from Brazil, 2018 field trials for the cross of events MON 95379 × MON89034, event MON89034 and negative control.				
Event	FAW (LDR)	Neonates	Larvae >2 mm and ≤1.5 cm	Larvae >1.5 cm
MON 95379 × MON89034	0.62	0	0	0
MON 89034	1.47	6	16	0
Negative Control	5.20	0	22	9

As can be seen in Table 14, the cross of events MON 95379×MON89034 provided resistance to FAW under natural infestation conditions relative to the negative control. The cross of events MON 95379×MON89034 also performed better than event MON89034 under conditions where event MON89034-resistant FAW are within the population of FAW. With respect to neonates and larvae, none were observed on the plants corresponding to the cross of events MON 95379×MON89034. Neonates and larvae between two (2) millimeters and one and a half (1.5)

centimeters were observed on event MON89034 plants. The negative control plants were observed to have even more larvae than event MON89034 plants, and had larvae that had grown greater than 1.5 centimeters.

Example 5

Assay of Activity of Corn Event MON 95379 Against Lesser Cornstalk Borer

This Example describes the assay of activity of transgenic corn event MON 95379 against the Lepidopteran insect pest, Lesser Cornstalk Borer (LSCB, *Elasmopalpus lignosellus*).

Event MON 95379 was grown in a greenhouse along with negative control plants and infested with LSCB neonates. Ten (10) event MON 95379 plants and nine (9) negative control plants were grown in individual pots. Nine (9) days after planting, each plant was infested with ten (10) LSCB neonates per plant. Twenty-two (22) days after infestation, the plants were examined and rated for damage using a 0-4 damage rating scale as presented in Table 15.

TABLE 15

LSCB plant damage rating scale.		
LSCB Damage Rating	Severity of Injury	Description
0	No damage	Plants without injury
1	Slight injury	Plants with scratches in leaves and/or stalk
2	Average damage	Plants with stalk parts damaged
3	Serious damage	Plants with stalk parts damage and dead hear symptoms
4	Dead plant	Dead plants

The resulting LSCB damage ratings for each plant is presented in Table 16.

TABLE 16

LSCB plant damage for each infested plant.			
Plant	MON 95379	Negative Control	
1	0	4	
2	0	2	
3	0	3	
4	0	3	
5	0	4	
6	0	3	
7	0	3	
8	0	4	
9	0	4	
10	0		

As can be seen in Table 16, LSCB produced extensive damage to the negative control plants, four (4) of which were rated as "Dead," four (4) of which were rated as "Serious damage," and only one (1) rated as "Average damage." In contrast, the event MON 95379 LSCB infested plants showed no damage.

Transgenic corn event MON 95379 provides resistance to Lesser Cornstalk Borer (LSCB, *Elasmopalpus lignosellus*).

Example 6

Corn Event MON 95379 Provides Consistent Yield and Similar Agronomics to Untransformed LH244 Corn Plants

This Example demonstrates that transgenic corn event MON 95379 provides consistent yields and similar agronomics in the field to untransformed LH244 corn plants.

Field trials were conducted with plants corresponding to event MON 95379 prior to Cre-excision of the glyphosate selection cassette to determine various aspects of yield and agronomics in comparison to control plants. Measurements of yield were calculated and expressed as bushels per acre (bu/acre). Plant height and ear height were measured in inches (in). Fifty percent (50%) pollen shed and fifty percent (50%) silking were expressed as days after planting (DAP). Test weight, which is a measurement of bulk density, or the weight of a unit volume, of grain was expressed in pounds per bushel (lb/bu). The USDA established the standard test weight of a bushel of corn as fifty-six pounds per bushel (56 lb/bu) based upon a 15.5% moisture content. The percent moisture of the corn kernel was expressed on a wet weight basis. The moisture content is the amount of water in the seed and is usually expressed as a percentage. It can be expressed on either a wet weight basis (where it is expressed as a percentage of the fresh weight of the seed) or on a dry weight basis (where it is expressed as a percentage of the dry weight of the seed). Determination of percent moisture is destructive to the seed. Percent moisture (wet basis) can be calculated with the simple formula:

$$M_{wb} = (W_w / W_w + W_d) \times 100$$

Where  $W_w$  is equal to the weight of the water and  $W_d$  is equal to the weight of the dry matter.

In the growing season of 2016 in the United States, yield and agronomic measures were determined for event MON 95379 inbreds and hybrids pre-Cre-excision of the glyphosate maker cassette. Tables 17 and 18 show the yield and agronomic characteristics measured for event MON 95379 inbreds and hybrids, respectively. The negative control plants for the inbred comparisons was untransformed variety LH244. Hybrids containing event MON 95379 were created by cross pollinating the inbred event MON 95379 with corn variety 93IDI3, and the untransformed control was an LH244x93IDI3 cross.

TABLE 17

Yield and agronomics for event MON 95379 inbreds relative to non-transgenic controls.										
Event	Yield (bu/acre)		Plant Height (in)		Ear Height (in)		50% Pollen Shed (DAP)		50% Silking (DAP)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
MON 95379	111.08	5.17	85.40	2.72	41.29	1.45	65.11	2.00	65.77	2.05
LH244	114.18	5.02	82.05	2.32	38.73	1.33	62.60	1.79	63.54	1.82

45

TABLE 18

Yield and agronomic for event MON 95379 hybrids relative to non-transgenic controls.						
Event	Yield (bu/acre)		Test Weight (lb/bu)		Percent Moisture	
	Mean	SE	Mean	SE	Mean	SE
MON 95379 × 93IDI3	203.48	7.04	58.78	0.61	16.04	0.42
LH244 × 93IDI3	196.45	6.81	58.31	0.61	15.89	0.43

As can be seen in Tables 17 and 18, the yield and other agronomic measures for event MON 95379 in the 2016 United States field trials were relatively the same for both inbreds and hybrids relative to the controls. The variability between the inbreds and hybrids and their respective controls was within acceptable limits and demonstrate there were no negative impacts on yield and other agronomic characteristics caused by insertion of the T-DNA into the corn genome of event MON 95379.

Yield and agronomics were also studied in Argentina during the 2016 to 2017 growing season for event MON 95379 inbreds and hybrids pre-Cre-excision of the glyphosate marker cassette. Tables 19 and 20 show the yield and agronomic characteristics measured for event MON 95379 inbreds. The negative control plants for the inbred comparisons was untransformed variety LH244. Hybrids containing event MON 95379 were created by cross-pollinating events MON89034×MON895379. The transgenic control was event MON88017×event MON89034. The non-transgenic control was a LH244×93IDI3 cross. Table 21 shows the yield and agronomic characteristics measured for event MON 95379 hybrids, wherein “NC” indicates not calculated.

TABLE 19

Yield and agronomics for event MON 95379 inbreds relative to non-transgenic controls.										
Event	Yield (bu/acre)		Plant Height (in)		Ear Height (in)		50% Pollen Shed (DAP)		50% Silking (DAP)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
MON 95379	113.39	4.57	73.89	1.05	38.38	0.95	64.32	0.60	64.68	0.62
LH244	105.49	NC	71.69	NC	36.66	NC	64.86	NC	65.17	NC

TABLE 20

Yield and agronomics for event MON 95379 inbreds relative to non-transgenic controls.				
Event	Test Weight (lb/bu)		Percent Moisture	
	Mean	SE	Mean	SE
MON 95379	59.28	0.10	16.54	0.26
LH244	59.32	NC	16.01	NC

46

TABLE 21

Yield and agronomics for event MON 95379 hybrids relative to transgenic and non-transgenic controls.							
Event	Yield (bu/acre)		Test Weight (lb/bu)		Percent Moisture		
	Mean	SE	Mean	SE	Mean	SE	
MON89034 × MON 95379	165.80	5.94	60.06	0.20	15.12	0.18	
MON88017 × MON89034	164.87	5.94	59.62	0.20	15.45	0.18	
LH244 × 93IDI3	166.82	NC	59.70	NC	15.38	NC	

As can be seen in Tables 19 through 21, the measures of yield and other agronomic characteristics were relatively the same for event MON 95379 inbreds and hybrids relative to the controls.

In 2017, yield and agronomics were again measured in field trials in the United States for event MON 95379 inbreds and hybrids pre-Cre-excision of the glyphosate marker cassette. Inbred and hybrid controls were similar to those used in the 2016 United States field trials. Table 22 shows the yield and agronomic characteristics for event MON 95379 inbreds relative to non-transgenic controls, and Tables 23 and 24 show the yield and agronomic characteristics measured for event MON 95379 hybrids relative to non-transgenic controls in the 2017 United States field trials.

TABLE 22

Yield and agronomics for event MON 95379 inbreds relative to non-transgenic controls.							
Event	Yield (bu/acre)		Test Weight (lb/bu)		Percent Moisture		
	Mean	SE	Mean	SE	Mean	SE	
MON 95379	116.49	5.45	57.86	0.59	21.16	0.76	
LH244	124.46	5.39	58.60	0.58	20.57	0.76	

TABLE 23

Yield and agronomic for event MON 95379 hybrids  
relative to non-transgenic controls.

Event	Yield		Plant Height		Ear Height		50% Pollen Shed		50% Silking	
	(bu/acre)		(in)		(in)		(DAP)		(DAP)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
MON 95379 × 93IDI3	213.15	4.84	99.14	2.50	45.27	1.77	56.65	0.93	57.17	1.06
LH244 × 93IDI3	217.86	4.44	97.42	2.28	45.61	1.45	55.87	0.91	56.47	1.04

TABLE 24

Yield and agronomic for event MON 95379 hybrids relative  
to non-transgenic controls.

Event	Test Weight (lb/bu)		Percent Moisture	
	Mean	SE	Mean	SE
MON 95379 × 93IDI3	57.98	0.37	19.73	0.51
LH244 × 93IDI3	57.53	0.36	19.74	0.50

20

TABLE 26

Yield and agronomics for event MON 95379 inbreds relative  
to non-transgenic controls.

Event	Test Weight (lb/bu)		Percent Moisture	
	Mean	SE	Mean	SE
MON 95379	59.59	1.52	17.49	0.96
LH244	60.34	0.97	17.24	0.69

25

30

As can be seen in Tables 22 through 24, the yield and other agronomic properties event MON 95379 demonstrated in the 2017 United States field trials were similar to the untransformed controls for both inbred and hybrid lines.

During the 2018 to 2019 growing season in Argentina, agronomics and yield were measured in field trials for event MON 95379 inbreds and hybrids post-Cre-excision of the glyphosate marker cassette. Inbred controls were similar to those used in the 2017 United States field trials. The hybrids were produced through crosses with the elite variety 80IDM2. Tables 25 and 26 show the yield and agronomic characteristics for event MON 95379 inbreds relative to non-transgenic controls, and Table 27 shows the yield and agronomic characteristics measured for event MON 95379 hybrids relative to non-transgenic controls in the 2018 to 2019 Argentina field trials.

35

40

45

TABLE 27

Yield and agronomics for event MON 95379 hybrids  
relative to non-transgenic controls.

Event	Yield (bu/acre)		Test Weight (lb/bu)		Percent Moisture	
	Mean	SE	Mean	SE	Mean	SE
MON 95379 × 80IDM2	206.86	8.37	59.42	0.46	19.03	0.84
LH244 × 80IDM2	207.81	8.11	59.22	0.45	19.12	0.83

TABLE 25

Yield and agronomics for event MON 95379 inbreds  
relative to non-transgenic controls.

Event	Yield		Plant Height		Ear Height		50% Pollen Shed		50% Silking	
	(bu/acre)		(in)		(in)		(DAP)		(DAP)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
MON 95379	92.81	7.30	85.14	2.45	40.57	2.06	62.74	0.61	63.77	0.65
LH244	103.46	7.68	83.93	1.72	38.57	1.24	62.93	0.61	63.88	0.71

As can be seen in Tables 25 through 27, the yield and other agronomic properties event MON 95379 demonstrated in the 2017 to 2018 Argentina field trials were similar to the untransformed controls for both inbred and hybrid lines.

Thus, in sum, corn event MON 95379 demonstrated similar yield and other agronomic properties over four (4) separate growing seasons in the United States and Argentina. Event MON 95379 does not negatively affect yield or cause a change in other agronomic properties measured compared to non-transgenic and transgenic controls.

Example 7

Corn Event MON 95379 Event-Specific Endpoint TAQMAN® Assays

The following Example describes methods useful in identifying the presence of event MON 95379 in a corn sample. A pair of PCR primers and a probe were designed for the purpose of identifying the unique junction formed between the corn genomic DNA and the inserted DNA of event MON 95379 in an event-specific endpoint TAQMAN® PCR. Examples of conditions utilized for identifying the presence of event MON 95379 in a corn sample in an event-specific endpoint TAQMAN® PCR are described in Table 28 and Table 29.

The sequence of the oligonucleotide forward primer SQ21529 (SEQ ID NO:15) is identical to the nucleotide sequence corresponding to positions 833-852 of SEQ ID NO:10. The sequence of the oligonucleotide reverse primer SQ21524 (SEQ ID NO:16) is identical to the reverse complement of the nucleotide sequence corresponding to positions 905-934 of SEQ ID NO:10. The sequence of the oligonucleotide probe PB10269 (SEQ ID NO:17) is identical to the reverse complement of the nucleotide sequence corresponding to positions 886-901 of SEQ ID NO:10. The primers SQ21529 (SEQ ID NO:15) and SQ21524 (SEQ ID NO:16) with probe PB10269 (SEQ ID NO:17), which may be fluorescently labeled (e.g., a 6-FAM™ fluorescent label), can be used in an endpoint TAQMAN® PCR assay to identify the presence of DNA derived from event MON 95379 in a sample.

In addition to SQ21529 (SEQ ID NO:15), SQ21524 (SEQ ID NO:16), and PB10269 (SEQ ID NO:17), it should be apparent to persons skilled in the art that other primers and/or probes can be designed to either amplify or hybridize to sequences within SEQ ID NO:10 which are unique to, and useful for, detecting the presence of DNA derived from event MON 95379 in a sample.

Following standard molecular biology laboratory practices, PCR assays for event identification were developed for detection of event MON 95379 in a sample. Parameters of either a standard PCR assay or a TAQMAN® PCR assay were optimized with each set of primer pairs and probes (e.g., probes labeled with a fluorescent tag such as 6-FAM™) used to detect the presence of DNA derived from event MON 95379 in a sample. A control for the PCR reaction includes internal control primers and an internal control probe (e.g., VIC®-labeled) specific to a region within the corn genome that is used as an internal control, and are primers SQ20222 (SEQ ID NO:18), SQ20221 (SEQ ID NO:19), and VIC® labeled probe PB50237 (SEQ ID NO:20).

Generally, the parameters which were optimized for detection of event MON 95379 in a sample included primer and probe concentration, amount of templated DNA, and PCR amplification cycling parameters. The controls for this

analysis include a positive control from corn containing event MON 95379, a negative control from non-transgenic corn, and a negative control that contains no template DNA.

TABLE 28

MON 95379 event-specific endpoint TAQMAN® PCR reaction components.					
Step	Reagent	Stock Concentration (µM)	Volume (µl)	Final Concentration (µM)	Comments
1	Reaction volume		5		
	2x Master Mix		2.28		1x final concentration
2	Event Specific Primer SQ51219	100	0.05	0.9	
3	Event Specific Primer SQ21524	100	0.05	0.9	
4	Event Specific 6FAM™ probe PB10269	100	0.01	0.2	Probe is light sensitive
5	Internal Control Primer SQ20222	100	0.05	0.9	
6	Internal Control Primer SQ20221	100	0.05	0.9	
7	Internal Control VIC® probe PB50237	100	0.01	0.2	Probe is light sensitive
8	Extracted DNA (template): Leaf Samples to be analyzed		2.5		Separate reactions are made for each template.
	Negative control (non-transgenic DNA)				
	Negative water control (No template control)				
	Positive Qualitative control(s) MON 95379 DNA				

TABLE 29

Endpoint TAQMAN® thermocycler conditions.		
Step No.	Number of Cycles	Settings
1	1	95° C. 20 seconds
2	40	95° C. 3 seconds 60° C. 20 seconds
3	1	10° C.

Example 8

Assays for Determining Zygosity for Event MON 95379 Using TAQMAN® and Detection of the Insect Toxin Transgenes

The following Example describes methods useful in identifying the zygosity of event MON 95379 and detection of the insect toxin transgenes in event MON 95379 in a corn sample. Pairs of PCR primers and a probe are designed for the purpose of identifying specific properties of alleles positive and negative for the T-DNA insertion that gave rise to event MON 95379.

A zygosity assay is useful for determining if a plant comprising an event is homozygous for the event DNA (i.e., comprising the exogenous DNA in the same location on

51

each chromosome of the chromosomal pair), heterozygous for the event DNA (i.e., comprising the exogenous DNA on only one chromosome of the chromosomal pair), or wild-type (i.e., null for the event DNA).

An endpoint TAQMAN® thermal amplification method was used to develop a zygosity assay for event MON 95379. The assay uses a primer pair and a probe to detect amplicons corresponding to one of the two insect toxin coding sequences encoding Cry1B.868 and Cry1Da\_7 comprised within the T-DNA used to generate corn event MON 95379. In addition, a primer pair and probe are used to detect a single-copy internal control that is located within the corn genome and is known to be present as a homozygous allele.

For this assay two (2) primer pairs and two (2) probes were mixed together with the sample. The DNA primers used in the zygosity assay which detects the presence of the Cry1B.868 toxin coding sequence were primers SQ50998 (SEQ ID NO:21) and SQ50997 (SEQ ID NO:22). The VIC®-labeled DNA probe used in the zygosity assay which detects the presence of the Cry1B.868 toxin coding sequence was PB54340 (SEQ ID NO:23). The DNA primers used in the zygosity assay which detect the presence of the Cry1Da\_7 toxin coding sequence were primers SQ50485 (SEQ ID NO:24) and SQ50484 (SEQ ID NO:25). The VIC®-labeled DNA probe used in the zygosity assay which detects the presence of the Cry1Da\_7 toxin coding sequence was PB50138 (SEQ ID NO:26). Both zygosity detection assays use the same internal control. The primers for the internal control were SQ20222 (SEQ ID NO:18) and SQ20221 (SEQ ID NO:19), and the 6FAM™-labeled probe for the internal control was PB50237 (SEQ ID NO:20). The DNA primers and probe for either Cry1B.868 or Cry1Da\_7 were mixed with the primers and probe for the internal control as shown in Tables 30 and 31.

TABLE 30

Corn event MON 95379 zygosity TAQMAN® PCR for the detection of Cry1B.868.					
Step	Reagent	Stock Concentration (µl)	Volume (µl)	Final Concentration (µM)	Comments
1	Reaction volume 2x Master Mix		5 2.4		1x final concentration
2	Cry1B.868 specific primer SQ50998	100	0.0225	0.45	
3	Cry1B.868 specific primer SQ50997	100	0.0225	0.45	
4	Cry1B.868 6FAM™ probe PB50340	100	0.005	0.1	Probe is light sensitive
5	Internal Control Primer SQ20222	100	0.0225	0.45	
6	Internal Control Primer SQ20221	100	0.0225	0.45	
7	Internal Control VIC® probe PB50237	100	0.005	0.1	Probe is light sensitive
8	Extracted DNA (template): Leaf Samples to be analyzed Negative control (non-transgenic DNA) Negative water control (No template control)		2.5		Separate reactions are made for each template.

52

TABLE 30-continued

Corn event MON 95379 zygosity TAQMAN® PCR for the detection of Cry1B.868.					
Step	Reagent	Stock Concentration (µl)	Volume (µl)	Final Concentration (µM)	Comments
	Positive copy number control(s) Cry1B.868				

TABLE 31

Corn event MON 95379 zygosity TAQMAN® PCR for the detection of Cry1Da_7.					
Step	Reagent	Stock Concentration (µl)	Volume (µl)	Final Concentration (µM)	Comments
1	Reaction volume 2x Master Mix		5 2.4		1x final concentration
2	Cry1Da_7 specific Primer SQ50485	100	0.0225	0.45	
3	Cry1Da_7 specific Primer SQ50484	100	0.0225	0.45	
4	Cry1Da_7 specific 6FAM™ probe PB50138	100	0.005	0.1	Probe is light sensitive
5	Internal Control Primer SQ20222	100	0.0225	0.45	
6	Internal Control Primer SQ20221	100	0.0225	0.45	
7	Internal Control VIC® probe PB50138	100	0.005	0.1	Probe is light sensitive
8	Extracted DNA (template): Leaf Samples to be analyzed Negative control (non-transgenic DNA) Negative water control (No template control) Positive copy number control(s) Cry1Da_7		2.5		Separate reactions are made for each template.

Separate reactions are mixed using DNA derived from a leaf sample for which zygosity is not known, a negative control of DNA derived from an untransformed corn plant, a negative control lacking DNA, and a positive control using DNA derived from a transgenic plant homozygous for Cry1B.868 or Cry1Da\_7, depending upon which toxin coding sequence is used for detection. The reactions are then subjected to the thermal cycles presented in Table 32.

TABLE 32

Zygosity TAQMAN® Thermocycler conditions.		
Step No.	Number of Cycles	Settings
1	1	95° C. 20 seconds
2	40	95° C. 3 seconds 60° C. 20 seconds
3	1	10° C.

After amplification, the cycle thresholds (Ct values) were determined for the amplicon corresponding to the toxin coding sequence and the single-copy, homozygous internal standard. The difference ( $\Delta$ Ct) between the Ct value of the single-copy, homozygous internal standard amplicon and the Ct value of the toxin coding sequence amplicon was determined. With respect to zygosity, a  $\Delta$ Ct of around zero (0) indicated homozygosity of the inserted event MON 95379 T-DNA and  $\Delta$ Ct of around one (1) indicated heterozygosity of the inserted event MON 95379 T-DNA. Lack of an amplicon corresponding to the insect toxin coding sequence indicated the sample is null for the inserted event MON 95379 T-DNA. The Ct values in the TAQMAN® thermal amplification method will have some variability due to multiple factors such as amplification efficiency and ideal annealing temperatures. Therefore, the range of “about one (1)” is defined as a  $\Delta$ Ct of 0.75 to 1.25.

For each progeny derived from a cross with event MON 95379, assays were performed for both toxin coding sequences to assure accuracy in the determination of zygosity of the progeny.

#### Example 9

##### Assays for Determining Zygosity for Corn Event MON 95379 Using TAQMAN®

The following Example describes a method useful in identifying the zygosity of event MON 95379 in a corn sample.

Pairs of PCR primers and a probe are designed for the purpose of identifying specific properties of alleles positive and negative for the T-DNA insertion that gave rise to event MON 95379. Examples of conditions that may be used in an event-specific zygosity TAQMAN® PCR are provide in Tables 33 and 34. For this assay, three primers and two probes were mixed together with the sample. The DNA primers used in the zygosity assay were primers SQ50219 (SEQ ID NO:15), SQ21524 (SEQ ID NO:16), and PWTDNA (SEQ ID NO:27). The probes used in the zygosity assay were 6FAM™-labeled probe PB10269 (SEQ ID NO:17) and VIC®-labeled probe PRWTDNA (SEQ ID NO:28). Primers SQ50219 (SEQ ID NO:15) and SQ21524 (SEQ ID NO:16) and the 6FAM™-labeled probe PB10269 (SEQ ID NO:17) are diagnostic for event MON 95379 DNA. SQ50219 (SEQ ID NO:15) and PWTDNA (SEQ ID NO:27) and the VIC®-labeled probe PRWTDNA (SEQ ID NO:28) are diagnostic when there is no copy of event MON 95379; i.e., they are diagnostic for the wild type allele.

When the three primers and two probes are mixed together in a PCR reaction with DNA extracted from a plant heterozygous for event MON 95379, there is a fluorescent signal from both the 6FAM™-labeled probe PB10269 (SEQ ID NO:17) and the VIC®-labeled probe PRWTDNA (SEQ ID NO:28) which is indicative of and diagnostic for a plant heterozygous for event MON 95379. When the three primers and two probes are mixed together in a PCR reaction with DNA extracted from a plant homozygous for event MON 95379, there is a fluorescent signal from only the 6FAM™-labeled probe PB10269 (SEQ ID NO:17) and not the VIC®-labeled probe PRWTDNA (SEQ ID NO:28). When the three primers and the two probes are mixed together in a PCR reaction with DNA extracted from a plant which is null for event MON 95379 (i.e., the wild-type), there is a fluorescent signal from only the VIC®-labeled probe PRWTDNA (SEQ ID NO:28). The template DNA samples and controls for this analysis were a positive control from corn containing event

MON 95379 DNA (from both a known homozygous and a known heterozygous sample), a negative control from non-transgenic corn and a negative control that contains no template DNA.

TABLE 33

Event MON 95379 zygosity TAQMAN® PCR					
Step	Reagent	Stock Concentration (μl)	Volume (μl)	Final Concentration (μM)	Comments
1	Reaction volume 18 megohm water		5 0.33		Adjust for final volume 1× final concentration
2	2× Master Mix		2.5		
3	Event Specific Primer SQ51219	100	0.05	0.9	
4	Event Specific Primer SQ21524	100	0.05	0.9	
5	Event Specific 6FAM™ probe PB10269	100	0.01	0.2	Probe is light sensitive
6	WT allele Primer PNEG DNA	100	0.05	0.9	
7	WT allele VIC® probe PRBNEG DNA	100	0.01	0.2	Probe is light sensitive
8	Extracted DNA (template): Leaf Samples to be analyzed Negative control (non-transgenic DNA) Negative water control (No template control) Positive Qualitative control(s) MON 95379 DNA		2.5		Separate reactions are made for each template.

TABLE 34

Zygosity TAQMAN® thermocycler conditions		
Step No.	Number of Cycles	Settings
1	1	95° C. 20 seconds
2	40	95° C. 3 seconds 60° C. 20 seconds
3	1	10° C.

#### Example 10

##### Identification of Corn Event MON 95379 in any MON 95379 Breeding Event

The following Example describes how one may identify the MON 95379 event within progeny of any breeding activity using corn event MON 95379.

DNA primer pairs are used to produce an amplicon diagnostic for corn event MON 95379. An amplicon diagnostic for event MON 95379 comprises at least one junction sequence. The junction sequences for event MON 95379 are SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8 ([1], [2], [3], [4], [5], [6], [7], and [8], respectively in FIG. 1). SEQ ID NO:1 is a fifty (50) nucleotide sequence

representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:1 is positioned in SEQ ID NO:10 at nucleotide position 838-887. SEQ ID NO:2 is a fifty (50) nucleotide sequence representing the 3' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:2 is positioned in SEQ ID NO:10 at nucleotide position 14156-14205. SEQ ID NO:3 is a one hundred (100) nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:3 is positioned in SEQ ID NO:10 at nucleotide position 813-912. SEQ ID NO:4 is a one hundred (100) nucleotide sequence representing the 3' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:4 is positioned in SEQ ID NO:10 at nucleotide position 14,131-14,230. SEQ ID NO:5 is a two hundred (200) nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:5 is positioned in SEQ ID NO:10 at nucleotide position 763-962. SEQ ID NO:6 is a two hundred (200) nucleotide sequence representing the 3' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:6 is positioned in SEQ ID NO:10 at nucleotide position 14,081-14,280. SEQ ID NO:7 is a one thousand one hundred sixty (1,160) nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette. SEQ ID NO:7 is positioned in SEQ ID NO:10 at nucleotide positions 1-1,160. SEQ ID NO:8 is a one thousand one hundred seventy eight (1,178) nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA. SEQ ID NO:8 is positioned in SEQ ID NO:10 at nucleotide positions 14,039-15,216.

Primer pairs that will produce an amplicon diagnostic for event MON 95379 include primer pairs based upon the flanking sequences (SEQ ID NO:11 and SEQ ID NO:12) and the inserted T-DNA (SEQ ID NO:9). To acquire a diagnostic amplicon in which SEQ ID NO:1, or SEQ ID NO:3, or SEQ ID NO:5, or SEQ ID NO:7 is found, one would design a forward primer molecule based upon the 5' flanking corn genomic DNA (SEQ ID NO:11; from bases 1 to 862 of SEQ ID NO:10) and a reverse primer molecule based upon the inserted T-DNA (SEQ ID NO:9; from positions 863 through 14,180 of SEQ ID NO:10) in which the primer molecules are of sufficient length of contiguous nucleotides to specifically hybridize to SEQ ID NO:11 and SEQ ID NO:9. To acquire a diagnostic amplicon in which SEQ ID NO:2, or SEQ ID NO:4, or SEQ ID NO:6, or SEQ ID NO:8 is found, one would design a forward primer molecule based upon the inserted T-DNA (SEQ ID NO:9; from positions 863 through 14,180 of SEQ ID NO:10) and a reverse primer molecule based upon the 3' flanking corn genomic DNA (SEQ ID NO:12; from positions 14,181 through 15,216 of SEQ ID NO:10) in which the primer molecules are of sufficient length of contiguous nucleotides to specifically hybridize to SEQ ID NO:9 and SEQ ID NO:12.

For practical purposes, one should design primers which produce amplicons of a limited size range, preferably between 200 to 1000 bases. Smaller sized amplicons in general are more reliably produced in PCR reactions, allow for shorter cycle times, and can be easily separated and visualized on agarose gels or adapted for use in endpoint TAQMAN®-like assays. In addition, amplicons produced using said primer pairs can be cloned into vectors, propagated, isolated and sequenced, or can be sequenced directly with methods well established in the art. Any primer pair derived from the combinations of SEQ ID NO:11 and SEQ ID NO:9 or SEQ ID NO:12 and SEQ ID NO:9 that are useful in a DNA amplification method to produce an amplicon

diagnostic for event MON 95379 or progeny thereof is an aspect of the present invention. Any single isolated DNA polynucleotide primer molecule comprising at least eleven (11) contiguous nucleotides of SEQ ID NO:11, SEQ ID NO:9 or SEQ ID NO:12 or their complements that is useful in a DNA amplification method to produce an amplicon diagnostic for event MON 95379 or progeny thereof is an aspect of the present invention.

An example of the amplification conditions for this analysis is illustrated in Tables 28 and 29. Any modification of these methods or the use of DNA primers homologous or complementary to SEQ ID NO:11 or SEQ ID NO:12, or DNA sequences of the genetic elements contained in the transgene insert (SEQ ID NO:9) of event MON 95379, that produce an amplicon diagnostic for event MON 95379 is within the art. A diagnostic amplicon comprises a DNA molecule homologous or complementary to at least one transgene/genomic junction DNA or a substantial portion thereof.

An analysis for an event MON 95379 containing plant tissue sample should include a positive tissue control from a plant that contains event MON 95379, a negative control from a corn plant that does not contain event MON 95379 (e.g., LH244), and a negative control that contains no corn genomic DNA. A primer pair will amplify an endogenous corn DNA molecule and will serve as an internal control for the DNA amplification conditions. Additional primer sequences can be selected from SEQ ID NO: 11, SEQ ID NO: 12, or SEQ ID NO: 9 by those skilled in the art of DNA amplification methods. Conditions selected for the production of an amplicon by the methods shown in Table 28 and Table 29 may differ, but result in an amplicon diagnostic for event MON 95379 DNA. The use of DNA primer sequences within or with modifications to the methods of Table 28 and Table 29 are within the scope of the invention. An amplicon produced by at least one DNA primer sequence derived from SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:9 that is diagnostic for event MON 95379 is an aspect of the invention.

DNA detection kits that contain at least one DNA primer of sufficient length of contiguous nucleotides derived from SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:9 that, when used in a DNA amplification method, produces a diagnostic amplicon for event MON 95379 or its progeny is an aspect of the invention. A corn plant or seed, wherein its genome will produce an amplicon diagnostic for event MON 95379, when tested in a DNA amplification method is an aspect of the invention. The assay for the event MON 95379 amplicon can be performed by using an Applied Biosystems GeneAmp™ PCR System 9700, Stratagene Robocycler®, Eppendorf® Mastercycler® Gradient thermocycler or any other amplification system that can be used to produce an amplicon diagnostic of event MON 95379 as shown in Table 29.

All publications and published patent documents cited in this specification, and which are material to the invention, are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 28

<210> SEQ ID NO 1  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 50 nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette.

<400> SEQUENCE: 1

aagcaaatgt gtggtagtgg tccaattttt tttcaattca aaaatgtaga 50

<210> SEQ ID NO 2  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 50 nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA.

<400> SEQUENCE: 2

gctattatgg gtattatggg taggcacatg ggaatatagt gggaggcaga 50

<210> SEQ ID NO 3  
 <211> LENGTH: 100  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 100 nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette.

<400> SEQUENCE: 3

cgtacgttcg aggtacctgt cctgcaagca aatgtgtggt agtgggtccaa ttttttttca 60

attcaaaaat gtagatgtcc gcagcgttat tataaaatga 100

<210> SEQ ID NO 4  
 <211> LENGTH: 100  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 100 nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA.

<400> SEQUENCE: 4

cgccaagaag aacgattggc aaacagctat tatgggtatt atgggtaggc acatgggaat 60

atagtgggag gcagaggcgt gctgtgcct gctcttaatt 100

<210> SEQ ID NO 5  
 <211> LENGTH: 200  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 200 nucleotide sequence representing the 5' junction region of corn genomic DNA and the integrated transgenic expression cassette.

<400> SEQUENCE: 5

tcgctageta tagcttggtg gtatgcacca acaacgacct agggagctag cgtacgttcg 60

aggtacctgt cctgcaagca aatgtgtggt agtgggtccaa ttttttttca attcaaaaat 120

-continued

---

```

gtagatgtcc gcagcgttat tataaaatga aagtacattt tgataaaacg acaaattacg   180
atccgctcgta tttataggcg                                     200

```

```

<210> SEQ ID NO 6
<211> LENGTH: 200
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 200 nucleotide sequence representing the 3'
      junction region of the integrated transgenic expression cassette
      and the corn genomic DNA.

```

```

<400> SEQUENCE: 6
aggcgcgggt accgctcggtc cgggcctagt aggccaaagca ggacgtggcg cgccaagaag   60
aacgattggc aaacagctat tatgggtatt atgggtaggc acatgggaat atagtgggag   120
gcagaggcgt gcctgtgctt gctcttaatt tgagctcctc ccctggcctt gatagggcat   180
gtgcatgctc acaataatca                                     200

```

```

<210> SEQ ID NO 7
<211> LENGTH: 1160
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 1160 nucleotide sequence representing the 5'
      junction region of corn genomic DNA and the integrated transgenic
      expression cassette.

```

```

<400> SEQUENCE: 7
cgcagcgcag gtcctgtggt cctgtttggt ttggettctg gtagcttctg gccacaaaa   60
gctgctgctg actgccaaac gctcagcttt tcagccagct tctataaaat ttgttggggg   120
gcaaaaacca tccaaaatca acataaacac ataatcagtt gtagctgtgt aatagtagga   180
attcgtcact ttctagatcc tgagccctat gaacaatttt atcttctctc acacgtaatc   240
gtaatgatat tcagattctc cccacagcca gattctctc acagccagat ttcagaaaa   300
gctggtcaga aaaaagctga accaaaacagg ccctaggaca cacataggaa tggatcaaaa   360
aacatagctc cgtaccgta cgtacgtgca gtcacagtc gctgctagct agctagctgc   420
gccggcggcg gcgtgtcgg tcttgactca acgacgtccg cggctggcgc cgtcggccag   480
cggcttgtg cgcgggtcc catccggcgg agtccggctg ccagtggccg gcggcggctt   540
cttcttctgc gtcgtctctc cctttgccat ccccgctgcc acctggacga gggttcttgg   600
tcccgtcgac gtcgacgaga gacgtggcgc cgcatttggc tgccgccgta tcgctcgtct   660
cgtcgagatt gaaatcgaga gggagcgcgc gggatggctg ccgcccgttg cgcgacatgg   720
tctcgcccgg ccggcgcctg atcgacttct ctccgccgc ctctcgctagc tatagcttgg   780
tagtagtcac caacaacgac ctaggagct agcgtacgtt cgaggtacct gtcctgcaag   840
caaatgtgtg gtagtggtcc aattttttt caattcaaaa atgtagatgt ccgcagcgtt   900
attataaaat gaaagtacat tttgataaaa cgacaaatta cgatccgtcg tatttatagg   960
cgaaagcaat aacaaaatta ttctaattcg gaaatcttta ttctgacgtg tctacattca  1020
cgtccaaatg ggggcttaga tgaaaaactt cagcatcgat gcggccacca cgagtcgaca  1080
taacttcgta tagcatacat tatacgaagt tatagcggcc gcaaatcaac ctcaactctat  1140
ttaaagtagg tggtaggatt                                     1160

```

-continued

---

<210> SEQ ID NO 8  
 <211> LENGTH: 1178  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 1178 nucleotide sequence representing the 3' junction region of the integrated transgenic expression cassette and the corn genomic DNA.

<400> SEQUENCE: 8

taagggtccc aagtaaggcc ggccaagtaa cggtcggaag taaggcgagg gtaccgtcgg	60
tccgggccta gtaggccaag caggacgtgg cgcgccaaga agaacgattg gcaaacagct	120
attatgggta ttatgggtag gcatatggga atatagtggg aggcagaggc gtgcctgtgc	180
ctgctcttaa tttgagctcc tcccctggcc ctgatagggc atgtgcatgc gcacaataat	240
cattggagct gcctgaatga tttgcatgcg ctctgtgtaa acgttcacgg tccatgccag	300
tccaagaccg tcggcaagag atgaggggcca tgcgcagtac gtacatttgt atacataac	360
atattttttt ggatgatgcg atgcgtgtgc atatctggct gcgcccctcc ctctgcccac	420
gtcgtcggcg tgtacaacga attttccgtg ggatgggctg aacttataat gggcctactg	480
ggcgggccgt atgaatagc acacaattca cctgctttt ctcaccagga atgcattttc	540
gcgggagcca tcgtcctaac ttggcagaca gacacacaaa cacaggacag gagttcgcca	600
accccagcat gagagaaacc tatccggaca gctcctccag cctatgggccc gttgggacct	660
cagcctcagc gccccggccc cgtggcgggg caaaacactc cacagtcacac acggctgctg	720
ctgctgcctt tgttccgtcg gacctggggg aactttatag ctatgatgtt cgccttccc	780
aaccaatgtg cgagcgtaa ggatgttaag taacactagt taggattatg tggccaata	840
cctactttgc taggaacttg tacaccctc ataatataga gaaaagaaag atgccagtgg	900
tccaactcta taccatgtgt ctctgtgtt tgttatgtcg tgcttatgtg aagggagacg	960
agtcttttac agcttctcac gcctaactcg ttgacgggag agaagagagc ggatctgggtg	1020
atccgtggta acgtagttct caacaaatga cacattgagc acaccgtgtg catctcctca	1080
gctaaacatt gagggggatt ttttttaag aaaggctgag attccttac gctcccttg	1140
gcagggctcg tccaacagct tcttgaaaaa ttctataa	1178

<210> SEQ ID NO 9  
 <211> LENGTH: 13318  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Sequence corresponding to the transgenic inserted T-DNA of corn event MON 95379.

<400> SEQUENCE: 9

tttttttca attcaaaaat gtatagtgcc gcagcgttat tataaaatga aagtacattt	60
tgataaaaacg acaaattacg atccgtcgta tttataggcg aaagcaataa acaaattatt	120
ctaattcggg aatctttatt tcgacgtgtc tacattcacg tccaaatggg ggcttagatg	180
agaaacttca cgatcgatgc ggccaccacg agtcgacata acttcgtata gcatacatta	240
tacgaagtta tagcggccgc aaatcaacct cactctattt aaatgagggtg gtaggatttg	300
ccgctgagga cctattaccc tgttatccct agtttaaaaca cctgcaggaa aagattaatt	360
aacatcgatc atcatgttca attgacagca aattaaacta gttatatata gggttcatga	420
gatgagagta gctaattatt aacacaagtg gaagaaactt ttatttgacg gtatacaaat	480

-continued

---

aacaagatac	agagatcagg	ctgatggatc	aatcatatca	tatcagtgat	agatcgatcg	540
ttcgacacat	atatatgtat	gacaaaacat	agtataagta	cgtgcgaaag	ggtttatata	600
tatatggata	ttgatgcattg	gtgtaattgt	atggtaatta	agggatcgga	ggatcgatta	660
atgggtccct	atcactcctc	catcagtagc	agctcgaccg	agtcaccgat	gaaagtgcct	720
tccgtctcgc	cgatctcgat	ccagactttg	tcggctcccg	ggaagtactc	cagctccttg	780
gtgacgtacc	cggcgggcag	cggagtgtag	tcaccgtagc	cacggttgga	ctcgcaaggg	840
ttgtccctcc	gtccgtcggg	gtacgccttc	tccctgtaag	cggacgcata	atccgcaggc	900
acgctgctgt	tggactcgta	cgcgccatca	tagcccgcat	tccgggaagt	gtaagtgcct	960
tcgtactcct	cttgctgggc	agtgtagtcg	ttgcaagtaa	cgggtgtgtt	cgggtacacc	1020
tccctcctca	cacagttcga	gaacttgagc	tcgtcgggtg	tggtctcgat	ctcgtggata	1080
gtcacgcagc	cttcaccgta	gccttccttg	tacgcagtca	cacggaggat	gtagccgcga	1140
ccagggcaga	cgcggactcc	ctggctcacc	tctgcctccc	actccggcac	gaccagcacg	1200
gaccgggtgt	tggtctgctc	ctcgacgtcc	acatggccct	tgacggtcca	gcaactcagc	1260
ccgttggtga	agtcaccggt	cttaataaca	ttcctggcat	cgtacagggt	gaatgctgtg	1320
aagatgcggc	cctccagttc	ctcgaagatc	gccgcgtcca	caccagggat	cactgacagc	1380
tcgggcaggt	acgectcagc	gatgctgtgg	accgccttgt	cagccgcgtg	gatcatggcg	1440
atggtggtgt	cagcctggag	gcggctcgta	tggtctgtga	cgaagagtgc	gtccactgac	1500
tccctggcct	ccttgtagac	gatggtggtc	tccactccca	gcttctcccg	cttgtctcgc	1560
catttctctc	eggcgcgctt	cacacgagcg	agggcctcgc	ccacgagagg	cttctctcca	1620
aggaactcca	ggttgcagag	gcgggcatgt	ccgtcctggg	tcttaatctt	gaagataacc	1680
cagacaccca	ggtcctcggt	gaggtcggtg	cagcccacgt	cgatgtccag	agagaagtgg	1740
tgcgagtggg	gagcgcactt	tccgattgga	gagggagcgc	tcaatggcca	aagtgagccc	1800
gtgcctggca	cgttcaactg	ctcgtgcttg	gcggtgtatc	ggatcaggta	gatttccagg	1860
tccctgactat	cctcgatgta	cccgcgaagc	tggtagcggg	tgtagcctt	caatttactc	1920
tcgtctatct	tctggtagag	atacgtcggg	tagcattcgt	cgaagtccc	gagcagcgtc	1980
acatagttct	ccttgaacac	atcgtcgcct	ccctggatgg	tgatgtctgt	actcccgcgc	2040
caaccacggg	cgagctggcg	ggtgatgcca	cggaaagtcc	ggctcctggg	caaatccgc	2100
tcgtcactca	gacgcttcgc	gtgtttgacc	ttctcggaca	gctccttctt	ctcgtcgagg	2160
cagaactcgt	cggacaggca	ctccacaagg	ttggacactt	ggtcaatgtg	atagtccgtg	2220
acgtcggctc	tgagcccaat	ctgggtgctg	ctggtgaaca	ggtcgttgac	agccttctga	2280
gcccgcctca	ggtcagactc	ggcctcgaac	gtcgcacccg	ccaggatgat	ctcaatcttg	2340
tcgatgtaca	ggtcgcagca	gctgatggag	ccagcaccga	acagcgggtg	ctcggagatc	2400
ccaatgatgt	cagggttcgc	tctgaatgag	aaagggttcg	agaaatcagt	gtaccggaag	2460
gttcggctag	tcaagttctc	gctatctccc	atcgtcttct	ggagaggcat	ggtgactgag	2520
acttggcctc	ccacgcctgt	ggacgcggct	cgggtcagga	caatgacct	cgcgtcgcgg	2580
cttgaagcgt	agcggaaact	cagacggtaa	cgtcgggtga	tcggcagagt	gatgttcaact	2640
tgcaacgaaa	cgaagtgcgc	gaaagtgttg	cgccgcagta	tgctctccacc	ggtgaagccc	2700
ggaccgggtg	tgacgctcgt	gccacccccc	acacggaagc	ccttcacgag	cgggatctgg	2760
ttaatggagt	cggaagagat	cgtgttggtc	cggtcggcgc	tgcggtgtgt	ccaggagtac	2820

-continued

---

accggcgcac	gcagcgtggt	gccagagatg	agccggatgt	tggacaggcg	gtggctgtag	2880
gactcgtagt	tcgggcttc	cgtggctcgc	ggcggcagct	cagtttact	gtcgaagagc	2940
tgggtgccga	cgcccggtga	cccgatggtg	tagagcaatg	accgcgcag	ggagttcaac	3000
gggttacgcc	agttgaaccg	ggcccacggg	acgccgttga	cgggcgttgt	cagaaggatg	3060
ttaatgcccg	cgaacgactc	agtcggtaa	acgtcgcggc	tcgtgaactg	tagcgttaact	3120
gggttgatgg	agggttggt	ggtgcctgg	gtactgggtc	tcaagctgcc	acggatcgtc	3180
ctactctcca	atcggtgccc	gaccagtag	ttcatgtact	gtgtggtgga	ccagcgtgag	3240
agcacggaga	agatcgtgag	ctgctcggga	aagtcgagca	ggtgaggagg	gcggatgacg	3300
gcggcctcga	ttgccgagaa	ggagggcgc	ttggtgtga	accaattcgt	ggatgcgaaa	3360
ccggagggtg	cgttcgtccg	gccgattggg	tctgtgtaga	tctcccgcgt	gagttgcgcg	3420
ctagtgttca	ttgggtacac	ccgtgtgctg	taagacggga	atagcgcac	caggcttagg	3480
acacccaaag	tcaagtgcgc	gcggaactgg	ttgtaacgga	gccatgactc	agcgttggtg	3540
ccgcgcaggt	tgttgagccc	tgtgtgtac	catcgagcgc	agtagtcaact	gtactcacgg	3600
gtcttctcca	cttggcgtc	gtagtaacgc	tggatttctt	gagatgtgag	tccgaactcg	3660
gagccgaaca	ggctggcgtc	ccgcagcagg	agcagatgta	agttggcggc	ctgggcgtac	3720
accataagga	gtggaacttc	ctggttctg	atggcgaaca	gcggcatagc	gttcaggaaag	3780
tcgagttcga	gagcagtgta	ctgggtgtag	agtaccgagc	gtgttcgcgc	gtcatccctg	3840
ttctccaacc	agtctccag	tgactgctgg	tatgccgaa	aggagtttcc	caggccctgt	3900
agcctggcca	gagcagtatc	gcggtgttc	tccgtgactt	gctgccggat	caactgctcc	3960
acatgctcca	ggaagatctc	ccacgggtca	cgaccgcgag	gccacagctc	gcccacaagg	4020
aacgagtaga	aggacgcgat	ctgaccggcg	aatgggacac	cgaggacgcc	gaggatacgg	4080
cccgcgatgt	tgatgccggg	ctgaaccgtg	ctggccgaca	cgaatgggtc	gatgtgttg	4140
ccctcggcta	tgcagaggga	gtcctcgatg	cgcgcgtcgg	tggacagggt	catctgcgcg	4200
gagtggtttg	aaacagcagg	gatcgagagg	gcgttgatga	tctcgttctc	gttcttcggg	4260
ttgctcgtca	tggtggtggc	cgccaagta	acggtccgac	ctgcagaagt	atcaccaaac	4320
aacaggtgga	gcatcggaca	aaagaacag	taccaagcaa	ataaatagcg	tatgaaggca	4380
gggctaaaaa	atccacatat	agctgctgca	tatgccatca	tccaagtata	tcaagatcaa	4440
aataattata	aaacatactt	gtttattata	atagataggt	actcaagggt	agagcatatg	4500
aatagatgcc	gcatatgcc	tcatgtatat	gcatcagtaa	aaccacatc	aacatgtata	4560
cctatcctag	atcaatattt	ccatccatca	tcttaaactc	gtaactatga	agatgtatgg	4620
cacacacata	aagatacaaa	attaataaat	ccaccaggta	gtttgaaaca	gtattctact	4680
ccgatctaga	acgaccgccc	aaccagacca	catcatcaca	accaagcgaa	aaaaagcatc	4740
tctgtatatg	catcagtaaa	accgcgatca	acatgtatac	ctatcctaga	tcgatatttc	4800
catccatcat	cttaaactcg	taactatgaa	gatgtaggca	catacataca	gatcctttaa	4860
taaatccacc	aggtagtttg	aaacagtatt	ctactccgat	ctagaacgac	cctttatcct	4920
ttaataaatc	caccaggtag	tttgaaacag	tattctactc	cgatctagaa	cgaccgcca	4980
accagacat	atcatcacia	ccaagcgaga	aaaaaagcat	ctctgtatat	gcatcagtaa	5040
aaccgcgcatc	aacatgtata	cctatcctag	atcgatattt	ccatccatca	tcttaaactc	5100
gtaactatga	agatgtatgg	cacacacata	cagatacaaa	attaataaat	ccaccaggta	5160
gcttgaaaca	gtatcctact	ccgatctaga	acgaccgccc	aaccagacca	catcatcaca	5220

-continued

---

aacatgaaca tgaacatggt tgctctaaca caaacatgaa cagaagtaga actaccgggc	5280
cctaaccatg gaccggatcg ccgatctaga gaaggtagag agaggggggg ggggaggatg	5340
agcggcgtac cttgaagcgg aggtgcccac ggctggattt gggggagatc tggttgcgtg	5400
tgtgtgcgct ccgaacgaac acgaggttgg gaaagagggg tgtggagggg gtgtctattt	5460
attacggcgg gcgaggaagg gaaagcgaag gagcgggtgg aaaggaatcc cccgtagctg	5520
ccggtgcccg gagaggagga agaggccgcc tgccgtgccg cctcacgtct gccgctccgc	5580
cacgcaattt ctggatgccg acagcgggac aagtccaacg gtggagcggg actctcgaga	5640
ggggtccaga ggcagctaca gagatgccgt gccgtctgct tcgcttgccc cgacgcgacg	5700
ctgctggttc gctggttggt gtcctttaga ctctcgacg gcgctgaaca gcctgtcatt	5760
atctactcga aacaagaaaa atgtttgctt agttttttta tttcttaaag ggtatttgtt	5820
ttattttag tcaatttatt ttatttcatt ttatatctaa attattaat aaaaaactaa	5880
aatagagttt tagttttaa aaatttagag actaaaaaga ataaatgga tgtactaaaa	5940
attagtctat agaaaccatc aaccctaaat cctaaatgga tgtactaata aaatggatga	6000
agtattatat aggtcaagct atttgcaaaa aaaacagaga gatcacatgc aactaaaaa	6060
gataaaactg tagattgtca aaatattcaa ttatccttta gaccatgtct agcagtttat	6120
ttatatgac ctctaaaaa ctaataattt tttagtatta tagactatat tttttaga	6180
gtgaagtta aatatatgta tagagataga taaactacac ttcaataag tgtgacaaaa	6240
aaatatgtgg taatttttta tacttttagac atgcaatgct ctttatctct agagaggggc	6300
acgacgcatg caagcttgtt aacgcggccg ctaattctca gtccaaagcc tcaacaaggt	6360
cagggtacag agtctccaaa ccattagcca aaagctacag gagatcaatg aagaatcttc	6420
aatcaagta aactactggt ccagcacatg catcatggtc agtaagtctc agaaaaagc	6480
atccaccgaa gacttaaaagt tagtgggcat ctttgaaagt aatcttgtca acatcgagca	6540
gctggcttgt ggggaccaga caaaaaagga atggtgcaga attgttaggc gcacctacca	6600
aaagcatctt tgcctttatt gcaagataa agcagattcc tctagtacaa gtggggaaca	6660
aaataacgtg gaaaagagct gtccctgacag cccactcact aatgcgtatg acgaacgag	6720
tgacgaccac aaaagaatta gcttgagctc aggatttagc agcattccag attgggttca	6780
atcaacaagg tacgagccat atcactttat tcaaattggt atcgccaaaa ccaagaagga	6840
actcccaccc tcaaaagttt gtaaggaagg tcgaggccgc tgtactgtca tattgtcgtg	6900
gtttttcaat tgctgtacct gatgcaaacg taatgggttt actaatcttg caccgcccgg	6960
cttcaaaatg aagagtgcta atttggtcca cgtcaccatc accggttcga actgtctaga	7020
atggcaggca aagatgattg gacaggcatg cagggaaaaa gagcaccggt gacgatgat	7080
gcgagttccc accattgcca gcaatgatta tcagccacac gacttactct tcagagctaa	7140
ccactgccat gcagagaaaa agtgaatcat attgtcatga tctacaacga agtgaacaaa	7200
tcaggcatgc taaagtgtg aaactttact gatctctcat gttggacaac aaagaatacg	7260
ggaatacatc agcaacgcaa ctcttgagct ttgcttgccc aatgaccagc tagaatttcc	7320
aagcatttac agaaacatga ctttaagttt cagaaaaaca aatacaaggc cactaaataa	7380
gcgtggggat aacatatcct ccagatgaca ggcaatctgc aactgcagc cattcaaatg	7440
tacgattaac aaaatattta agcgcacat gagataatat atcctccaat tagggccttt	7500
agtattgtca ttagctcata accatgggtc atcctcacat ggacgctgca taagaagttc	7560

-continued

---

ataatagcaa cagacatatg aacaaagcat ggtgcgcctg cccggccgga ctgctagta	7620
ctaccaatca tggaaatagc tagtaccta aatgaaatta aaatggtttt tagcgattat	7680
ccacgccgtc cagaatactc taatecaciaa gttgaggccg cccatgaagc cgcgagagg	7740
cgacgccatg tgtataaaag gggcctaagc tgagtggact tgctgcatca gattagtaag	7800
caatctcaag cgcagagagc caaagctttc ggtgtagctc gaagagcaaa gcgaaggcaa	7860
ggcgcgcctg taacctactc accaacctcg ccattcctct ccaaactggt gtgctgctgt	7920
ctagatctcc cacactaac tagttactcc tcgtagatct cggctacctg gctcaagatc	7980
cggggtcaga tccgggtccg gggattttct ttgtgcccta tggctgtatt ttggcgtctg	8040
tggctgatga cagcgtgtgt tctcgagtgc ggatgcaatc tgagttatat aggcaaatgg	8100
ccttgtcaac tcgggcagcg gcattgcttt gctcagtggtg tttgaatgtg ctgaaattca	8160
tgtagtaggc tgtaggctgt gcatttcttg atttgcgtct tgcataatc actgggggat	8220
tttctaaacc taacaagttt aaaattagac cattcaacca aagacaggag gaataagtga	8280
agctgttgta gtcacagctt atggccgatc caaaattgtg taggaatgtg aatatgtgat	8340
gctacaaaaca tacccttgta agctaccatg ctatttatca tgttocatca tggtgattgg	8400
tgagcactca tgaaaatttc agatccaaac ctagtgttac atgtggattt gtgctctgca	8460
atctatcgcc agtaataaaa tggttgagtg atccagctac taaaaatca cattgcatac	8520
tttttttttt tgtagattat gcactcctgt tttgggtggg gggttcctga tgcaggaat	8580
ataaatttag cctgctgatt taggtagcac tgccgggtgca cactttgggt tttgaatact	8640
tgtagtcttc cagcttcttg tagaactggg acaatgtggg ccatatataa gaagggtgt	8700
caactagcac atgctcacta attagtctaa acatttatgt ttttattcat tcaggtcagg	8760
tgcaatcata gaagtagtta atgacaatac tttagtgtgt ctaaatattat ttatgtatgg	8820
actcaaatta acatgcaaaa catatgagat tagtggcatg cattcttttt cttaaatgtg	8880
gaaaatacga gataatgata actgtgaagc tctgttagta ctcttcatta ctctatttga	8940
gtggcagcat atctcatgct agccataaag caagttctag acgtattctg ttgttaatta	9000
cttgtagcta tataaccocaa cctagtcatt ccagcttatg tctcttagag atcatgttta	9060
ttagcacctc aagatttctc ctgcacagta tagtaactat cgaaaaagat attatttctt	9120
tgtttttaat tgacaacctt cacgtgctac ttatttttgc agcttggccg gccaccaacc	9180
atggctgaga tcaacaacca gaaccagtgc gtcccgtaca actgctgag caaccctaag	9240
gagatcatcc tgggtgagga acgcctggag accggcaaca ccgtagccga cattagcctg	9300
ggcctcatca acttctctc cagcaacttc gtgcccggcg gtggcttcat cgtgggcctc	9360
ctggagctta tctggggctt catcgcccg tcccagtggg acatcttctc cgcccagatc	9420
gagcaactga tcagccagcg gatcgaggag ttcgctagga accaggccat ctcccgcctg	9480
gagggactct ccaacctcta caaggtgtac gtgcgcgcgt tcagcgactg ggagaaggac	9540
ccgaccaacc cggccctccg cgaggaaatg cgtatccagt tcaacgatat gaactcggcc	9600
ctcatcaccg ccattcccctc ctcccgctg cagaactacg aggtggccct cctgtccgtg	9660
tacgttcaag ccgccaacct ccacctctcc atcctccgcg acgtgagcgt gttcggcgag	9720
cgctggggct acgacaccgc caccatcaac aaccgctact ccgacctcac ctccctcatc	9780
cacgtttaca ccaacctctg cgtggacacg tacaaccagg gcctccgcg cctggagggc	9840
cgcttctctc ccgactggat cgtgtacaac cgcttccgcc gccagctcac catctccgtc	9900
ctggacatcg tcgcttctt tcccaactac gacatccgca cctaccctat ccagaccgcc	9960

-continued

---

```

accagctca cccgcgaggt ctacctogac ctcccgttca tcaacgagaa cctcagcccc 10020
gccgcgctct acccgacctt ctccgcoget gagtccgcca tcattcgag cccgcacctc 10080
gtggacttcc tcaactcctt caccatctac accgactccc tcgccccgag cgcctactgg 10140
ggcggtcacc tcgtgaactc ctccgcacc gccaccacta ccaacctcat cgcagccccg 10200
ctctacggcc gcgagggcaa caccgagcgc ccggtgacca tcaccgccag cccgagcgtg 10260
cccatcttcc gcaccctcag ctacccccacc gccctggaca acagcaaccc tgtggcgggc 10320
atcgaggcgg tggagttoca gaacaccatc tccaggagca tctaccgcaa gagcggccct 10380
atcgacagct tcagcgagct gcctctcag gacgccagcg tgagccctgc catcggtac 10440
agccacaggc tgtgccacgc caccttctctg gagcgcatca gcggccctcg catcgcgggc 10500
accgtgttct cgtggaccca ccgagcgcct tctctacga acgaggtgtc tctagtcgc 10560
atcaccaga tcccttgggt caaggccac accctggcta gtggcgtag tgtcatcaag 10620
ggccctggct tcaccggtgg tgacatcctg accaggaact ctatgggcga gctgggcact 10680
ctgagggtea ctttcaactg ccgctgcct cagtcttact acatccgctt ccgctacgct 10740
agtgtcgcta accgctctgg tactttccgc tactctcagc ctccgtctta cggtatctct 10800
ttccctaaga ctatggacgc tggtgagcct ctgaccagta ggagcttcgc tcacactact 10860
ctgttcaactc ctatcacttt ctctagggct caggaggagt tcgacctga catccagtct 10920
ggtgtgtaca tcgacaggat cgagttcact cccgtgaccg ccacgcttca gcccgagtac 10980
gaccttgagc gcgcccagaa ggtggtgaac gccctcttca ctagcaacta ccagctaggc 11040
ctgaagactg acgtgaccga ctaccacatc gaccaagtga gcaacctagt ggctgctc 11100
tccgacgagt tctgcctcga cgagaagcgc gagctgtccg agaaggtgaa gcacgccaag 11160
cgcctctccg acgagcga caactgctcag gacccccact tcaggggcat caacagggcag 11220
cccgaccgag gctggcggcg ctcccaccgac atcaccatcc agggcgggtga cgacgtattc 11280
aaggagaact acgttacctt ccccgccacc ttcgacgagt gttacccac ctacctctac 11340
cagaagatcg acgagtcaca gctgaaggcc tacaccgct accagctccg cggctacatc 11400
gaggactccc aggacctgga aatctaactc atccgctaca acgccaagca cgagatcgtg 11460
aacgtgcctg gcaccggcag cctctggcct ctccagcgtg agaaccagat cggcccttgc 11520
ggcgagccta accgctgcgc cctcaccctc gagtggaaacc ctgacctcca ctgctcgtgc 11580
agggacggcg agaagtgcgc ccaccatagc caccacttct ctctggacat cgacgtgggc 11640
tgcaccgacc tgaacgagga cctggcgctg tgggttatct tcaagatcaa gaccaggac 11700
ggtcacgcca ggctgggtaa cctggagttc cttgaggaaa agcctctgct ggggtgaggc 11760
ctggccaggg tcaagagggc tgagaagaaa tggagggata agagggagac cctgcagctg 11820
gagaccacta tcgtctacaa ggaggctaag gactctgtcg atgctctgtt cgtcaactct 11880
cagtacgata gactgcaagc tgataccaac atcgctatga tccacgctgc ggataagcgg 11940
gtccaccgga tccgggaggc ttaccttccg gagctttctg tcatcccggg tgtcaacgct 12000
gcgatcttcc aggaacttga ggaacggatc ttcactgcgt ttagtcttta cgatgcgagg 12060
aacatcatca agaacgggga cttcaacaat ggtctgctgt gctggaacgt caagggtcat 12120
gtcgaggtcg aggaacaaaa caatcatcgt agtgtccttg tcattcctga gtgggaggcg 12180
gaggtctctc aagaggtccg tgtttgcccg gggcgtgggt acattctctg tgttactgcg 12240
tacaaggagg ggtacgggga ggggtgcggt actattcatg agattgagaa caatactgat 12300

```

-continued

---

```

gagcttaagt tcaacaattg tgttgaggag gaggtttacc cgaacaatac tgttacgtgc 12360
atcaactaca cggcaacgca agaggaatac gaggggacgt acacctcgcg taatagaggg 12420
tatgatgagg cgtacggaaa caaccctcg gttccagcag attatgcctc ggtttatgag 12480
gagaagtcgt acacggatag acgacgcgag aatccatgtg agtcaaatcg aggatacggg 12540
gattacacac cattaccagc aggatacgtt acaaaggagt tggaatactt cccggaaca 12600
gataaagttt ggattgaaat cggagaaaca gaaggaacat tcatcgtcga ctcagtagaa 12660
ttgttgttga tggaagaatg atagggacc cggacccgcc aaaaccattg caaagactat 12720
agtttggggg ggagtatact tggttgtgta catgcctcgcg tgttccattg tacacacaaa 12780
acctagccac ctcttgactc ttgagtgtat gcttgttacc cgtgtgttga agtttgaag 12840
aggcaccatc actatagatg atggcttgtg tccctctttc atcaagattg aataatata 12900
gctactttga gagecgtatc ctgcttgcct gattgtgtta atacttacc cgtcccaca 12960
ctcccacaat ataaggaat aaggatattt ggcagtttag agcaaaatc ccttatattt 13020
ttgggacgga tgcctcttt tctgcatttt tttatgttca tatgttcctg aagagtaagg 13080
tggatcctga tcaacctgct ggtttatggt gattgatttg agtggaaatg aatgggcca 13140
cgtccggcat acagttatgc ttcagttaat taaaagtaag ggtcccaagt aaggccggc 13200
aagtaacggt ccgaagtaag gcgcggttac cgtcggccg gccctagtag gccaaagcag 13260
acgtggcgcg ccaagaagaa cgattggcaa acagctatta tgggtattat gggtaggc 13318

```

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 15216

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial

&lt;220&gt; FEATURE:

```

<223> OTHER INFORMATION: Sequence corresponding to the contig nucleotide
sequence of the 5' genomic flanking DNA nucleotide sequence, the
inserted T-DNA sequence in event MON 95379, and the 3' genomic
flanking DNA nucleotide sequence.

```

&lt;400&gt; SEQUENCE: 10

```

cgcagcgcag gtcctgtggt cctgtttgtt ttggttctg gtagcttctg gccacaaaa 60
gctgctgcgg actgccaac gctcagcttt tcagccagct tctataaaat ttgttggggg 120
gcaaaaacca tccaaatca acataaacac ataatcagtt gagtcgttgt aatagtagga 180
atcgtcact ttctagatcc tgagccctat gaacaatttt atcttcctcc acacgtaatc 240
gtaatgatat tcagattctc cccacagcca gattctctc acagccagat ttcagaaaa 300
gctggtcaga aaaaagctga accaaacagg ccctaggaca cacataggaa tggatcaaaa 360
aacatagcgc cgtaccgta cgtacgtgca gtcacagtca gctgctagct agctagctgc 420
gccggcggcg gcgttgcgg tcttgactca acgacgtccg cggtggcgc cgtcggccag 480
cggcttgttg ccgccggtcc catccggcgg agtccggctg ccagtggccg gcggcggctt 540
cttcttcgtc gtcgtctctc cctttgccat ccccgctgcc acctggacga gggttctggc 600
tcccgtcgac gtcgacgaga gacgtggcgc cgcatttggc tgccgcccga tcgtcgtcgt 660
cgtcgagatt gaaatcgaga gggagcgcgc gggatggctg ccgcccgttg cgcgacatgg 720
tctcgcggcg ccggcgcctg atcgacttct ctccgccgc ctccgctagc tatagcttgg 780
tagtagtcac caacaacgac ctaggagct agcgtacgtt cgaggtacct gtcctgcaag 840
caaatgtgtg gtagtggtcc aatttttttt caattcaaaa atgtagatgt ccgcagcgtt 900
attataaaat gaaagtacat tttgataaaa cgacaaatta cgatccgtcg tatttatagg 960

```

-continued

---

cgaagcaat aaacaaatta ttctaattcg gaaatcttta tttcgacgtg tctacattca	1020
cgtccaaatg ggggcttaga tgagaaactt cacgatcgat gcggccacca cgagtcgaca	1080
taacttcgta tagcatacat tatacgaagt tatagcggcc gcaaatcaac ctcactctat	1140
ttaaagagg tggtaggatt tgccgcgtga gacctattac cctgttatcc ctagttaaa	1200
cacctgcagg aaaagattaa ttaacatcga tcatcatggt caattgacag caaattaaac	1260
tagttatata tagggttcat gagatgagag tagctaatta ttaacacaag tggaagaaac	1320
ttttatttga cggatatacaa ataacaagat acagagatca ggctgatgga tcaatcata	1380
catatcagtg atagatcgat cgttcgacac atatatatgt atgacaaaac atagtataag	1440
tacgtgcgaa agggtttata tatatatgga tattgatgca tgggtgtaat gtatggtaat	1500
taagggatcg gaggatcgat taatgggtcc ctatcactcc tccatcagta gcagctcgac	1560
cgagtccacg atgaaagtgc cctccgtctc gccgatctcg atccagactt tgtcggctctc	1620
cgggaagtac tccagctcct tggtagcgtc cccggcgggc agcggagtgt agtcaccgta	1680
gccacggttg gactcgcaag ggttgtcctt ccgtccgtcg gtgtacgctt tctcctcgta	1740
agcggacgca taatccgacg gcacgctgct gttggactcg tacgcgccat catagccgcy	1800
attccgggaa gtgtaagtgc cctcgtactc ctcttgctg gcagtgtagt cgttgcaagt	1860
aacggtgttg ttcgggtaca cctcctcctc cacacagttc gagaacttga gctcgtcgg	1920
gttgttctcg atctcgtgga tagtcacgca gccctcaccg tagccttctt tgtacgcagt	1980
cacacggagg atgtagccgc gaccagggca gacgcggacc tcttggtca cctctgctc	2040
ccactccgcy acgaccagca cggaccggtg gttgttctgc tctcgcagct ccaatggcc	2100
cttgacgttc cagcaactca gcccggtgtt gaagtcaccg ttcttaataa cattcctggc	2160
atcgtacagg ctgaatgctg tgaagatgcy gccctccagt tctcgaaga tcgccgcyt	2220
cacaccaggy atcactgaca gctcggcag gtacgcctca cggatgctgt ggaccgctt	2280
gtcagccgcy tggatcatgg cgatgttgg ttcagcctgg aggcggctgt actggtggt	2340
gacgaagagt gcgtccactg actccttggc ctctctgtac acgatgttg tctccactc	2400
cagcttctcc cgttctctc gccatttctt ctccgcygc ttcacacgag cgaggcctc	2460
gcccacgaga ggcttctcct caaggaactc caggttgccg aggcggccat gtcgctcctg	2520
ggtcttaatc ttgaagataa ccagacacc caggtcctcg ttgaggtcgg tgcagccac	2580
gtcagtgctc agagagaagt ggtgcyagt gtgagcgcac tttccgattg gagagggagc	2640
gctcaatggc caaagtgagc ccgtgcctgg cacgttctc gtctcgtgct tggcgttgta	2700
tccgatcagg tagatttoca ggtcctgact atcctcgatg taccgcgcaa gctggtagcy	2760
ggtgtacgcc ttcaatttac tctcgtctat ctcttggtag agatacgtcg ggtagcattc	2820
gtcgaagtc ccgagcagcy tccatagtt ctcttgaac acatcgtcgc ctccctggat	2880
ggtgatgtct gtactccgc gccaccacg gtcgagctgy cggttgatgc cacggaagtt	2940
cgggtcctgg agcaattcc gctcgtcact cagacgcttc gcgtggttga ccttctcgga	3000
cagctccttc ttctcgtcga ggcagaactc gtcggacagc cactcccaa ggttgacac	3060
ttggtcaatg tgatagtcg tgacgtcgg cttagacca atctggttgc tgcgttgtaa	3120
cagttcgttg acagccttct gagcccgctc caggtcagac tcggcctcga acgtcgcac	3180
cgccaggatg atctcaatct tgtcagtgta cagttcgcga gagctgatgg agccagcacc	3240
gaacagcggg tgcctcgaga tcccaatgat gtcagggctc gctctgaatg agaaaggtt	3300
cgagaaatca gtgtaccgga aggttcggct agtcaagttc tcgcctatct ccatcgtctt	3360

-continued

---

ctggagagggc atgttgactg agacttggcc tcccacgcct gtggacgagg ctcgggtcag	3420
gacaatgacc ctcgcgtcgc ggcttgaagc gtagcggaac ctcagacggg aacgctgggt	3480
gatcggcgag ttgatgttca cttgcaacga aacgaagtcg ccgaaagtgt tgcgcccag	3540
tatgtctcca cgggtgaagc cgggaccggg gatgacgctc gtgccacccc acacacggaa	3600
gcccttcaag agcgggatct ggttaatgga gtcggaagag atcgtgttgg tccggtcggc	3660
gctgcgggtg gtccaggagt acaccggcgc acgcagcgtg ttgccagaga tgagccggat	3720
gttgacaggg cgggtggtgt aggactcgtg gttcggggcgt tccgtggtct cggggcgag	3780
ctcagtttca ctgtcgaaga gctgggtgcc gacgcccgtg taccgatgg tgtagagcaa	3840
tgaccgcgc agggagtcca acgggttacg ccagttgaac cgggcccacg ggacccggt	3900
gacggcggtt gtcagaagga tgttaatgcc cgcgaacgac tcagtccggg aaacgtcgcg	3960
gctcgtgaac tgtagcgtaa ctgggtgat ggaggtgttg gtgtgccgt gggactggt	4020
gctcaagctg ccacggatcg tctactctc caatcgggac ccgaccaggt agttcatgta	4080
ctgtgtgttg gaccagcgtg agagcacgga gaagatcgtg agctgctcgg gaaagtcgag	4140
caggtgagga gggcggatga cggcggcctc gattgccgag aaggaggggt cgttgttgt	4200
gaaccaatc gtggatgcca aaccggaggg tgcgttcgtc cggccgattg ggtctgtgta	4260
gatctcccgc gtgagttgcg cgtagtggtt cattgggtac acccgtgtgt cgtaaagcgg	4320
gaatagcgc accaggtcta ggacacccaa agtcaagtcg cggcggaaact ggttgaacg	4380
gagccatgac tcagcgttgg tgccgcgag gttgttgagc cctgtgtgt accatcgagc	4440
gcagtagtca ctgtactcac gggctctctc cacttggcgc tcgtagtaac gctggatttc	4500
ttgagatgtg agtccgaact cggagccgaa caggctggcg tcccgcagca ggagcagatg	4560
taagttggcg gcctgggctg acaccataag gagggaaact tccctggtcc tgatggcgaa	4620
cagcggcata gcgttcagga agtcgagttc gagagcagat tactgggtgt agagtaccga	4680
gcgtgttcgc gcgtcatccc tgttctccaa ccagtcctcc agtgactgct ggtatgcccg	4740
aaaggagttt cccaggccct gtgacctggc cagagcagta tcgagggtgt tctccgtgac	4800
ttgctgcccg atcaactgct ccacatgctc caggaagatc tcccacgggt cacgaccgcg	4860
aggccacagc tcgcccacaa ggaacgagta gaaggacgcg atctgaccgg cgaatgggac	4920
accgaggacg ccgaggatac ggcgcgcgat gttgatgccg gtctgaaccg tgctggccga	4980
cacgaatggg tcgatgttgt tgccctcggc tatgcagagg gagtctcga tgcgcgctc	5040
ggtggacagg ttcactctgc cggagtggtt tgaaacagca gggatcgaga gggcgttgat	5100
gatctcgttc tcgttcttcc ggttgctcgt catggtggtg gccggccaag taacggtccg	5160
acctgcagaa gtatcaccaa acaacagggt gagcatcgga caaaagaaac agtaccaagc	5220
aaataaatag cgtatgaagg cagggctaaa aaatccacat atagctgctg catatgccat	5280
catccaagta tatcaagatc aaaataatta taaaacatac ttgtttatta taatagatag	5340
gtactcaagg ttagagcata tgaatagatg ccgcatatgc catcatgtat atgcatcagt	5400
aaaaccacac tcaacatgta tacctatcct agatcaatat ttccatccat catcttaaac	5460
tcgtaactat gaagatgtat ggcacacaca taaagataca aaattaataa atccaccag	5520
tagtttgaag cagtattcta ctccgatcta gaacgaccgc ccaaccagac cacatcatca	5580
caaccaagcg aaaaaagca tctctgtata tgcacagta aaaccgcat caacatgtat	5640
acctatccta gatcgatatt tccatccatc atcttaaacg cgtaactatg aagatgtagg	5700

-continued

---

cacatacata	cagatccttt	aataaatcca	ccaggtagtt	tgaaacagta	ttctactccg	5760
atctagaacg	accctttatc	ctttaataaa	tccaccaggt	agtttgaac	agtattctac	5820
tccgatctag	aacgaccgcc	caaccagacc	atatcatcac	aaccaagcga	gaaaaaaagc	5880
atctctgtat	atgcatcagt	aaaaccogca	tcaacatgta	tacctatcct	agatcgatat	5940
ttccatccat	catcttaaac	tcgtaactat	gaagatgtat	ggcacacaca	tacagataca	6000
aaattaataa	atccaccagg	tagcttgaaa	cagtatccta	ctccgatcta	gaacgaccgc	6060
ccaaccagac	cacatcatca	caaacatgaa	catgaacatg	tttgccttaa	cacaaacatg	6120
aacagaagta	gaactaccgg	gccctaacca	tggaccggat	cgccgatcta	gagaaggtag	6180
agagaggggg	ggggggagga	tgagcggcgt	accttgaagc	ggaggtgccg	acggctggat	6240
ttgggggaga	tctggttgcg	tgtgtgtgcg	ctccgaacga	acacgaggtt	ggggaaagag	6300
ggtgtggagg	gggtgtctat	ttattacggc	gggcgaggaa	gggaaagcga	aggagcggtg	6360
ggaaaggaat	ccccctagc	tgccggtgcc	gtgagaggag	gaagaggccg	cctgccgtgc	6420
cgctcactgc	ctgccgctcc	gccacgcaat	ttctggatgc	cgacagcggg	gcaagtccaa	6480
cgggtggagc	gaactctcga	gaggggtcca	gaggcagcta	cagagatgcc	gtgccgtctg	6540
cttcgcttgg	cccgcgcga	cgetgctggt	tcgctggttg	gtgtccgtta	gactcgtcga	6600
cggcgttgaa	cagcctgtca	ttatctactc	gaaacaagaa	aaatgtttgc	ttagtttttt	6660
tatttcttaa	agggtatttg	ttttatttgt	agtcaattta	ttttatttca	ttttatatct	6720
aaattattaa	ataaaaaact	aaaatagagt	tttagtttta	aaaaatttag	agactaaaaa	6780
gaataaaaatg	gatgtactaa	aaattagtct	atagaacca	tcaaccctaa	atcctaatag	6840
gatgtactaa	taaaatggat	gaagtattat	ataggtcaag	ctatttgcaa	aaaaaacaga	6900
gagatcacat	gcacactaaa	aagataaaac	tgtagattgt	caaaatattc	aattatcctt	6960
tagaccatgt	ctagcagttt	atttatatga	tcctctaaaa	cactaatatt	attttagtat	7020
tatagactat	attatttcta	gagtgaagtt	taaatatatg	tatagagata	gataaactac	7080
acttcaaata	agtgtgacaa	aaaaatatgt	ggtaattttt	tatactttag	acatgcaatg	7140
ctctttatct	ctagagaggg	gcacgacgca	tgcaagcttg	ttaacgcggc	cgctaattct	7200
cagtccaaag	cctcaacaag	gtcagggtac	agagtctcca	aaccattagc	caaaagctac	7260
aggagatcaa	tgaagaatct	tcaatcaaag	taaactactg	ttccagcaca	tgcacatggg	7320
tcagtaagtt	tcagaaaaag	acatccaccg	aagacttaaa	gttagtgggc	atctttgaaa	7380
gtaatcctgt	caacatcgag	cagctggcct	gtggggacca	gacaaaaaag	gaatggtgca	7440
gaattgttag	gcccacctac	caaaagcctc	tttgccctta	ttgcaaagat	aaagcagatt	7500
cctctagtac	aagtggggaa	caaaataacg	tgaaaaagag	ctgtcctgac	agcccactca	7560
ctaattgcgta	tgacgaacgc	agtgacgacc	acaaaagaat	tagcttgagc	tcaggattta	7620
gcagcattcc	agattggggt	caatcaacaa	ggtacgagcc	atatcacttt	attcaaattg	7680
gtatcgccaa	aaccaagaag	gaactcccat	cctcaaaggt	ttgtaaggaa	ggtcgaggcc	7740
gctgtactgt	catattgtcg	tggtttttca	attgctgtac	ctgatgcaaa	cgtaaatgggt	7800
ttactaatct	tgaccccgcc	ggcttcaaaa	tgaagagtgc	taatttggtc	cacgtcacca	7860
tcaccggttc	gaactgtcta	gaatggcagg	caaagatgat	tggacaggca	tcaggggaaa	7920
aagagcaccg	ttgacgatgt	atgagagttc	ccaccattgc	gagcaatgat	tatcagccac	7980
acgacttact	cttcagagct	aaccactgcc	atgcagagaa	aaagtgaatc	atattgtcat	8040
gatctacaac	gaagtgaaac	aatcaggcat	gctaaagtgc	tgaaaactta	ctgatctctc	8100

-continued

---

atggttgaca	acaagaata	cggaataca	tcagcaacgc	aactcttgag	ctttgcttgc	8160
cgaatgacca	gctagaattt	ccaagcattt	acagaaacat	gactttaagt	ttcagaaaaa	8220
caaatacaag	gccactaaat	aagcgtgggg	ataacatata	ctccagatga	caggcaatct	8280
gcaacttgca	gccattcaaa	tgtacgatta	acaaaatatt	taagcgccac	atgagataat	8340
atatcctcca	attagggcct	ttagtattgt	cattagctca	taaccatggt	gcatcctcac	8400
atggacgctg	cataagaagt	tcataatagc	aacagacata	tgaacaaagc	atggtgcgcc	8460
tgcccggcgg	gactagctag	tactaccaat	catggaataa	gctagtaccc	taaatgaaat	8520
taaaatggtt	tttagcgatt	atccacgctg	tccagaatac	tctaateccac	aagttgaggg	8580
cgcccatgaa	gcccgagag	ggcgacgcca	tgtgtataaa	aggggcctaa	gctgagtgga	8640
cttctgcat	cagattagta	agcaatctca	agcgagaga	gccaaagctt	tcggtgtagc	8700
tcgaagagca	aagcgaaggc	aagcgcgccc	tgtaacctac	ctaccaacct	cgccattcct	8760
ctccaaactg	ttgtgctgct	gtctagatct	cccacactac	actagttact	cctcgtagat	8820
ctcggctacc	tggetcaaga	tcgggggtca	gatccgggtc	cggggatttt	ctttgtgccc	8880
tatggctgta	ttttggcgtc	tgtggctgat	gacagcgtgt	gttctcgagt	gcggatgcaa	8940
tctgagttat	ataggcaaat	ggccttgtca	actcgggcag	cggcattgct	ttgctcagtg	9000
tgtttgaatg	tgctgaaatt	catgtagtag	gctgtaggct	gtgcatttct	tgatttgcgt	9060
cttgcataat	tcactggtgg	attttctaaa	cctaacaagt	ttaaaattag	accattcaac	9120
caaagacagg	aggaataagt	gaagctggtg	tagtcacagc	ttatggccga	tccaaaattt	9180
gttaggaatg	tgaatatgtg	atgctacaaa	catatccttg	taagctacca	tgctatttat	9240
catgttccat	catggtgatt	ggtgagcact	catgaaaatt	tcagatccaa	acctagtgtt	9300
acatgtggat	ttgtgctctg	caatctatcg	ccagtaataa	aatggttgag	tgatccagct	9360
actacaaaat	cacattgcat	actttttttt	ttttagatt	atgcatcctg	gttttgggtg	9420
gtgggttcct	gatgtcagga	atataaattt	agcctgctga	tttaggtagc	actgccggtg	9480
cacactttgg	tttttgaata	cttgtagtct	tccagcttct	tgtagaactg	gtacaatgtg	9540
ggccatatat	aagaagggct	gtcaactagc	acatgctcac	taattagtct	aaacatttat	9600
gtttttattc	attcagggtca	ggtgcaatca	tagaagtagt	taatgacaat	actttagtgt	9660
ttctaataat	atttatgtat	ggactcaaat	taacatgcaa	aacatagtag	attagtggca	9720
tgcatctctt	ttcttaatag	tggaaaatac	gagataatga	taactgtgaa	gctctgttag	9780
tactcttcat	tactctatct	gagtggcagc	atatctcatg	ctagccataa	agcaagttct	9840
agacgtattc	tgttgttaat	tacttgtagc	tatataacct	aacctagtca	ttccagctta	9900
tgtctcttag	agatcatggt	tattagcacc	tcaagatttc	ctctgcacag	tatagtaact	9960
atcgaaaaag	atattatttc	tttgttttta	attgacaacc	ttcacgtgct	acttattttt	10020
gcagcttggc	cggccaccaa	ccatggctga	gatcaacaac	cagaaccagt	gcgtcccgtg	10080
caactgctcg	agcaacccta	aggagatcat	cctgggtgag	gaacgcctgg	agaccggcaa	10140
caccgtagcc	gacattagcc	tgggcctcat	caacttcttc	tacagcaact	tcgtgcccg	10200
cgggtggctc	atcgtgggcc	tcttgagct	tatctggggc	ttcatcgccc	cgtcccagtg	10260
ggacatcttc	ctcggccaga	tcgagcaact	gatcagccag	cggatcgagg	agttcctag	10320
gaaccaggcc	atctcccgcc	tggagggact	ctccaacctc	tacaaggtgt	acgtgcgcgc	10380
gttcagcgac	tgggagaagg	acccgaccaa	cccggccctc	cgcgaggaaa	tgctatcca	10440

-continued

---

gttcaacgat atgaactcgg cctcatcac cgccatcccg ctctcccgcg tgcagaacta 10500  
 cgaggtggcc ctctgtcccg tgtacgttca agccgccaac ctccacctct ccatacctccg 10560  
 cgacgtgagc gtgttcggcg agcgcgtggg ctacgacacc gccaccatca acaaccgcta 10620  
 ctccgacctc acctccctca tccacgttta caccaaccac tgcgtggaca cgtacaacca 10680  
 gggcctccgc cgctggagg gccgcttct ctccgactgg atcgtgtaca accgcttccg 10740  
 ccgccagctc accatctccg tctggacat cgtcgccttc tttcccaact acgacatccg 10800  
 cacctacctc atccagacc ccaccagct caccgcgag gtctacctc acctcccgct 10860  
 catcaacgag aacctcagcc cggccgccgt ctaccgacc ttctccgccc ctgagtcgca 10920  
 catcattcgc agcccgacc tctggactt cctcaactcc ttcacctct acaccgactc 10980  
 cctcgcgccg agcgcctact ggggcggta cctcgtgaac tcctccgca ccggcaccac 11040  
 taccaacctc atccgcagcc cgctctacgg ccgaggggg aacaccgagc gcccggtgac 11100  
 catcacgccg agcccgagcg tccccattt ccgaccctc agctacccca ccggcctgga 11160  
 caacagcaac cctgtggcgg gcatcgaggc cgtggagttc cagaacacca tctccaggag 11220  
 catctaccgc aagagcggcc ctatcgacag cttcagcgag ctgcctctc aggacgccag 11280  
 cgtgagccct gccatcggct acagccacag gctgtgccac gccaccttc tggagcgcat 11340  
 cagcggccct cgcacgcggg gcaccgtgtt ctcgtggacc caccgcagcg cctctcctac 11400  
 gaacgaggtg tctcctagtc gcatcaccca gatccctgg gtcaaggccc acaccctggc 11460  
 tagtggcgct agtgtcatca agggccctgg cttcaccggt ggtgacatcc tgaccaggaa 11520  
 ctctatgggc gagctgggca ctctgagggt cactttcact ggccgcctgc ctcagtctta 11580  
 ctacatccgc ttcgctacg ctagtctcgc taaccgctct ggtactttcc gctactctca 11640  
 gcctccgtct tacggtatct ctttcctaa gactatggac gctgggtgagc ctcgaccag 11700  
 taggagcttc gctcacacta ctctgttca cctatcact ttctctaggc ctcaggagga 11760  
 gttcgacctg tacatccagt cttggtgtga catcgacagg atcgagtca tccccgtgac 11820  
 cgccacgttc gagccgagc acgacctga gcgcgccag aaggtggtga acgcccctct 11880  
 cactagcact aaccagctag gcctgaagac tgacgtgacc gactaccaca tcgaccaagt 11940  
 gagcaacctc gtggcctgcc tctccgacga gttctgcctc gacgagaagc gcgagctgtc 12000  
 cgagaaggtg aagcacgcca agcgcctctc cgacgagcgc aacctgctcc aggaccccaa 12060  
 cttcaggggc atcaacaggc agcccagcc cggctggcgc ggctccaccg acatcacct 12120  
 ccagggcggt gacgacgtat tcaaggagaa ctacgttacc ctccccgca ctttcgacga 12180  
 gtgttaccoc acctacctc accagaagat cgacgagctc aagctgaagg cctacaccgc 12240  
 ctaccagctc cgcggctaca tcgaggactc ccaggacctg gaaatctacc tcatacctca 12300  
 caacgccaaag cacgagatcg tgaacgtgcc tggcaccggc agcctctggc ctctcagcgt 12360  
 ggagaaccag atcggccctt gcggcgagcc taaccgctgc gccctcacc tcgagtgga 12420  
 ccctgacctc cactgctcgt gcagggacgg cgagaagtgc gccaccata gccaccactt 12480  
 ctctctggac atcgacgtgg gctgcaccga cctgaacgag gacctggcg tgtgggttat 12540  
 cttcaagatc aagaccagc acggctcagc caggctgggt aacctggagt tccttgagga 12600  
 aaagcctctg ctgggtgagg cctggccag ggtcaagagg gctgagaaga aatggagggga 12660  
 taagagggag acctgcagc tggagaccac tategtctac aaggaggcta aggagtctgt 12720  
 cgatgctctg ttcgtcaact ctcagtacga tagactgca gctgatacca acatcgctat 12780  
 gatccacgct gcggataagc ggtccaccg gatccgggag gcttaccttc cggagcttcc 12840

-continued

---

tgteatcccg	ggtgtcaacg	ctgcatctt	cgaggaactt	gaggaacgga	tcttcaactgc	12900
gtttagtctt	tacgatgcgc	ggaacatcat	caagaacggg	gacttcaaca	atggtctgct	12960
gtgctggaac	gtcaaggggc	atgctcaggt	cgaggaacaa	aacaatcatc	gtagtgtcct	13020
tgteattcct	gagtgggagg	cggaggtctc	tcaagaggtc	cgtgtttgcc	cggggcgtgg	13080
gtacattctt	cgtgttactg	cgtacaagga	ggggtacggg	gaggggtgcg	ttactattca	13140
tgagattgag	aacaatactg	atgagcttaa	gttcaacaat	tgtgttgagg	aggaggttta	13200
cccgaacaat	actgttacgt	gcatcaacta	cacggcaacg	caagaggaat	acgaggggac	13260
gtacacctcg	cgtaatagag	ggtatgatga	ggcgtacgga	aacaaccctg	cggttccagc	13320
agattatgoc	tccgtttatg	aggagaagtc	gtacacggat	agacgacgcg	agaatccatg	13380
tgagtcaaat	cgaggatacg	gagattacac	accattacca	gcaggatcag	ttacaaagga	13440
gttgaatac	ttcccggaaa	cagataaagt	ttggattgaa	atcggagaaa	cagaagggaac	13500
attcatcgtc	gactcagtag	aattgttgtt	gatggaagaa	tgatagggac	cccggaccgc	13560
ccaaaacct	tgcaagact	atagtttggg	gtggagtata	cttggttgtg	tacatgctg	13620
cgtgttccat	tgtacacaca	aaacctagcc	acctcttgac	tcttgagtgt	atgcttgta	13680
tccgtgtgtt	gaagtttga	agaggcacca	tcaactataga	tgatggcttg	tgtccctctt	13740
tcatcaagat	tgaataatat	atgctacttt	gagagcgcta	tcctgcttgc	ctgattgtgt	13800
taatacttac	atccgtccca	cactcccaca	atataaggaa	ataaggtatt	ttggcagttt	13860
agagcaaaat	tcccttatat	ttttgggacg	gatgtcctct	tttctgcatt	tttttatggt	13920
catatgttoc	tgaagagtaa	ggtggatctt	gatcaacctg	tccgtttatg	gtgattgatt	13980
tgagtggaat	agaatgggcc	aacgtccggc	atacagttat	gcttcagtta	attaaaagta	14040
agggccccaa	gtaaggccgg	ccaagtaacg	gtccgaagta	aggcgcgggt	accgtcggtc	14100
cgggcctagt	aggccaagca	ggcgtggcgc	cgccaagaag	aacgattggc	aaacagctat	14160
tatgggtatt	atgggtaggc	acatgggaat	atagtgggag	gcagaggcgt	gcctgtgcct	14220
gctcttaatt	tgagctcttc	ccttgccct	gatagggcat	gtgcatgcgc	acaataatca	14280
ttggagctgc	ctgaatgatt	tgcctgcgct	cgtgtaaaac	gttcacggtc	catgccagtc	14340
caagaccgtc	ggcaagagat	gagggccatg	cgcagtagct	acatttgtat	acatatacat	14400
atTTTTTgg	atgatgcgat	gcgtgtgcat	atctggctgc	gccgctccct	cgcccatcgt	14460
cgctcgggtg	tacaacgaat	tttccgtggg	atgggctgaa	cttataatgg	gcctactggg	14520
cgggcctgat	gaaatagcac	acaattcacc	ctgcttttct	caccaggaat	gcattttcgc	14580
gggagccatc	gtcctaactt	ggcagacaga	cacacaaaca	caggacagga	gttcgccaac	14640
cccagcatga	gagaaaccta	tccggacagc	tcctccagcc	tatgggccgt	tgggccttca	14700
gcctcagcgc	cccggccccg	tgggcgggca	aaacactcca	cagtccacac	ggctgctgct	14760
gctgcccttg	ttccgtcgga	cctggggtaa	ctttatagct	atgatgttcg	cccttcccaa	14820
ccaatgtgcg	agcgttaagg	atgttaagta	acactagtta	ggattatgtg	gccaataacc	14880
tactttgcta	ggaacttgta	cacccctcat	aatatagaga	aaagaaagat	gccagtggtc	14940
caactctata	ccatgtgtct	cgtgtgtttg	ttatgtcgtg	cttatgtgaa	gggagacgag	15000
tcttctacag	cttctcacgc	ctaactctgt	gacgggagag	aagagagcgg	atctggtgat	15060
ccgtggtaac	gtagtcttca	acaaatgaca	cattgagcac	accgttgtca	tctcctcagc	15120

-continued

---

```

taaacattga gggggatttt tttttaagaa aggctgagat tcccttaocg tccctttggc 15180
agggctcgtc caacagcttc ttgaaaaatt ctataa                               15216

```

```

<210> SEQ ID NO 11
<211> LENGTH: 862
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(862)
<223> OTHER INFORMATION: 862 nucleotide sequence representing the 5'
flanking corn genomic DNA.

```

```

<400> SEQUENCE: 11
cgcagcgcag gtcctgtggt cctgttgtt ttggcttctg gtagcttctg gccacaaaaa   60
gctgtctcgg actgccaaac gctcagcttt tcagccagct tctataaaat ttgttggggg   120
gcaaaaacca tccaaaatca acataaacac ataatcagtt gagtcgtgtg aatagtagga   180
attcgtcact ttctagatcc tgagccctat gaacaatfff atcttctctc acacgtaatc   240
gtaatgatat tcagattctc cccacagcca gattctctc acagccagat ttcagaaaaa   300
gctggtcaga aaaaagctga accaaaacag ccttaggaca cacataggaa tggatcaaaa   360
aacatagcgc cgtaccggta cgtacgtgca gtcacagtca gctgctagct agctagctgc   420
gccggcggcg gcgttgctcg tcttgactca acgacgtccg cggctggcgc cgtcggccag   480
cggttgttg ccgcccgtcc catccggcgg agtccggctg ccagtggcgc gcggcggctt   540
cttcttcgtc gtcgtctctc cctttgccat ccccgtgcc acctggacga gggttctggc   600
tcccgtcgac gtcgacgaga gacgtggcgc cgcatttggc tgcccgcgta tcgtcgtcgt   660
cgtcgagatt gaaatcgaga gggagcgcgc gggatggctg ccgcccgttg cgcgacatgg   720
tctcgcgccg ccggcgcctg atcgacttct cttccgccgc cttegttagc tatagcttgg   780
tagtagtcaac caacaacgac ctaggagct agcgtacgtt cgaggtaacct gtctgcaag   840
caaatgtgtg gtagtgggtcc aa                                           862

```

```

<210> SEQ ID NO 12
<211> LENGTH: 1036
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1036)
<223> OTHER INFORMATION: 1036 nucleotide sequence representing the 3'
flanking corn genomic DNA.

```

```

<400> SEQUENCE: 12
acatgggaat atagtgggag gcagaggcgt gcctgtgcct gctcttaatt tgagctctc   60
ccctggccct gatagggcat gtgcatgcgc acaataatca ttggagctgc ctgaatgatt   120
tgcatgcgct cgtgtaaaac gttcacggtc catgccagtc caagaccgtc ggcaagagat   180
gagggccatg cgcagtacgt acatttgtat acatatacat atttttttgg atgatgcgat   240
gogtgtgcat atctggctgc gccgctccct cgtccatcgt cgtcgcggtg tacaacgaat   300
ttcccgctgg atgggctgaa cttataatgg gcctactggg cgggcccgtat gaaatagcac   360
acaattcacc ctgcttttct caccaggaat gcattttcgc gggagccatc gtctaaactt   420
ggcagacaga cacacaaaca caggacagga gttcgccaac ccagcatga gagaaacct   480
tccggacagc tctccagcc tatgggcccgt tgggccttca gcctcagcgc cccggcccgc   540

```

-continued

---

tgggcgggca aaacactcca cagtccacac ggctgctgct gctgcccttg ttcgctcgga	600
cctggggtaa ctttatagct atgatgttcg cccttcccaa ccaatgtgcg agcgtaag	660
atgttaagta aactagtagta ggattatgtg gccaaatacc tactttgcta ggaacttgta	720
caccctcat aatatagaga aaagaaagat gccagtggtc caactctata ccatgtgtct	780
cgtgtgtttg ttatgtcgtg cttatgtgaa gggagacgag tctctacag cttctcacgc	840
ctaactctgtt gacgggagag aagagagcgg atctgggtgat ccgtggtaac gtagttctca	900
acaaatgaca cattgagcac accgttgta tctcctcagc taaacattga gggggatttt	960
tttttaagaa aggctgagat tcccttacgc tccctttggc agggctcgtc caacagcttc	1020
ttgaaaaatt ctataa	1036

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 18376

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Sequence representing the transgene cassette comprised within the binary plasmid transformation vector used to transform corn to produce corn event MON 95379.

&lt;400&gt; SEQUENCE: 13

aaaagtccca tgtggatcac tccgttgccc cgctcgtcac cgtgttgggg ggaaggtgca	60
catggctcag ttctcaatgg aaattatctg cctaaccggc tcagttctgc gtagaaacca	120
acatgcaagc tccaccgggt gcaaacgggc agcggcgcca ggatatattc aattgtaaat	180
ggcttcatgt ccgggaaatc tacatggatc agcaatgagt atgatggta atatggagaa	240
aaagaaagag taattaccaa ttttttttca attcaaaaat gtagatgtcc gcagcgttat	300
tataaaatga aagtacatct tgataaaacg acaaattacg atccgctgta tttataggcg	360
aaagcaataa acaaattatt ctaattogga aatctttatt tcgacgtgtc tacattcacg	420
tccaaatggg ggcttagatg agaaaactca cgatcgatgc ggccaccacg agtcgacata	480
acttcgtata gcatacatta tacgaagtta tactagttgg tgttttctgt tacagagcaa	540
ctttatcata gttaggaact tgggtgtggc tctagccgct ccagtcgaa tgtttctaaa	600
ttaaaaatcc aaacaaaat cactgagggt tacttgggac acaatcttta cacagctacg	660
tattcacgcg ttctcttttc acacagctac gtattctggt gaggagcatt tacttaggaa	720
catcacttag ctcacctaag cagcagcat ctcaagattg cgaaccagag aatcagatat	780
cmetaactag tttcacgcca actaaaggtc acaattatc tcgagcaaaa aaacgatttc	840
aatgctatag acacaacttt acgaagttag ggcttctgtt tacgacaaac cacgaacttg	900
gatattaag atgccacaaa gaaggttcac ataaacaaca caggtcacag cagcgaagc	960
tacaagcaaa ccctacataa tcgaaaagca aatgtttttc aaaaccagac acacaaactg	1020
atagatagaa gccacacagg acagatcata ggatagcagc agtaatcaag cattgccaag	1080
gcaccaggca agaacctg agctcatcag gcagccttcg tatcggagag ttcgatcttc	1140
gcgccagcc cggccatcag gtccatgaac tccgggaagc tcgtggcgat catcgtggca	1200
tcgtccacgg tgacagggtt tccgacacg aggccatga cgaggaagct catggcgatg	1260
cggtgatcga gatgggtggc gacggcggcg cccgaggcgt tgccgagccc cttgcccgtca	1320
gggcggccac gcacgacgag cgacgtctcg cctcctcgc aatccacgcc attgagcttg	1380
agccattgg cgacggcca gagcggctcg ctttccttga cgcggagttc ttccagaccg	1440
ttcatcacgg tcgccccttc cgcgaaggcg gcggcgacag cgagaatcgg atattcgtcg	1500

-continued

---

atcatcgaag gcgcgcggtc ttcggcacc gtgacgcct tcagcgtgga ggagcgaacg	1560
cgcaggtccg ccacgtcttc gccgccgga aggcgcgggt tgatgacttc gatgtcggcg	1620
cccatttctc gcagcgtcag gatgagggcg gtgcccgggt ggttcatcag cactgtgagc	1680
atggtgacgt cggagcccg aacaagcagg gccgcaacca gcgggaaggc cgtcaggac	1740
gggtcgcgcg gcacgtcgt gacttggcgg gtgagcttgc cgcggccttc caggcggatg	1800
gtgcgcacgc cgtccgcac cgtctcgac gtaagggttg cgccaaagcc ctgcagcatc	1860
ttttccgtat gatecgcgt catgatcggc tcgatgaccg tcgtgatgcc gggcgtgtg	1920
aggccggcga gcagcacggc ggacttcacc tgtgcggagg ccatcggcac gcggtagggtg	1980
atcggcgtcg gcgtcttcgg cccgcgcaag gtaacgggaa gacggtcacc gtcttcgat	2040
ttcaactgca cgcctatttc gcgcagcggg ttcaacacgc ggccatcgg gcgctttgtg	2100
agcgaggcgt cgcgatgaa ggtgctgtcg aaatcgtaga ccccgacgag gcccatcgtc	2160
aggcggcagc ccgtggcggc attgccgaaa tcgagcggcg cctcaggcgc caggaggccg	2220
ccattgcccga cgcctcgtat gatccagggt tcgcttctc tacggatgcg ggcgcccac	2280
gcctgcattg ccttgccgt attgatgac tcctcgcct ccagaaggcc ggtgatgccc	2340
gtttcaccgc tcgcgagacc gccgaacatg aaggaccggt gggagatcga cttgtgcgcg	2400
ggaatgcgga cggttccgga aaggccagag gatttgcggg cggttgcggg ccggtcgtt	2460
gcaccgtgaa gcatgcacgc cgtggaaca gaagacatga ccttaagagg acgaagctca	2520
gagccaatta acgtcatccc actctcttc aatccccacg acgacgaaat cggataagct	2580
cgtggatgct gctgcgtctt cagagaaacc gataagggag atttgcgttg actggatttc	2640
gagagattgg agataagaga tgggttctgc acaccattgc agattctgct aacttgccc	2700
atggatcagg ctgcatagga caaaacaaga tttcaacaaa ttacgcctta agaattttt	2760
ttatcacagg atttatcaag acaagacgaa caggagtcga caggatgaacc aatccacata	2820
atacagcctt tcagatagcg ttctaatac ctttaagaat ataatagcac ctatcttcag	2880
atcagatcaa aagattacta cgcagaaacg agcactacaa agcatgatac caaagaccat	2940
caaatagttg ccaatctgt ctaaatattt ttttctgtgc ttgctctgaa atagttctta	3000
gctcgcagca atgtacatac tacagatcca catactcgt ccatacatgt gcctaagtta	3060
cagatctatt cataataagc aacaaattaa gtataaacgg gagtacaacat ctagggtgac	3120
agcagtaaaa tatcaagac catcaaccaa aggggcattt cagcacggat catggcaatc	3180
caaccagatc cgcattgcat tcctatacaa cctaggtaga aaatttgaaa aaacaaaca	3240
gcattgatca gatctatccg aaacaaacat aactaacca gatctacca aaaggaacag	3300
attatcagca gggagacaga ctaattcagt ggatcccat ccgaaacccc cctaatttg	3360
catcatttcc cgattcttag cgtctcactg ccagatccaa aacatttcgc cagatccacg	3420
aactaccact gatcaaacg cagatccgga catcgacaac cacagaaggc cagatccacc	3480
gacagatcta tcaaacagaa cgaaccacac cccaggatca cccagatcca cccacagatc	3540
gaagcgcgt actacgagaa acggaggaat ggaagcatca gatccgtttt tgtttgttt	3600
acctggatgg ctacccgggt gcggcgggaga cgaacggcgg cgagcggcgg cggaggagg	3660
cgcgagaga ggcgagtacg aagacgcctt tctctcgggt gtggtagggt ttccacctcc	3720
cactcctccc ctttctatag agcgggaggc gcctcggcgt ttttgcgttt aacccccctc	3780
attaagtctt ccaattttac actccgctcc actttgctc ctccaacggg caggaaactgt	3840

-continued

---

actggtttgg aatatcgga ttttctgag ggttttagtg caaaatattc atgccgtttt	3900
gtttagtggt ggtgggtaag agtccagatc taccgcgcgc tcaactgggag gtgggcccac	3960
tggttgctgc gagcgactga cgtctcgctt gtaggtgggc ccacaagggt gggaccggat	4020
tgtcagtggc gtgtgggttg gtgggcgtag ttggggtaga attcagagggg ttgttccgtc	4080
atthtccgcc cgctccgtcg gcggccagtc gtgccgatat ttttggattt tgaattcaa	4140
gagcgggagc caggaggggg aggggtgggg caggcctggc ccatgatcta ttgtttgctc	4200
ggatgggcct tgggcctaac tgttagtttt gattctgttt gtaattaact ctttttga	4260
ggaataaaaa gaacctgct acctgacggg ttgatgtttt accattgaga tttcaaagg	4320
gttaaagtga aattgaaact caattcgaac atgattttta ttgttttgaa ctcccttca	4380
attgagggca gtttttagtg ttataaatct tcatcgagag aatthttctg atgtttttat	4440
ggaatttgat gtcctaagag cacctctaac aggcaacatc ctctccaaat ggctctcaaa	4500
ctatthtttt ttgacaattc tagtaaaaag atactacccc aataacctt ccaaccggat	4560
ggccaaatcc ctccccttat tggcctaact tggcaaccca aaataacttg ccaacggta	4620
ggccccatgt actatctctc ctcccacat ggctactctc ctcttctctc tctctcttat	4680
gttgacggag agggggcgtt gacgggtggc gttggggcaa gctagggtag tgcgggtgtg	4740
gaggtgaagc ggtggctgtg ctaaagctcg cccctggcat tgggtgtgtg ggagagctgg	4800
ccagtggcta gtgtgacaat ggaggtgacc aggcgacact ttctccctcc ctcttcctc	4860
catgttgatg agaagcatgt tgtgtccct aggataactt cgtatagcat acattatacg	4920
aagttatagc ggccgcaaat caacctcact ctatttaaat gaggtggtag gatttgccgc	4980
gtgagaccta ttaccctggt atccctagtt taaacacctg caggaaaaga ttaattaaca	5040
togatcatca tgttcaattg acagcaaat aaactagtta tataatgggt tcatgagatg	5100
agagtagcta attattaaca caagtggaag aaacttttat ttgacgggat acaaataaca	5160
agatacagag atcaggctga tggatcaatc atatcatatc agtgatagat cgatcgttcg	5220
acacatatat atgtatgaca aaacatagta taagtacgtg cgaaagggtt tatatatata	5280
tggatattga tgcattggtt aatggtatgg taattaaggg atcggaggat cgattaatgg	5340
gtccctatca ctctccatc agtagcagct cgaccgagtc cactgatgaa gtgccctccg	5400
tctcgccgat ctcgatccag actttgtcgg tctccgggaa gtaactccagc tctttgtga	5460
cgtaccggc gggcagcggg gtgtagtca cgtagccacg gttggactcg caagggttgt	5520
ccctccgtcc gtcgggtgac gccttctcct cgtaagcggg cgcataatcc gcaggcacgc	5580
tgtgtttgga ctctgacgag ccatcatagc cgcgattccg ggaagtgtaa gtgccctcgt	5640
actcctcttg cgtggcagtg tagtcgttgc aagtaacggg gttgttcggg tacacctcct	5700
cctccacaca gttcagagaac ttgagctcgt cgggtgtgtt ctcgatctcg tggatagtea	5760
cgcagccttc accgtagcct tctttgtacg cagtcacacg gaggatgtag ccgagaccag	5820
ggcagacgag gacctcctgg ctacctctg cctcccactc cggcacgacc agcacggacc	5880
ggtggttgtt ctgctcctcg acgtccacat ggccttgac gttccagcaa ctacagccgt	5940
tgttgaagtc accgttctta ataacattcc tggcatcgta caggctgaat gctgtgaaga	6000
tgcggccctc cagttcctcg aagatcgccg cgttcacacc agggatcact gacagctcgg	6060
gcaggtaacg ctacaggatg ctgtggaccc gcttgcagc cgcgtggatc atggcgatgt	6120
tgggtgcagc ctggagggcg tcgtaactggc tgttgacgaa gagtgcgtcc actgactcct	6180
tggcctcctt gtacacgatg ttggtctccc actccagett ctcccgttg tctcgccatt	6240

-continued

---

tcttctcggc	gcgcttcaca	cgagcgaggg	cctcgccac	gagaggcttc	tctcaagga	6300
actccaggtt	gccgagggcg	gcattgccc	cctgggtctt	aatcttgaag	ataaccaga	6360
caccaggtc	ctcgttgagg	tcgggtgcgc	ccacgtcgat	gtccagagag	aagtgggtgc	6420
agtggtagc	gcactttccg	attggagagg	gagcgctcaa	tggccaaagt	gagcccgctc	6480
ctggcacgtt	cactgtctcg	tgcttggcgt	tgtatcgat	caggtagatt	tccaggtcct	6540
gactatcctc	gatgtaccgc	cgaagctggg	agcgggtgta	cgcttcaat	ttactctcgt	6600
ctatctctg	gtagagatac	gtcgggtagc	attcgtcgaa	agtcccgagc	agcgtcacat	6660
agttctcctt	gaacacatcg	tcgcctccct	ggatgggtgat	gtctgtactc	ccgcgccaac	6720
cacggtcgag	ctggcgggtg	atgccacgga	agttcgggtc	ctggagcaaa	ttccgctcgt	6780
cactcagacg	cttcgcgtgt	ttgaccttct	cggacagctc	cttctctcgc	tcgaggcaga	6840
actcgtcggg	caggcactcc	acaaggttgg	acacttggtc	aatgtgatag	tccgtgacgt	6900
cggctcttgg	cccaatctgg	ttgctgctgg	tgaacagttc	gtagacagcc	ttctgagccc	6960
gctccaggtc	agactcggcc	tcgaacgtcg	catccgccag	gatgatctca	atcttgtcga	7020
tgtacagttc	gccagagctg	atggagccag	caccgaacag	cggttgctcg	gagatcccaa	7080
tgatgtcagg	gttcgctctg	aatgagaaa	ggttcgagaa	atcagtgtac	cggaagggtc	7140
ggctagtcaa	gttctcgctc	atctccatcg	tcttctggag	aggcatgttg	actgagactt	7200
ggcctccac	gcctgtggac	gcggctcccg	tcaggacaat	gaccctcgcg	tcgcggcttg	7260
aagcgtagcg	gaacctcaga	cggtaacgct	gggtgatcgg	cgagttgatg	ttcacttgca	7320
acgaaacgaa	gtcgcgcaaa	gtgttgccgc	gcagtatgtc	tccaccgggtg	aagcccgagc	7380
cgggtgatgac	gctcgtgcca	ccccacacac	ggaagccctt	cacgagcggg	atctggttaa	7440
tggagtcgga	agagatcgtg	ttggctccgt	cggcgcctcg	gtgtgtccag	gagtacaccg	7500
gcgcacgcag	cgtgttgcca	gagatgagcc	ggatgttggg	caggcgggtg	ctgtaggact	7560
cgtagttcgg	gcgttccgtg	gtctcggcgc	gcagctcagt	ttcactgtcg	aagagctggg	7620
tgccagcgc	cgtgtaccgc	atgggtgtaga	gcaatgaccc	gcgcaggag	ttcaacgggt	7680
taccgcagtt	gaaccgggcc	cacgggacgc	cgttgacggg	cgttgcaga	aggatgttaa	7740
tgcccgcaaa	cgactcagtc	cggtaaacgt	cgcggctcgt	gaactgtagc	gtaactgggt	7800
tgatggaggt	gttgggtgtg	ccgtgggtac	tgggtctcaa	gctgccacgg	atcgtcttac	7860
tctccaatcg	gtgcccagcc	cagtagttca	tgtactgtgt	gttggaccag	cgtgagagca	7920
cggagaagat	cgtgagctgc	tcgggaaagt	cgagcaggtg	aggaggcggg	atgacggcgg	7980
cctcgattgc	cgagaaggag	ggtgcggttg	tgttgaacca	attcgtggat	gcgaaaccgg	8040
agggtagcgt	cgtccggcgc	attgggtctg	tgtagatctc	ccgcgtgagt	tgcgcgctag	8100
tgttcattgg	gtacaccgct	gtgtcgtaac	acgggaatag	cgccaccagg	tctaggacac	8160
ccaaagtcaa	gtcgcggcgg	aactgggttg	aacggagcca	tgactcagcg	ttggtgccgc	8220
gcaggttgtt	gagccctgtg	ttgtaccatc	gagcgcagta	gtcactgtac	tcacgggtct	8280
tctccacttg	gcgctcgtag	taacgtggga	tttcttgaga	tgtgagtcgc	aactcggagc	8340
cgaacaggct	ggcgtcccg	agcaggagca	gatgtaagtt	ggcggcctgg	gcgtacacca	8400
taaggagtgg	aacttcctgg	ttcctgatgg	cgaacagcgg	catagcgttc	aggaagtcca	8460
gttcgagagc	gatgtactgg	gtgtagagta	ccgagcgtgt	tcgcgcgtca	tccctgttct	8520
ccaaccagtc	ctccagtgac	tgctgggtatg	cccgaagga	gtttcccagg	ccctgtagcc	8580

-continued

---

tggccagagc	agtatcggcg	gtgttctccg	tgacttgctg	ccggatcaac	tgctccacat	8640
gctccaggaa	gatctcccac	gggtcacgac	cgcgaggcca	cagctcgccc	acaaggaacg	8700
agtagaagga	cgcgatctga	ccggcgaatg	ggacaccgag	gacgccgagg	atacggcccg	8760
cgatgttgat	gccggctctga	accgtgctgg	ccgacacgaa	tgggtcgatg	ttgttgccct	8820
cggctatgca	gagggagtc	tcgatgcgcg	cgctcggtga	caggttcac	tgccgggagt	8880
ggtttgaaac	agcagggatc	gagagggcgt	tgatgatctc	gttctcgttc	ttccggttgc	8940
tcgtcatggt	ggtggccggc	caagtaacgg	tccgacctgc	agaagtatca	ccaaacaaca	9000
gggtgagcat	cggacaaaag	aaacagtacc	aagcaataa	atagcgtatg	aaggcagggc	9060
taaaaaatcc	acatatagct	gctgcatatg	ccatcatcca	agtatatcaa	gatcaaaaata	9120
attataaaac	atacttgttt	attataatag	ataggtactc	aaggtagag	catatgaata	9180
gatgccgcat	atgccatcat	gtatatgcat	cagtaaaacc	cacatcaaca	tgtataccta	9240
tcctagatca	atatttccat	ccatcatctt	aaactcgtaa	ctatgaagat	gtatggcaca	9300
cacataaaga	tacaaaatta	ataaatccac	caggtagttt	gaaacagtat	tctactccga	9360
tctagaacga	ccgcccaacc	agaccacatc	atcacacca	agcgaaaaaa	agcatctctg	9420
tatatgcatc	agtaaaacc	gcatcaacat	gtatacctat	cctagatcga	tatttccatc	9480
catcatctta	aaactgtaac	tatgaagatg	taggcacata	catacagatc	ctttaataaa	9540
tccaccaggt	agtttgaaac	agtattctac	tccgatctag	aacgacctt	tatcctttaa	9600
taaatccacc	aggtagttt	aaacagtatt	ctactccgat	ctagaacgac	cgcccaacca	9660
gaccatatca	tcacaaccaa	gcgagaaaa	aagcatctct	gtatatgcat	cagtaaaacc	9720
cgcatacaaca	tgtataccta	tcctagatcg	atatttccat	ccatcatctt	aaactcgtaa	9780
ctatgaagat	gtatggcaca	cacatacaga	tacaaaatta	ataaatccac	caggtagctt	9840
gaaacagtat	cctactccga	tctagaacga	ccgcccaacc	agaccacatc	atcacaaaca	9900
tgaacatgaa	catgtttgct	ctaacacaaa	catgaacaga	agtagaacta	ccgggcccta	9960
accatggacc	ggatgcgcga	tctagagaag	gtagagagag	gggggggggg	aggatgagcg	10020
gcgtaccttg	aagcggaggt	gccgacggct	ggattgggg	gagatctggt	tgctgtgtg	10080
tgctctccga	acgaacacga	ggttggggaa	agaggggtg	gagggggtg	ctatttatta	10140
cggcggggcga	ggaagggaaa	gcgaaggagc	ggtgggaaag	gaatcccccg	tagctgccgg	10200
tgccgtgaga	ggaggaagag	gccgcctgcc	gtgccgcctc	acgtctgccg	ctccgccacg	10260
caatttctgg	atgccgacag	cggagcaagt	ccaacgggtg	agcggaaactc	tcgagagggg	10320
tccagaggca	gctacagaga	tgccgtgccg	tctgcttcgc	ttggcccgc	gcgacgctgc	10380
tggttcgctg	gttggtgtcc	gtagactcgc	tcgacggcgt	tgaacagcct	gtcattatct	10440
actcgaaaaca	agaaaaatgt	ttgcttagtt	tttttatttc	ttaaagggtg	ttgtttttat	10500
ttgtagtcaa	tttattttat	ttcattttat	atctaaatta	ttaaataaaa	aactaaaata	10560
gagttttagt	tttaaaaaat	ttagagacta	aaaagaataa	aatggatgta	ctaaaaatta	10620
gtctatagaa	accatcaacc	ctaaatccta	aatggatgta	ctaataaaa	ggatgaagta	10680
ttatataggt	caagctat	ttgcaaaaaa	cagagagatc	acatgcacac	taaaaagata	10740
aaactgtaga	ttgtcaaaat	attcaattat	cctttagacc	atgtctagca	gtttattttat	10800
atgatcctct	aaaacactaa	tattattttta	gtattataga	ctatattatt	tgtagagtga	10860
agtttaata	tatgtataga	gatagataaa	ctacacttca	aataagtggtg	acaaaaaaat	10920
atgtggtaat	tttttatact	ttagacatgc	aatgctcttt	atctctagag	aggggcacga	10980

-continued

---

cgcattgcaag cttgttaacg cggccgctaa ttctcagtc aaagcctcaa caaggtcagg 11040  
 gtacagagtc tccaaacccat tagccaaaag ctacaggaga tcaatgaaga atcttcaatc 11100  
 aaagtaaaact actgttcacg cacatgcac atggtcagta agtttcagaa aaagacatcc 11160  
 accgaagact taaagttagt gggcatcttt gaaagtaac ttgtcaacat cgagcagctg 11220  
 gcttggtggg accagacaaa aaaggaatgg tgcagaattg ttaggcgcac ctacaaaag 11280  
 catctttgcc tttattgcaa agataaagca gattcctcta gtacaagtgg ggaacaaaat 11340  
 aacgtgaaa agagctgtcc tgacagccca ctactaatg cgtatgacga acgcagtgac 11400  
 gaccacaaaa gaattagctt gagctcagga tttagcagca ttccagattg ggttcaatca 11460  
 acaaggtagc agccatatca ctttattcaa attggtatcg ccaaaaccaa gaaggaactc 11520  
 ccactctcaa aggtttgtaa ggaaggtcga ggcgctgta ctgtcatatt gtcgtggttt 11580  
 ttcaattgct gtacctgatg caaacgtaac gggtttacta atcttgacc cgccgcttc 11640  
 aaaatgaaga gtgctaattt ggtccacgct accatcaccc gttcgaactg tctagaatgg 11700  
 caggcaaaaga tgattggaca ggcattgcagg gaaaagagc accgttgacg atgtatgca 11760  
 gttcccacca ttgcgagcaa tgattatcag ccacacgact tactcttcag agctaaccac 11820  
 tgccatgcag agaaaaagtg aatcatattg tcatgatcta caacgaagtg aaacaatcag 11880  
 gcattgctaaa gtgctgaaac tttactgatc tctcatggtg gacaacaaag aatcgggaa 11940  
 tacatcagca acgcaactct tgagctttgc ttgccgaatg accagctaga atttccaagc 12000  
 atttacagaa acatgacttt aagtttcaga aaaacaaata caaggccact aaataagcgt 12060  
 ggggataaca tatctccag atgacaggca atctgcaact tgcagccatt caaatgtacg 12120  
 attaacaaaa tatttaagcg ccacatgaga taatatatcc tccaattagg gcctttagta 12180  
 ttgtcattag ctacatacca tgggtgatcc tcacatggac gctgcataag aagttcataa 12240  
 tagcaacaga catatgaaca aagcatggtg cgcctgccc gccggactag ctagtactac 12300  
 caatcatgga ataagctagt accctaaatg aaatataat ggtttttagc gattatccac 12360  
 gccgtccaga atactctaat ccacaagtg aggcgcccga tgaagccgag agaggcgac 12420  
 gccatgtgta taaaaggggc ctaagctgag tggacttgct gcattcagatt agtaagcaat 12480  
 ctcaagcgca gagagccaaa gctttcgggt tagctcgaag agcaaagcga aggcaaggcg 12540  
 cgcctgtaac ctacctacca acctcgccat tctctccaa actgttgctg tgctgtctag 12600  
 atctcccaca ctacactagt tactctcgt agatctcggc tacttgctc aagatccggg 12660  
 gtcagatccg ggtccgggga tttctttgt gccctatggc tgtattttg cgtctgtggc 12720  
 tgatgacagc gtgtgtctc gagtgccgat gcaatctgag ttatataggc aaatggcctt 12780  
 gtcaactcgg gcagcggcat tgtttgctc agtgtgttg aatgtgctga aattcatgta 12840  
 gtaggctgta ggctgtgcat ttcttgattt gcgtcttgca taattcactg gtggattttc 12900  
 taaacctaac aagtttaaaa ttgaccatt caaccaaga caggaggaat aagtgaagct 12960  
 gttgtagca cagcttagg ccgatccaaa atttgttagg aatgtgaata tgtgatgcta 13020  
 caaacatata cttgtaagct accatgctat ttatcatggt ccatcatggt gattgggtgag 13080  
 cactcatgaa aatttcagat ccaaacctag tgttacatgt ggatttgctc tctgcaatct 13140  
 atcggcagta ataaaatggt tgagtgatcc agctactaca aaatcactt gcatactttt 13200  
 tttttttgta gattatgcat cctggttttg ggtggtgggt tctgatgctc aggaataaa 13260  
 atttagcctg ctgatttagg tagcactgcc ggtgcacact ttggtttttg aatcttga 13320

-continued

---

gtcttccagc ttcttgtaga actggtacaa tgtgggcat atataagaag ggctgtcaac 13380  
tagcacatgc tcactaatta gtctaaacat ttatgttttt attcattcag gtcaggtgca 13440  
atcatagaag tagttaatga caatacttta gttgttctaa tattatttat gtatggactc 13500  
aaattaacat gcaaaacata tgagattagt ggcatgcatt ctttttctta atagtggaaa 13560  
atacgagata atgataactg tgaagctctg ttagtactct tcattactct atttgagtgg 13620  
cagcatatct catgctagcc ataagcaag ttctagacgt attctgtgtg taattacttg 13680  
tagctatata acccaaccta gtcattccag cttatgtctc ttagagatca tgtttattag 13740  
cacctcaaga tttcctctgc acagtatagt aactatcgaa aaagatatta tttctttgtt 13800  
tttaattgac aaccttcacg tgctacttat ttttgacgtg tggcgggcca ccaacctggt 13860  
ctgagatcaa caaccagaac cagtgcgtcc cgtacaactg cctgagcaac cctaaggaga 13920  
tcatcctggg tgaggaacgc ctggagaccg gcaaacaccg agccgacatt agcctgggcc 13980  
tcatcaactt cctctacagc aacttcgtgc ccggcgggtg cttcatcgtg ggccctctgg 14040  
agcttatctg gggcttcate ggcccgtccc agtgggacat cttcctcggc cagatcgagc 14100  
aactgatcag ccagcggatc gaggagtctg cttaggaacca ggccatctcc cgccctggag 14160  
gactctccaa cctctacaag gtgtacgtgc gcgcgttcag cgactgggag aaggaccoga 14220  
ccaaccggc cctccgcgag gaaatgcgta tccagttcaa cgatatgaac tcggccctca 14280  
tcaccgcat cccgctcttc cgcgtgcaga actacagagt ggccctcctg tccgtgtacg 14340  
ttcaagccgc caacctccac ctctccatcc tccgcgacgt gagegtgttc ggcgagcgt 14400  
ggggctacga caccgccacc atcaacaacc gctactccga cctcacctcc ctcatccacg 14460  
tttacaccaa cactgcgtg gacacgtaca accagggcct ccgcgcctg gagggcgcgt 14520  
tcctctccga ctggatcgtg tacaaccgct tccgcgcca gctcaccatc tccgtcctgg 14580  
acatcgtcgc cttctttccc aactacgaca tccgcaccta ccctatccag accgccacc 14640  
agctcaccgc cgagggtctac ctgcacctcc cgttcatcaa cgagaacctc agcccgccgc 14700  
ccgtctacc gacctctcc gccgctgagt ccgcatcat tcgcagccc caccctcgtg 14760  
acttctcaa ctccttcacc atctacaccg actccctcgc ccgcagcgc tactggggcg 14820  
gtcacctcgt gaactcctc cgcaccggca ccaactacaa cctcatcgc agcccgctct 14880  
acggcgcga gggcaacacc gagcgcggg tgaccatcac cgcagccc agcgtgccca 14940  
tcttccgac cctcagctac cccaccggcc tggacaacag caaccctgtg gcgggcatcg 15000  
agggcgtgga gttccagaac accatctcca ggagcatcta ccgcaagagc ggccctatcg 15060  
acagctcag cgagctgcct cctcaggacg ccagcgtgag ccctgccatc ggctacagcc 15120  
acaggctgtg ccacgccacc ttctggagc gcatcagcgg ccctcgcac gcgggcaccg 15180  
tgctctcgtg gacccaccgc agcgcctctc ctacgaacga ggtgtctcct agtcgcatca 15240  
cccagatccc ttgggtcaag gccacaccc tggctagtgg cgctagtgtc atcaagggcc 15300  
ctggcttcac cgggtgtgac atcctgacca ggaactctat gggcgagctg ggcactctga 15360  
gggtcactt cactggcgcg ctgcctcagt cttactacat ccgcttcgcg tacgctagt 15420  
tcgctaaccg ctctggtact ttccgctact ctacgcctcc gtcttaagggt atctctttcc 15480  
ctaagactat ggaagctggt gagcctctga ccagtaggag cttegcctcac actactctgt 15540  
tcaactctat cactttctct agggctcagg aggagtcca cctgtacatc cagtctggtg 15600  
tgtacatcga caggatcgag ttcacccccg tgaccgccac gttcggggc gagtacgacc 15660  
ttgagcgcgc ccagaaggtg gtgaacgcc tcttactag cactaacccg ctaggcctga 15720

-continued

---

agactgacgt gaccgactac cacatcgacc aagtgagcaa cctagtggcc tgcctctccg 15780  
 acgagttctg cctcgacgag aagcgcgagc tgtccgagaa ggtgaagcac gccaaagccc 15840  
 tctccgacga gcgcaactcg ctccaggacc ccaacttcag gggcatcaac aggcagcccc 15900  
 accgcgctg gcgcgctcc accgacatca ccatccaggg cggtgacgac gtattcaagg 15960  
 agaactacgt taccctccc ggcacctcg acgagtgtta cccacactac ctctaccaga 16020  
 agatcgacga gtccaagctg aaggcctaca cccgctacca gctccgccc tacatcgagg 16080  
 actcccagga cctggaaatc tacctcatcc gctacaacgc caagcacgag atcgtgaacg 16140  
 tgcttgacc cggcagcctc tggcctctca gcgtggagaa ccagatcggc ccttgccggc 16200  
 agcctaaccg ctgcgcccct cacctcgagt ggaaccctga cctccactgc tcgtgcaggg 16260  
 acggcgagaa gtgcgcccac catagccacc acttctctct ggacatcgac gtgggctgca 16320  
 ccgacctgaa cgaggacctg ggcgtgtggg ttatcttcaa gatcaagacc caggacggtc 16380  
 acgccaggct gggtaactcg gagttccttg aggaaaagcc tctgctgggt gaggcctcgg 16440  
 ccagggtcaa gagggctgag aagaaatgga gggataagag ggagaccctg cagctggaga 16500  
 ccactatcgt ctacaaggag gctaaggagt ctgtcgatgc tctgttcgtc aactctcagt 16560  
 acgatagact gcaagctgat accaacatcg ctatgatcca cgctcgggat aagcgggtcc 16620  
 accggatcgg ggaggcttac cttccggagc tttctgtcat cccgggtgtc aacgctcga 16680  
 tcttcgagga acttgaggaa cggatcttca ctgcgtttag tctttacgat gcgcggaaca 16740  
 tcatcaagaa cggggacttc aacaatggtc tgctgtgctg gaacgtcaag ggatcatgctg 16800  
 aggtcgagga acaaaacaat catcgtagtg tccttgcctc tcctgagtgaggaggcgagg 16860  
 tctctcaaga ggtccgtggt tgcccgggce gtgggtacat tcttcgtggt actgcgtaca 16920  
 aggaggggta cggggagggg tgcggtacta ttcatgagat tgagaacaat actgatgagc 16980  
 ttaagttaa caattgtggt gaggaggagg tttaccgaa caatactggt acgtgcatca 17040  
 actacacggc aacgcaagag gaatacaggg ggacgtacac ctccgtaat agagggtatg 17100  
 atgaggccta cggaaacaac ccgtcgggtc cagcagatta tgccctcgggt tatgaggaga 17160  
 agtcgtacac ggatagacga cgcgagaatc catgtgagtc aaatcgagga tacggagatt 17220  
 acacaccatt accagcagga tacgttaca aggagttgga atacttccc gaaacagata 17280  
 aagtttgat tgaatcgga gaaacagaag gaacattcat cgctgactca gtagaattgt 17340  
 tgttgatgga agaatgatag ggaccccga cccgccaaa ccattgcaa gactatagtt 17400  
 tgggtggag tatacttggg tgtgtacatg cctgcgtggt ccattgtaca cacaaaacct 17460  
 agccacctct tgactcttga gtgtatgctt gttatccgtg tggtgaagtt tgtaagaggc 17520  
 accatcacta tagatgatgg cttgtgtccc tctttcatca agattgaata atatatgcta 17580  
 ctttgagagc gctatcctgc ttgcctgatt gtgtaatac ttacatccgt cccacactcc 17640  
 cacaataaa ggaataaagg tattttggca gtttagagca aaattccctt atatttttgg 17700  
 gacggatgtc ctctttctg cattttttta tgttcatatg ttectgaaga gtaaggtgga 17760  
 tcttgatcaa cctgtcgggt tatggtgatt gatttgagtg gaatagaatg ggccaacgtc 17820  
 cggcatacag ttatgcttca gtttaataaa agtaagggtc ccaagtaagg ccggccaagt 17880  
 aacggtccga agtaaggcgc gggtaaccgtc ggtccgggccc tagtaggcca agcaggacgt 17940  
 ggccgcgcaa gaagaacgat tggcaaacag ctattatggg tattatgggt aggcctgccc 18000  
 gggctcagcc taagtcgcta ccttaggacc gttatagtta cgggaaact atcagtgttt 18060

-continued

---

```

gacaggatat attggcgggt aaacctaaga gaaaagagcg tttattagaa taatcggata 18120
tttaaaaggg cgtgaaaagg tttatccgtt cgtccatttg tatgtgcatg ccaaccacag 18180
ggttcccctc gggagtgcct ggcattccgt gcgataatga cttctgttca accaccctaaa 18240
cgtcggaaaag cctgacgacg gacgagcatt ccaaaaagat cccttggtc gtctgggtcg 18300
gctagaaggt cgagtgggct gctgtggctt gatccctcaa cgcggtcgcg gacgtagcgc 18360
agcgccgaaa aatcct 18376

```

```

<210> SEQ ID NO 14
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Enterobacteria phage P1
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(35)
<223> OTHER INFORMATION: 35 nucleotide sequence representing the LoxP
sites used for Cre-mediated excision and recombination.

```

```

<400> SEQUENCE: 14

```

```

ataaactcgt atagcataca ttatacgaag ttata 35

```

```

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 20 nucleotide sequence corresponding to a
thermal amplification primer referred to as SQ51219 used to
identify corn event MON 95379 DNA in a sample.

```

```

<400> SEQUENCE: 15

```

```

cctgcaagca aatgtgtggt 20

```

```

<210> SEQ ID NO 16
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 30 nucleotide sequence corresponding to a
thermal amplification primer referred to as SQ21524 used to
identify corn event MON 95379 DNA in a sample.

```

```

<400> SEQUENCE: 16

```

```

gtcgttttat caaaatgtac tttcatttta 30

```

```

<210> SEQ ID NO 17
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 16 nucleotide sequence corresponding to a probe
referred to as PB10269 used to identify corn event MON 95379 DNA
in a sample.

```

```

<400> SEQUENCE: 17

```

```

taacgctgcg gacatc 16

```

```

<210> SEQ ID NO 18
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 24 nucleotide sequence corresponding to a
thermal amplification primer referred to as SQ20222 used as an
internal control for the event and zygosity assay for MON 95379
and hybridizes to a region of the corn genome.

```

-continued

---

<400> SEQUENCE: 18

gccctatgac ttaccgagag ttca 24

<210> SEQ ID NO 19

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 28 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ20221 used as an internal control for the event and zygosity assay for MON 95379 and hybridizes to a region of the corn genome.

<400> SEQUENCE: 19

gttgctatgt actaacagaa ctgcatgt 28

<210> SEQ ID NO 20

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 29 nucleotide sequence corresponding to a probe referred to as PB50237 used as an internal control for the event and zygosity assay for MON 95379 and hybridizes to a region of the corn genome.

<400> SEQUENCE: 20

ttgttggtg gctccattct gacttgta 29

<210> SEQ ID NO 21

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 20 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50998 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1B.868.

<400> SEQUENCE: 21

cgggtagcat tcgctgaaag 20

<210> SEQ ID NO 22

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 20 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50997 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1B.868.

<400> SEQUENCE: 22

ggcgacgatg tgttcaagga 20

<210> SEQ ID NO 23

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 18 nucleotide sequence corresponding to a probe referred to as PB50340 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1B.868.

-continued

---

<400> SEQUENCE: 23  
aactatgtga cgctgctc 18

<210> SEQ ID NO 24  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: 19 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50485 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1Da\_7.

<400> SEQUENCE: 24  
ggcttacctt cggagctt 19

<210> SEQ ID NO 25  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: 18 nucleotide sequence corresponding to a thermal amplification primer referred to as SQ50484 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1Da\_7.

<400> SEQUENCE: 25  
cgaagatcgc agcgttga 18

<210> SEQ ID NO 26  
<211> LENGTH: 14  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: 14 nucleotide sequence corresponding to a probe referred to as PB50138 used in the zygosity assay for event MON 95379 and hybridizes to the coding sequence of Cry1Da\_7.

<400> SEQUENCE: 26  
acccgggatg acag 14

<210> SEQ ID NO 27  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: 21 nucleotide sequence corresponding to a thermal amplification primer referred to as PNEGDNA used in the zygosity assay for event MON 95379.

<400> SEQUENCE: 27  
ccactgcgcg cgcgccatgg t 21

<210> SEQ ID NO 28  
<211> LENGTH: 14  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: 14 nucleotide sequence corresponding to a probe referred to as PRBNEGDNA used in the zygosity assay for event MON 95379.

<400> SEQUENCE: 28  
cgtcctccga tcca 14

---

What is claimed is:

1. A recombinant DNA molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10 and a complete complement thereof.

2. A corn plant, corn plant part, or corn cell comprising a recombinant polynucleotide molecule comprising SEQ ID NO:9 or SEQ ID NO:10.

3. The corn plant, corn plant part, or corn cell, or part thereof of claim 2, wherein:

a) the corn plant, corn plant part, or corn cell, or part thereof comprises corn event MON 95379; or

b) the corn plant is a progeny of any generation of a corn plant comprising corn event MON 95379; wherein a representative sample of seed of a corn plant comprising corn event MON 95379 has been deposited under ATCC Accession No. PTA-125027.

4. A method for protecting a corn plant from insect infestation, wherein said method comprises providing in the diet of a Lepidopteran insect pest an insecticidally effective amount of cells or tissue of a corn plant comprising a recombinant DNA molecule comprising SEQ ID NO:9 or SEQ ID NO:10.

5. The method of claim 4, wherein said Lepidopteran insect pest is selected from the group consisting of Fall Armyworm (*Spodoptera frugiperda*), Corn Earworm (*Helioverpa zea*), Southwestern Corn Borer (*Diatraea grandiosella*), Surgarcane Borer (*Diatraea saccharalis*), and Lesser Cornstalk Borer (*Elasmopalpus lignosellus*).

6. A method of producing an insect resistant corn plant, said method comprising:

a) sexually crossing two different corn plants, wherein at least one of the two different corn plants comprises SEQ ID NO:9 or SEQ ID NO:10;

b) sampling seed or tissue from progeny of said crossing;

c) detecting the presence of a DNA segment diagnostic for corn event MON 95379 DNA in said sample from step b) to identify progeny comprising corn event MON 95379 DNA; and

d) selecting said progeny comprising corn event MON 95379 DNA;

wherein a representative sample of seed of a corn plant comprising corn event MON 95379 has been deposited under ATCC Accession No. PTA-125027.

7. A corn seed comprising a detectable amount of SEQ ID NO:9 or SEQ ID NO:10.

8. A nonliving corn plant material comprising a detectable amount of SEQ ID NO:9 or SEQ ID NO:10.

9. A microorganism comprising a detectable amount of SEQ ID NO:9 or SEQ ID NO:10.

10. The microorganism of claim 9, wherein the microorganism is a plant cell.

11. A commodity product comprising SEQ ID NO:9 or SEQ ID NO:10.

12. The commodity product of claim 11, further selected from the group consisting of whole or processed corn seed, animal feed comprising corn, corn oil, corn meal, corn flour, corn flakes, corn bran, corn biomass, and fuel products produced using corn and corn parts.

13. The corn plant, corn plant part, or corn cell of claim 2, comprising SEQ ID NO:10.

14. A method of determining the zygosity of a corn plant or corn seed comprising event MON 95379, said method comprising:

a) contacting a sample comprising DNA from said corn plant or corn seed comprising event MON 95379,

wherein a representative sample of seed of a corn plant comprising corn event MON 95379 has been deposited under ATCC Accession No. PTA-125027, with a primer pair that is capable of producing an amplicon of the toxin coding sequence encoding Cry1B.868 or an amplicon of the toxin coding sequence encoding Cry1Da<sub>7</sub>;

b) contacting said sample comprising corn DNA with a primer pair that is capable of producing an amplicon of an internal standard known to be single-copy and homozygous in the corn plant;

c) contacting the DNA sample with a probe set which contains at least a first probe that specifically hybridizes to the toxin coding sequence encoding Cry1B.868 or specifically hybridizes to the toxin coding sequence encoding Cry1 Da<sub>7</sub>, and a second probe that specifically hybridizes to the internal standard genomic DNA known to be single-copy and homozygous in the corn plant;

d) performing a DNA amplification reaction using real-time PCR and determining the cycle thresholds (Ct values) of the amplicon corresponding to the toxin coding sequence and the single-copy, homozygous internal standard;

e) calculating the difference ( $\Delta$ Ct) between the Ct value of the single-copy, homozygous internal standard amplicon and the Ct value of the toxin coding sequence amplicon; and

f) determining zygosity, wherein a  $\Delta$ Ct of zero (0) indicates homozygosity of the inserted T-DNA of event MON 95379 and a  $\Delta$ Ct of about one (1) indicates heterozygosity of the inserted T-DNA of event MON 95379.

15. The method of claim 14, wherein:

a) the primer pairs are selected from the group consisting of SEQ ID NO:18 combined with SEQ ID NO:19, and SEQ ID NO:21 combined with SEQ ID NO:22; and the probes are SEQ ID NO:20 and SEQ ID NO:23; or

b) the primer pairs are selected from the group consisting of SEQ ID NO:18 combined with SEQ ID NO:19, and SEQ ID NO:24 combined with SEQ ID NO:25; and the probes are SEQ ID NO:20 and SEQ ID NO:26.

16. A method of determining the zygosity of a corn plant or corn seed comprising event MON 95379, wherein a representative sample of seed of a corn plant comprising corn event MON 95379 has been deposited under ATCC Accession No. PTA-125027, said method comprising:

a) contacting a sample comprising DNA from said corn plant or corn seed comprising event MON 95379 with a set of primer pairs comprising at least two different primer pairs capable of producing a first amplicon diagnostic for event MON 95379 and a second amplicon diagnostic for native corn genomic DNA not comprising event MON 95379;

b) performing a nucleic acid amplification reaction with the sample and the set of primer pairs; and

c) detecting in the nucleic acid amplification reaction the first amplicon diagnostic for event MON 95379, or the second amplicon diagnostic for native corn genomic DNA not comprising event MON 95379, wherein the presence of only the first amplicon is diagnostic of a corn plant or corn seed homozygous for event MON 95379, and the presence of both the first amplicon and the second amplicon is diagnostic of a corn plant or corn seed heterozygous for event MON 95379;

or said method comprising:

- i) contacting a sample comprising corn DNA comprising event MON 95379 with a probe set which contains at least a first probe that specifically hybridizes to event MON 95379 DNA and at least a second probe that specifically hybridizes to corn genomic DNA that was disrupted by insertion of the heterologous DNA of event MON 95379 and does not hybridize to event MON 95379 DNA; and
- ii) hybridizing the probe set with the sample under stringent hybridization conditions, wherein detecting hybridization of only the first probe under the hybridization conditions is diagnostic for a corn plant or corn seed homozygous for event MON 95379, and wherein detecting hybridization of both the first probe and the second probe under the hybridization conditions is diagnostic for a corn plant or corn seed heterozygous for event MON 95379.

**17.** The method of claim **16**, wherein:

- a) the set of primer pairs comprises SEQ ID NO:15 combined with SEQ ID NO:16, and SEQ ID NO:15 combined with SEQ ID NO:27; or
- b) the probe set comprises SEQ ID NO:17 and SEQ ID NO:28.

\* \* \* \* \*