

(12) **Patent Application Publication**
Ellis et al.

(43) **Pub. Date:** **Aug. 8, 2019**

Publication Classification

(51) **Int. Cl.**

C12N 15/82 (2006.01)

C120 1/6876 (2006.01)

C12Q 1/6874 (2006.01)

C12Q 1/686 (2006.01)

(52) U.S. Cl.

CPC *C12N 15/8274* (2013.01); *C12Q 1/6876*
(2013.01); *C12N 15/8223* (2013.01); *C12Q*
1/686 (2013.01); *C12Q 1/6874* (2013.01)

(57)

ABSTRACT


The invention provides recombinant DNA molecules that are unique to maize event MON87429 and transgenic maize plants, maize plant parts, maize seeds, maize cells, and agricultural products containing maize event MON87429 as well as methods of using and detecting maize event MON87429. Transgenic maize plants containing maize event MON87429 exhibit tolerance to inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS).

Specification includes a Sequence Listing.

(22) Filed: **Jan. 28, 2019**

Related U.S. Application Data

(60) Provisional application No. 62/625,537, filed on Feb. 2, 2018.






Figure 1




Figure 2




Figure 3

MAIZE EVENT MON87429 AND METHODS OF USE THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of priority of U.S. Provisional Application No. 62/625,537, filed Feb. 2, 2018, the disclosure of which is hereby incorporated by reference in its entirety.

INCORPORATION OF SEQUENCE LISTING

[0002] The sequence listing that is contained in the file named "MONS430US-seq.txt", which is 44.2 KB (measured in MS-Windows) and created on Jan. 17, 2019, is filed herewith by electronic submission and incorporated herein by reference.

FIELD OF THE INVENTION

[0003] The invention relates to recombinant DNA molecules of maize event MON87429. The invention also relates to transgenic maize plants, parts, seeds, cells, and agricultural products containing the maize event MON87429 as well as methods of using transgenic maize plants, parts, seeds, cells, and agricultural products containing the maize event MON87429 and detecting maize event MON87429. Transgenic maize plants, parts, seeds, and cells containing maize event MON87429 exhibit tolerance to inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group such as quizalofop and haloxyfop; synthetic auxins such as dicamba and 2,4-D; inhibitors of glutamine synthetase such as glufosinate; and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) such as glyphosate.

BACKGROUND OF THE INVENTION

[0004] Maize (*Zea mays*) is an important crop in many areas of the world, and the use of herbicides for weed control in crop production is a well-established tool. The methods of biotechnology have been used to produce transgenic maize that are tolerant to a specific herbicide due to the expression of a heterologous gene, also known as a transgene. An herbicide tolerance trait can be used alone or combined with other traits, such as tolerance to another herbicide or resistance to pests or pathogens. Combinations of herbicide tolerance traits are desirable to provide weed control options that increase grower flexibility and enable the use of multiple herbicide mode of actions for controlling challenging weeds. A combination of traits can be achieved by breeding together each individual trait. Breeding together individual traits in maize and maintain this combination during breeding with a diverse pool of elite germplasm is a time-consuming and expensive process. A combination of traits can also be achieved by combining multiple traits at one location, or locus, in the genome, thereby simplifying the breeding process. One way to achieve this in through a single transgenic insertion containing multiple transgenes. The combination of multiple herbicide tolerance traits at a single locus in maize would provide a useful tool in weed control that is much simpler and less expensive to maintain during breeding with a diverse pool of elite germplasm.

[0005] The expression of a transgene in a transgenic plant, part, seed, or cell, and therefore its effectiveness, may be influenced by many different factors, such as the elements

used in the transgene's expression cassette and the interaction of those elements. This is complicated further for a transgenic insertion containing two or more expression cassettes with each expression cassette having a transgene conferring a separate trait, also known as a multi-gene transgenic event. A commercially useful multi-gene transgenic event requires that each of the transgenes in the transgenic insertion express in the manner necessary for each trait. To achieve this, individual expression cassettes first are designed and tested in plants, and the best expression cassettes are selected for each trait. Next, the selected expression cassettes for one trait are combined with the selected expression cassettes for the other trait(s) into one construct, and the construct is tested to ensure that all the expression cassettes function well together and each transgene is properly expressed. Then, the selected combination of expression cassettes is used as a single transgenic insert to produce hundreds of multi-gene transgenic events, each event the result of a random insertion of the foreign DNA in a different genomic location.

[0006] Each transgenic event is unique. The unique events are then analyzed through multiple generations of plants to select a superior event for commercial use. The performance of an event in a transgenic plant, part, seed, or cell, and therefore its effectiveness, may be influenced by the genomic location of the transgenic insertion. Events can have the same transgenic insertion and nonetheless have different transgene expression levels and performance across tissues and developmental stages, in various germplasm, or under specific growth conditions. The creation of a multi-gene event for commercial use requires rigorous molecular characterization, greenhouse testing, and field trials over multiple years, in multiple locations, and under a variety of conditions so extensive agronomic, phenotypic, and molecular data may be collected. The resulting data must then be analyzed by scientists and agronomists to select an event that is useful for commercial purposes. The commercial multi-gene event can then be introgressed as a single locus having multiple herbicide tolerance traits into new germplasm using plant breeding methods.

BRIEF SUMMARY OF THE INVENTION

[0007] The invention provides recombinant DNA molecules comprising a sequence selected from the group consisting of SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:8, SEQ ID NO:7, SEQ ID NO:6, SEQ ID NO:5, SEQ ID NO:4, SEQ ID NO:3, SEQ ID NO:2, and SEQ ID NO:1. In one embodiment, the recombinant DNA molecule is derived from a plant, seed, or cell comprising maize event MON87429, a representative sample of seed comprising the event having been deposited as ATCC Accession No. PTA-124635. In another embodiment, the recombinant DNA molecule is in a plant, cell, seed, or plant part comprising maize event MON87429, a representative sample of seed comprising the event having been deposited as ATCC PTA-124635. In another embodiment, the recombinant DNA molecule is an amplicon diagnostic for the presence of maize event MON87429.

[0008] The invention provides a DNA construct comprising four expression cassettes, wherein the first expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Erianthus ravennae*, (II) a phosphinothricin N-acetyltransferase coding sequence, and (III) a fructose-bisphosphate aldolase 3' UTR from *Setaria*

italica; the second expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Coix lacryma-jobi*, (II) an albino and pale green 6 chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (III) a dicamba monooxygenase coding sequence, and (IV) a metallothionein-like protein 3' UTR from *Oryza sativa*; the third expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Arundo donax*, (II) a malate dehydrogenase chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (III) a FT_T protein coding sequence, and (IV) a no apical meristem protein 3' UTR from *Oryza sativa*; and the fourth expression cassette comprises in operable linkage (I) a CaMV 35S promoter and leader, (II) a chlorophyll a/b-binding protein leader from *Triticum aestivum*, (III) an actin 1 intron from *Oryza sativa*, (IV) a ShkG chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (V) a glyphosate tolerant 5-enolpyruvylshikimate-3-phosphate synthase coding sequence from *Agrobacterium* sp strain CP4, (VI) a male tissue specific siRNA target from *Zea mays*, and (VII) a glycine-rich RNA binding protein 3'UTR from *Oryza sativa*.

[0009] The invention provides a DNA molecule having a sufficient length of contiguous nucleotides of SEQ ID NO:10 to function as a DNA probe specific for SEQ ID NO:10 in a sample of DNA derived from a maize plant, maize seed, or maize cell. In one embodiment, the DNA probe comprises SEQ ID NO:13.

[0010] The invention provides a pair of DNA molecules comprising a first DNA molecule and a second DNA molecule, wherein the first and second DNA molecules each comprise a fragment SEQ ID NO:10 and function as DNA primers when used together in an amplification reaction with DNA containing maize event MON87429 to produce an amplicon diagnostic for maize event MON87429 in a sample. In one embodiment, at least one DNA primer comprises a fragment of a sequence selected from the group consisting of SEQ ID NO:7 and SEQ ID NO:8. In another embodiment, the first DNA molecule comprises SEQ ID NO:11 and the second DNA molecule comprises SEQ ID NO:12.

[0011] The invention provides a method of detecting the presence of maize event MON87429 in a sample of DNA derived from a maize plant, maize seed, or maize cell, the method comprising: contacting the sample with a DNA probe; subjecting the sample and the DNA probe to stringent hybridization conditions; and detecting hybridization of the DNA probe to a DNA molecule in the sample, wherein the hybridization of the DNA probe to the DNA molecule indicates the presence of maize event MON87429 in the sample of DNA.

[0012] The invention provides a method of detecting the presence of maize event MON87429 in a sample of DNA derived from a maize plant, maize seed, or maize cell, the method comprising: contacting the sample with a pair of DNA molecules that function as DNA primers; performing an amplification reaction sufficient to produce a DNA amplicon comprising a sequence selected from the group consisting of SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:8, SEQ ID NO:7, SEQ ID NO:6, SEQ ID NO:5, SEQ ID NO:4, SEQ ID NO:3, SEQ ID NO:2, and SEQ ID NO:1; and detecting the presence of the DNA amplicon, wherein the presence of the DNA amplicon indicates the presence of maize event MON87429 in the sample.

[0013] The invention provides a method of detecting the presence of maize event MON87429 in a sample derived from a maize plant, maize seed, or maize cell, the method comprising: contacting the sample with at least one antibody specific for at least one or more protein encoded by maize event MON87429; and detecting binding of the antibody to the protein in the sample, wherein the binding of the antibody to the protein indicates the presence of maize event MON87429 in the sample. In another embodiment, the method comprises additionally contacting the sample with at least a second antibody specific for a second protein encoded by maize event MON87429. In another embodiment, the method comprises additionally contacting the sample with at least a second antibody and a third antibody specific for a second protein and a third protein, respectively, encoded by maize event MON87429. In another embodiment, the method comprises additionally contacting the sample with at least a second antibody, a third antibody, and a fourth antibody specific for a second protein, a third protein, and a fourth protein, respectively, encoded by maize event MON87429.

[0014] The invention provides a kit for detecting the presence of maize event MON87429 comprising a DNA probe specific for SEQ ID NO:10, a pair of DNA primers that produce an amplicon diagnostic for maize event MON87429, or at least one antibody specific for at least one protein encoded by maize event MON87429.

[0015] The invention provides a plant, seed, cell, plant part, or commodity product comprising a DNA molecule comprising a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10. In one embodiment, the plant, seed, cell, or plant part is tolerant to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof. In another embodiment, the plant, seed, cell, or plant part is tolerant to quizalofop, haloxyfop, dicamba, 2,4-D, and glufosinate.

[0016] The invention provides a method for controlling weeds in a crop-growing area comprising planting maize comprising maize event MON87429 in the crop-growing area and applying to the crop-growing area, or any portion thereof, an effective amount of at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, to control the weeds in the area without injuring the maize. In one embodiment, the method comprises applying at least two or more herbicides selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) over a growing season. In another embodiment, the method comprises applying an herbicide selected from the group consisting of quizalofop, haloxyfop, dicamba, 2,4-D, glufosinate, and glyphosate, or any combination thereof. In one embodiment, the effective amount of dicamba is about 0.1 lb

ae/acre to about 16 lb ae/acre over a growing season. In one embodiment, the effective amount of dicamba is about 0.5 lb ae/acre to about 2 lb ae/acre over a growing season. In one embodiment, the effective amount of glufosinate is about 0.1 lb ae/acre to about 16 lb ae/acre over a growing season. In one embodiment, the effective amount of glufosinate is about 0.4 lb ae/acre to about 1.59 lb ae/acre over a growing season. In one embodiment, the effective amount of 2,4-D is about 0.1 lb ae/acre to about 10 lb ae/acre over a growing season. In one embodiment, the effective amount of 2,4-D is about 0.75 lb ae/acre to 1.0 lb ae/acre over a growing season. In one embodiment, the effective amount of the FOP herbicide is about 0.01 lb ai/acre to about 1.0 lb ai/acre over a growing season. In one embodiment, the effective amount of the FOP herbicide is about 0.034 lb ai/acre to about 0.083 lb ai/acre of quizalofop over a growing season. In one embodiment, the effective amount of the FOP herbicide is about 0.018 ai/acre to about 0.07 lb ai/acre of haloxyfop over a growing season.

[0017] The invention provides a method for controlling volunteer maize comprising maize event MON87429 in an area comprising applying an herbicidally effective amount of at least one cyclohexanedione (DIM) herbicide, where the herbicide application prevents growth of maize comprising maize event MON87429. In one embodiment, the cyclohexanedione (DIM) herbicide is selected from the group consisting of clethodim, sethoxydim, and tralkoxydim.

[0018] The invention provides a method of producing a plant that is tolerant to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, the method comprising: breeding a plant comprising maize event MON87429 with itself or a second plant to produce seed; and identifying progeny seed that comprise maize event MON87429. In one embodiment, identifying progeny seed that comprise maize event MON87429 is by growing the progeny seed to produce progeny plants; treating the progeny plants with an effective amount of at least one herbicide selected from the group consisting of quizalofop, haloxyfop, dicamba, 2,4-D, glufosinate, glyphosate, or any combination thereof; and selecting a progeny plant that is tolerant to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS). In one embodiment, identifying progeny seed that comprise maize event MON87429 is by detecting the presence of maize event MON87429 in a sample derived from the progeny seed. In one embodiment, identifying progeny seed that comprise maize event MON87429 is by detecting the presence of at least one protein encoded by maize event MON87429 in a sample derived from the progeny seed.

[0019] The invention provides a method of producing hybrid seed comprising: growing a plant comprising SEQ ID NO:10; applying an effective amount of glyphosate prior to or during the development of the male reproductive tissue of the plant thereby inducing male-sterility in the plant; fertilizing the plant with pollen from a second plant; and harvesting hybrid seed from the plant. In one embodiment, the glyphosate is applied prior to or during the development at

an effective amount of about 0.25 lb ae/acre to about 11.0 lb ae/acre. In one embodiment, the glyphosate is applied prior to or during the development at an effective amount of about 0.5 lb ae/acre to about 2.5 lb ae/acre total in one or more applications. In one embodiment, the effective amount of glyphosate is applied at a developmental stage selected from the group consisting of the V4, V5, V6, V7, V8, V9, V10, V11, V12, V13, and V14 stage of maize plant development. The invention provides hybrid seed comprising SEQ ID NO:10 and produced by using the method of producing hybrid seed comprising: growing a plant comprising SEQ ID NO:10; applying an effective amount of glyphosate prior to or during the development of the male reproductive tissue of the plant thereby inducing male-sterility in the plant; fertilizing the plant with pollen from a second plant; and harvesting hybrid seed from the plant.

[0020] The invention provides a method of determining zygosity of a plant for maize event MON87429 comprising: contacting a sample comprising DNA derived from the plant with a primer set capable of producing a first amplicon diagnostic for the presence of maize event MON87429 and a second amplicon diagnostic for the wild-type maize genomic DNA not comprising maize event MON87429; performing a nucleic acid amplification reaction; detecting the first amplicon and the second amplicon, wherein the presence of both amplicons indicates the sample is heterozygous for maize event MON87429 and the presence of only the first amplicon indicates the sample is homozygous for maize event MON87429. In one embodiment, the primer set comprises SEQ ID NO:11 and SEQ ID NO:12.

[0021] The invention provides a method of improving tolerance to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, in a maize plant comprising: (a) obtaining a DNA construct comprising four expression cassettes, wherein the first expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Erianthus ravennae*, (II) a phosphinothricin N-acetyltransferase coding sequence, and (III) a fructose-bisphosphate aldolase 3' UTR from *Setaria italica*; the second expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Coix lacrym-jobi*, (II) an albino and pale green 6 chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (III) a dicamba monooxygenase coding sequence, and (IV) a metallothionein-like protein 3' UTR from *Oryza sativa*; the third expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Arundo donax*, (II) a malate dehydrogenase chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (III) a FT_T protein coding sequence, and (IV) a no apical meristem protein 3' UTR from *Oryza sativa*; and the fourth expression cassette comprises in operable linkage (I) a CaMV 35S promoter and leader, (II) a chlorophyll a/b-binding protein leader from *Triticum aestivum*, (III) an actin 1 intron from *Oryza sativa*, (IV) a ShkG chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (V) a glyphosate tolerant 5-enolpyruvylshikimate-3-phosphate synthase coding sequence from *Agrobacterium* sp strain CP4, (VI) a male tissue specific siRNA target from *Zea mays*, and (VII) a glycine-rich RNA binding protein 3'UTR from *Oryza sativa*;

(b) inserting the DNA construct into the genome of a maize cell; (c) regenerating the maize cell into a maize plant; and (d) selecting a maize plant comprising the DNA construct. In one embodiment, selecting is by treating the maize plant with an effective amount of at least one herbicide selected from the group consisting of quizalofop, haloxyfop, dicamba, 2,4-D, glufosinate, or glyphosate. In one embodiment, the invention provides a maize plant, maize seed, or maize cell tolerant to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, and obtainable by the method, wherein the maize plant, maize seed, or maize cell comprises the DNA construct. In a further embodiment, the maize plant, maize seed, or maize cell produced by the method comprises SEQ ID NO:10.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] FIG. 1 represents the sequence of maize event MON87429. Horizontal lines correspond to the positions of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9 relative to SEQ ID NO:10; the horizontal arrows labeled SQ51062 (SEQ ID NO:11) and SQ51053 (SEQ ID NO:12) represent the approximate position of a pair of primers that can be used to detect maize event MON87429; and the horizontal line labeled PB50370 (SEQ ID NO:13) represents the approximate position of a DNA probe that can be used to detect maize event MON87429.

[0023] FIG. 2 represents the four expression cassettes of maize event MON87429 relative to SEQ ID NO:9 with their respective genetic elements labeled as described in Table 1.

[0024] FIG. 3 represents the creation, testing, characterization, and selection of the MON87429 event as described herein. All times are approximate.

BRIEF DESCRIPTION OF THE SEQUENCES

[0025] SEQ ID NO:1 is a thirty nucleotide DNA sequence representing the 5' junction of maize genomic DNA and the transgene insert. SEQ ID NO:1 corresponds to nucleotide positions 1015 to 1044 of SEQ ID NO:10.

[0026] SEQ ID NO:2 is a thirty nucleotide DNA sequence representing the 3' junction of maize genomic DNA and the transgene insert. SEQ ID NO:2 corresponds to nucleotide positions 15023 to 15052 of SEQ ID NO:10.

[0027] SEQ ID NO:3 is a sixty nucleotide DNA sequence representing the 5' junction of maize genomic DNA and the transgene insert. SEQ ID NO:3 corresponds to nucleotide positions 1000 to 1059 of SEQ ID NO:10.

[0028] SEQ ID NO:4 is a sixty nucleotide DNA sequence representing the 3' junction of maize genomic DNA and the transgene insert. SEQ ID NO:4 corresponds to nucleotide positions 15008 to 15067 of SEQ ID NO:10.

[0029] SEQ ID NO:5 is a one-hundred nucleotide DNA sequence representing the 5' junction of maize genomic DNA and the transgene insert. SEQ ID NO:5 corresponds to nucleotide positions 980 to 1079 of SEQ ID NO:10.

[0030] SEQ ID NO:6 is a one-hundred nucleotide DNA sequence representing the 3' junction of maize genomic DNA and the transgene insert. SEQ ID NO:6 corresponds to nucleotide positions 14988 to 15087 of SEQ ID NO:10.

[0031] SEQ ID NO:7 is a 1350 nucleotide DNA sequence representing 1029 nucleotides of the 5' flanking maize genomic DNA and 321 nucleotides of the 5' end of the transgene insert.

[0032] SEQ ID NO:8 is a 1069 nucleotide DNA sequence representing 38 nucleotides of the 3' end of the transgene insert and 1031 nucleotides of the 3' flanking maize genomic DNA.

[0033] SEQ ID NO:9 is a 14008 nucleotide DNA sequence corresponding to the transgene insert of the maize MON87429 event.

[0034] SEQ ID NO:10 is a 16068 nucleotide DNA sequence corresponding to the maize MON87429 event; the sequence contains the 5' flanking genomic DNA sequence from positions 1 to 1029, the transgenic DNA insert from positions 1030 to 15037, and the 3' flanking genomic DNA sequence from positions 15038 to 16068.

[0035] SEQ ID NO:11 is a 29 nucleotide DNA sequence corresponding to a primer referred to as SQ51062 and used to identify maize MON87429 event DNA in a sample; it corresponds to positions 15038 to 15066 of SEQ ID NO:10.

[0036] SEQ ID NO:12 is a 17 nucleotide DNA sequence corresponding to a primer referred to as SQ51053 and used to identify maize MON87429 event DNA in a sample; it corresponds to positions 14987 to 15003 of SEQ ID NO:10.

[0037] SEQ ID NO:13 is a 16 nucleotide DNA sequence corresponding to a probe referred to as PB50370 and used to identify maize MON87429 event DNA in a sample; it corresponds to positions 15009 to 15024 of SEQ ID NO:10.

DETAILED DESCRIPTION OF THE INVENTION

[0038] The following definitions and methods are provided to better define the invention and to guide those of ordinary skill in the art in the practice of the invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

[0039] Plant transformation techniques are used to insert foreign DNA (also known as transgenic DNA) randomly into a chromosome of the genome of a cell to produce a genetically engineered cell, also referred to as "transgenic" or "recombinant" cell. Using this technique, many individual cells are transformed, each resulting in a unique transgenic event due to the random insertion of the foreign DNA into the genome. A transgenic plant is then regenerated from each individual transgenic cell. This results in every cell of the transgenic plant containing the uniquely inserted transgenic event as a stable part of its genome. This transgenic plant can then be used to produce progeny plants, each containing the unique transgenic event. Maize event MON87429 was produced by: (i) transformation of thousands of maize cells with a DNA construct that includes four expression cassettes (each expression cassette having been selected after individual testing followed by testing in combination with the other three expression cassettes), (ii) regeneration of a population of transgenic plants each containing a unique transgenic event, and (iii) rigorous multi-year event selection involving the testing and analysis of molecular characteristics, herbicide tolerance efficacy, and agronomic properties in a variety of genetic backgrounds for thousands of events through tens of thousands of plants.

Maize event MON87429 was thus produced and selected as a uniquely superior event useful for broad-scale agronomic commercial purposes.

[0040] As used herein, a “transgenic event” or an “event” is a DNA molecule created by the act of inserting a transgenic DNA molecule into the genomic DNA of a plant cell using plant transformation methods known in the art. This insertion creates a new, transgenic genomic DNA sequence that consists of the inserted foreign DNA (referred to as the “transgenic insert”) and the genomic DNA immediately adjacent to, or “flanking”, the transgenic insert on either side of the insertion location (referred to as the “flanking DNA”). The DNA sequence of an event is unique to and specific for the event and can be readily identified when compared to other DNA sequences, such as that of other events or untransformed maize genomic DNA. Maize event MON87429 has the new and unique DNA sequence provide as SEQ ID NO:10, which contains the transgenic insert sequence provided as SEQ ID NO:9 and the 5' and 3' flanking DNA sequence provided in SEQ ID NO:7 and SEQ ID NO:8, respectively. Maize event MON87429 is thus a DNA molecule that is an integral part of the chromosome of transgenic maize cells and plants comprising the event and as such is static and may be passed on to progeny cells and plants.

[0041] The present invention also provides progeny of the original transformed cell and plant that comprise maize

event MON87429. Such progeny may be produced by cell tissue culture, by selfing of a maize plant comprising the maize event MON87429, or by sexual outcrossing between a maize plant comprising maize event MON87429 and another plant that does or does not contain the event. Such other plant may be a transgenic plant comprising the same or different event(s) or a nontransgenic plant, such as one from a different variety. Maize event MON87429 is passed from the original parent through each generation to the progeny.

[0042] As used herein, the term “maize” means *Zea mays* (also referred to as corn) and includes all plant varieties that can be bred with *Zea mays*.

[0043] The invention provides maize event MON87429, which provides to maize cells, plants, and seeds that comprise the event tolerance to inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group such as quizalofop and haloxyfop; synthetic auxins such as dicamba and 2,4-D; inhibitors of glutamine synthetase such as glufosinate; and the 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitor glyphosate. Maize event MON87429 contains four expression cassettes. As used herein, an “expression cassette” or “cassette” is a recombinant DNA molecule comprising a combination of distinct elements that are to be expressed by a transformed cell. Table 1 provides a list of the elements contained in SEQ ID NO:10 and as illustrated in FIG. 2.

TABLE 1

Description of maize event MON87429		
Element	Position in SEQ ID NO: 10	Description
5' Flanking DNA	1-1029	DNA sequence flanking the 5' end of the transgenic insert
Left Border Region	1030-1288	DNA region from <i>Agrobacterium tumefaciens</i> containing the left border sequence
Intervening Sequence	1289-1359	Sequence used in DNA cloning
P-Ea.ubq	1360-3541	Promoter, 5' UTR, and intron sequences of a ubiquitin gene from <i>Erianthus ravennae</i>
Intervening Sequence	3542-3546	Sequence used in DNA cloning
CS-Sv.pat	3547-4098	Coding sequence for the phosphinothricin N-acetyltransferase (PAT) protein
T-Si.fba	4099-4475	3' UTR sequence of the fructose-bisphosphate aldolase gene from <i>Setaria italica</i>
Intervening Sequence	4476-4537	Sequence used in DNA cloning
P-Clj.ubq	4538-6463	Promoter, 5' UTR, and intron sequences of a ubiquitin gene from <i>Coix lacryma-jobi</i>
Intervening Sequence	6464-6473	Sequence used in DNA cloning
TS-At.apg6	6474-6677	Codon optimized targeting sequence of the Albino and pale green 6 gene from <i>Arabidopsis thaliana</i>
CS-Sm.dmo	6678-7700	Coding sequence for the dicamba monooxygenase (DMO) protein
Intervening Sequence	7701-7708	Sequence used in DNA cloning
T-Os.mt	7709-8008	3' UTR sequence of the metallothionein-like protein from <i>Oryza sativa</i>
Intervening Sequence	8009-8016	Sequence used in DNA cloning
P-Ad.ubq	8017-9973	Promoter, 5' UTR, and intron sequences of a ubiquitin gene from <i>Arundo donax</i>
Intervening Sequence	9974-9986	Sequence used in DNA cloning
TS-At.mdh	9987-10229	Sequence of the transit peptide of the malate dehydrogenase gene from <i>Arabidopsis thaliana</i>

TABLE 1-continued

Description of maize event MON87429		
Element	Position in SEQ ID NO: 10	Description
CS-Sh.ft_t	10230-11117	Coding sequence for the FT_T protein
Intervening Sequence	11118-11132	Sequence used in DNA cloning
T-Os.nam	11133-11649	3' UTR sequence of the no apical meristem protein from <i>Oryza sativa</i>
Intervening Sequence	11650-11655	Sequence used in DNA cloning
P-CaMV.35S	11656-11979	Promoter and leader from the 35S RNA of cauliflower mosaic virus
Intervening Sequence	11980-12001	Sequence used in DNA cloning
L-Ta.cab	12002-12062	5' UTR leader sequence from chlorophyll a/b-binding protein of <i>Triticum aestivum</i>
Intervening Sequence	12063-12078	Sequence used in DNA cloning
I-Os.ract1	12079-12558	Intron and UTR sequence of the Actin 1 protein from <i>Oryza sativa</i>
Intervening Sequence	12559-12567	Sequence used in DNA cloning
TS-At.CTP2	12568-12795	Transit peptide sequence of the ShkG gene from <i>Arabidopsis thaliana</i>
CS-cp4epsps	12796-14163	Coding sequence for the 5-enolpyruvylshikimate-3- phosphate synthase (CP4-EPSPS) protein
Intervening Sequence	14164-14169	Sequence used in DNA cloning
mts-siRNA	14170-14370	Sequence of the male tissue specific siRNA target
Intervening Sequence	14371-14378	Sequence used in DNA cloning
T-Os.grp3	14379-14989	3' UTR sequence of the glycine-rich RNA-binding protein (Grp3) gene from <i>Oryza sativa</i>
Intervening Sequence	14990-15030	Sequence used in DNA cloning
Right Border Region	15031-15037	DNA region from <i>Agrobacterium tumefaciens</i> containing the right border sequence
3' Flanking DNA	15038-16068	Flanking DNA

[0044] As used herein, the term “recombinant” refers to a non-natural DNA, protein, or organism that would not normally be found in nature and was created by human intervention. As used herein, a “recombinant DNA molecule” is a DNA molecule comprising a combination of DNA molecules that would not naturally occur together and is the result of human intervention, for example, a DNA molecule that is comprised of a combination of at least two DNA molecules heterologous to each other, such as a DNA molecule that comprises a transgene and the plant genomic DNA adjacent to the transgene. An example of a recombinant DNA molecule is a DNA molecule comprising at least one sequence selected from SEQ ID NO:1-10. As used herein, a “recombinant plant” is a plant that would not normally exist in nature, is the result of human intervention, and contains a transgenic DNA molecule. As a result of such genomic alteration, the recombinant plant is something new and distinctly different from the related wild-type plant. An example of a recombinant plant is a maize plant containing the maize event MON87429.

[0045] As used herein, the term “transgene” refers to a DNA molecule artificially incorporated into an organism’s genome as a result of human intervention, such as by plant transformation methods. A transgene may be heterologous to the organism. The term “transgenic insert” as used herein refers to the foreign DNA inserted by plant transformation techniques into the maize genome to produce maize event

MON87429. The sequence for the transgenic insert of maize event MON87429 is provided as SEQ ID NO:9. The term “transgenic” refers to comprising a transgene, for example a “transgenic plant” refers to a plant comprising a transgene.

[0046] As used herein, the term “heterologous” refers to a first molecule not normally associated with a second molecule or an organism in nature. For example, a DNA molecule may be from a first species and inserted into the genome of a second species. The DNA molecule would thus be heterologous to the genome and the organism.

[0047] As used herein, the term “chimeric” refers to a single DNA molecule produced by fusing a first DNA molecule to a second DNA molecule, where neither first nor second DNA molecule would normally be found in that configuration fused to the other. The chimeric DNA molecule is thus a new DNA molecule not normally found in nature. An example of a chimeric DNA molecule is a DNA molecule comprising at least one sequence selected from SEQ ID NO:1-10.

[0048] As used herein, the term “isolated” refers to separating a molecule from other molecules that are normally associated with it in its native or natural state. The term “isolated” thus may refer to a DNA molecule that has been separated from other DNA molecule(s) that it is associated with it in its native or natural state. Such a DNA molecule may be present in a recombined state, such as a recombinant DNA molecule. Thus, a DNA molecule removed from its

natural state and fused to another DNA molecule with which it is not normally associated would be an isolated DNA molecule. Such an isolated DNA molecule could result from the use of biotechnology techniques, such as making recombinant DNA or integrating a foreign DNA molecule into the chromosome of a cell, plant, or seed.

[0049] The invention provides DNA molecules and their corresponding DNA sequences. As used herein, the terms “DNA” and “DNA molecule” refer to a deoxyribonucleic acid (DNA) molecule. A DNA molecule may be of genomic or synthetic origin and is by convention from the 5' (upstream) end to the 3' (downstream) end. As used herein, the term “DNA sequence” refers to the nucleotide sequence of a DNA molecule. The nomenclature used is that required by Title 37 of the United States Code of Federal Regulations § 1.822 and set forth in the tables in WIPO Standard ST.25 (1998), Appendix 2, Tables 1 and 3. By convention, the DNA sequences of the invention and fragments thereof are disclosed with reference to only one strand of the two complementary DNA sequence strands. By implication and intent, the complementary sequences of the sequences provided here (the sequences of the complementary strand), also referred to in the art as the reverse complementary sequences, are within the scope of the invention and are expressly intended to be within the scope of the subject matter claimed. Thus, as used herein references to SEQ ID NO:1-10 and fragments thereof include and refer to the sequence of the complementary strand and fragments thereof.

[0050] As used herein, the term “fragment” refers to a smaller piece of a whole. For example, fragments of SEQ ID NO:10 would include sequences that are at least about 10 consecutive nucleotides, at least about 11 consecutive nucleotides, at least about 12 consecutive nucleotides, at least about 13 consecutive nucleotides, at least about 14 consecutive nucleotides, at least about 15 consecutive nucleotides, at least about 16 consecutive nucleotides, at least about 17 consecutive nucleotides, at least about 18 consecutive nucleotides, at least about 19 consecutive nucleotides, at least about 20 consecutive nucleotides, at least about 25 consecutive nucleotides, at least about 30 consecutive nucleotides, at least about 35 consecutive nucleotides, at least about 40 consecutive nucleotides, at least about 45 consecutive nucleotides, at least about 50 consecutive nucleotides, at least about 60 consecutive nucleotides, at least about 70 consecutive nucleotides, at least about 80 consecutive nucleotides, at least about 90 consecutive nucleotides, or at least about 100 consecutive nucleotides of the complete sequence of SEQ ID NO:10.

[0051] The DNA sequence for the transgenic insert of maize event MON87429 is provided as SEQ ID NO:9. The DNA sequence of the transgenic insert and the maize genomic DNA flanking each side of the transgenic insert is provided as SEQ ID NO:10. The DNA sequences of a portion of flanking DNA and the 5' end of the transgenic insert are provided as SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, and SEQ ID NO:7. The DNA sequences of a portion of flanking DNA and the 3' end of the transgenic insert are provided as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, and SEQ ID NO:8.

[0052] The DNA sequence of the region spanning the connection by phosphodiester bond linkage of one end of the transgenic insert to the flanking maize genomic DNA is referred to herein as a “junction”. A junction is the connec-

tion point of the transgenic insert and flanking DNA as one contiguous molecule. One junction is found at the 5' end of the transgenic insert and the other is found at the 3' end of the transgenic insert, referred to herein as the 5' and 3' junction, respectively. A “junction sequence” refers to a DNA sequence of any length that spans the 5' or 3' junction of an event. Junction sequences of maize event MON87429 are apparent to one of skill in the art using SEQ ID NO:10. Examples of junction sequences of maize event MON87429 are provided as SEQ ID NO:1-8. FIG. 1 illustrates the physical arrangement of SEQ ID NO:1-10 arranged from 5' to 3'. The junction sequences of maize event MON87429 may be present as part of the genome of a plant, seed, or cell containing maize event MON87429. The identification of any one or more of SEQ ID NO:1-8 or 10 in a sample from a plant, plant part, seed, or cell indicates that the DNA was obtained from maize containing maize event MON87429 and is diagnostic for the presence of maize event MON87429.

[0053] The plants, seeds, cells, plant parts, and commodity products of the invention may be used for detection of DNA or protein molecules indicative of the presence of maize event MON87429. Provided are exemplary DNA molecules that can be used either as primers or probes for detecting the presence of maize event MON87429 in a sample. Such primers or probes are specific for a target nucleic acid sequence and as such are useful for the identification of maize event MON87429 by the methods described here. Detection of the presence of maize event MON87429 may be done by using methods known in the art, such as thermal amplification of nucleic acid or nucleic acid hybridization techniques (such as northern blotting and southern analysis).

[0054] A “primer” is a DNA molecule that is designed for use in annealing or hybridization methods that involve an amplification reaction. An amplification reaction is an in vitro reaction that amplifies template DNA to produce an amplicon. As used herein, an “amplicon” is a DNA molecule that has been synthesized using amplification techniques. Amplicons of the invention have a DNA sequence comprising one or more of SEQ ID NO:1-10, or fragments thereof. A pair of primers may be used with template DNA, such as a sample of maize genomic DNA, in an amplification reaction, such as polymerase chain reaction (PCR), to produce an amplicon, where the amplicon produced would have a DNA sequence corresponding to sequence of the template DNA located between the two sites where the primers hybridized to the template. A primer is typically designed to hybridize to a complementary target DNA strand to form a hybrid between the primer and the target DNA strand. The presence of a primer is a point of recognition by a polymerase to begin extension of the primer using as a template the target DNA strand. Primer pairs refer to use of two primers binding opposite strands of a double stranded nucleotide segment for the purpose of amplifying the nucleotide segment between them. Examples of primer sequences are provided as SEQ ID NO:11 (SQ51062) and SEQ ID NO:12 (SQ51053). The primer pair provided as SEQ ID NO:11 and SEQ ID NO:12 are useful as a first DNA molecule and a second DNA molecule, where the first DNA molecule is a fragment of the transgenic insert DNA sequence of SEQ ID NO:10 and the second DNA molecule is a fragment of the flanking DNA sequence of SEQ ID NO:10, and each are of sufficient length to function as DNA primers when used together in an amplification reaction with

DNA containing maize event MON87429 to produce an amplicon diagnostic for maize event MON87429 in a sample. Primer pairs of the present invention may in certain embodiments also be defined as comprising a first and second DNA molecule, wherein the first DNA molecule is a fragment of the maize genomic portion of SEQ ID NO:10 and the second DNA molecule is a fragment of the transgene portion of SEQ ID NO:10, and each are of sufficient length to function as DNA primers when used together in an amplification reaction with DNA containing maize event MON87429 to produce an amplicon diagnostic for maize event MON87429 in a sample.

[0055] A “probe” is a nucleic acid molecule that is complementary to a strand of a target nucleic acid and useful in hybridization detection methods. Probes according to the invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and the detection of such binding can be useful in detecting the presence or absence of the target DNA sequence. A probe may be attached to a conventional detectable label or reporter molecule, such as a radioactive isotope, ligand, chemiluminescent agent, or enzyme. An exemplary DNA sequence useful as a probe for detecting maize event MON87429 is provided as SEQ ID NO:13 (PB50370).

[0056] Methods for designing and using primers and probes are well known in the art. DNA molecules comprising the full length of or fragments of SEQ ID NO:1-10 are useful as primers and probes for detecting maize event MON87429 and can readily be designed by one of skill in the art using the sequences provided herein.

[0057] Probes and primers according to the invention may have complete sequence identity with the target sequence, although primers and probes differing from the target sequence that retain the ability to hybridize preferentially to target sequences may be designed by conventional methods. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of transgenic DNA from maize event MON87429 in a sample. Probes and primers are generally at least about 11 nucleotides, at least about 18 nucleotides, at least about 24 nucleotides, or at least about 30 nucleotides or more in length. Such probes and primers hybridize specifically to a target DNA sequence under stringent hybridization conditions. Conventional stringency conditions are described by M R Green and J Sambrook, *Molecular cloning: a laboratory manual*, 4th Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2012). As used herein, two nucleic acid molecules are capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is the “complement” of another nucleic acid molecule if they exhibit complete complementarity. As used herein, two molecules exhibit “complete complementarity” if when aligned every nucleotide of the first molecule is complementary to every nucleotide of the second molecule. Two molecules are “minimally complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional “low-stringency” conditions.

Similarly, the molecules are “complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional “high-stringency” conditions. Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure.

[0058] Appropriate stringency conditions that promote DNA hybridization, for example, 6.0x sodium chloride/sodium citrate (SSC) at about 45° C., followed by a wash of 2.0xSSC at 50° C., are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0xSSC at 50° C. to a high stringency of about 0.2xSSC at 50° C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22° C., to high stringency conditions at about 65° C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed.

[0059] Provided are proteins that can be used to produce antibodies for detecting the presence of maize event MON87429 in a sample. Such antibodies are specific for one or more of the proteins that are encoded by maize event MON87429. The DNA sequence encoding such proteins is provided in SEQ ID NO:10 and the start positions and stop positions of the coding sequence are indicated in Table 1. The DNA sequence encoding each protein and the protein encoded by the sequence are useful to produce antibodies for detecting the presence of maize event MON87429 by the methods described here. Detection of the presence of maize event MON87429 may be done by using any protein detection techniques known in the art, such as western blotting, immuno-precipitation, enzyme-linked immunosorbent assay (ELISA), antibody attachment to a detectable label or reporter molecule (such as a radioactive isotope, ligand, chemiluminescent agent, or enzyme), or enzymatic action on a reporter molecule. One method provides for contacting a sample with an antibody that binds to the DMO, PAT, FT, T, or CP4-EPSPS protein encoded by maize event MON87429 and then detecting the presence or absence of antibody binding. The binding of such antibody is diagnostic for the presence of one or more proteins encoded by maize event MON87429.

[0060] Protein and nucleic acid detection kits for detecting the presence of maize event MON87429 are provided. Variations on such kits can also be developed using the compositions and methods disclosed herein and the methods well known in the art of protein and nucleic acid detection. Protein and nucleic acid detection kits can be applied to methods for breeding with plants containing maize event MON87429. Such kits contain primers or probes comprising fragments of SEQ ID NO:1-10 or antibodies specific for a protein encoded by maize event MON87429.

[0061] One example of a detection kit comprises at least one DNA molecule of sufficient length of contiguous nucleotides of SEQ ID NO:10 to function as a DNA probe useful for detecting the presence or absence of maize event MON87429 in a sample. An exemplary DNA molecule sufficient for use as a probe is one comprising the sequence provided as SEQ ID NO:13. Other probes may be readily designed by one of skill in the art. Another example of a

detection kit comprises at least one primer pair useful for producing an amplicon useful for detecting the presence or absence of maize event MON87429 in a sample. Such a method may also include sequencing the amplicon or a fragment thereof. Exemplary DNA molecules sufficient for use as a primer pair are ones comprising the sequences provided as SEQ ID NO:11 and SEQ ID NO:12, respectively. Other primer pairs may be readily designed by one of skill in the art. Kits of the invention may optionally also comprise reagents for performing the detection or diagnostic reactions described herein. Another example of a detection kit comprises at least one antibody specific for at least one protein encoded by maize event MON87429. For example, such a kit may utilize a lateral flow strip comprising reagents activated when the tip of the strip is contacted with an aqueous solution. Exemplary proteins sufficient for use in antibody production are ones encoded by the sequence provided as SEQ ID NO:10, or any fragment thereof.

[0062] The invention provides maize plants, progeny, seeds, cells, and plant parts containing maize event MON87429, and commodity products produced using these. The plants, progeny, seeds, cells, plant parts, and commodity products of the invention contain a detectable amount of DNA having at least one of the sequences provided as SEQ ID NO:1-8 and SEQ ID NO:10.

[0063] Plants, progeny, seeds, cells, and plant parts of the invention may also contain one or more additional transgenic traits, particularly those introduced by crossing a maize plant containing maize event MON87429 with another plant containing the additional transgenic trait(s). Such traits include but are not limited to increased insect resistance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, improved nutritional quality, hybrid seed production, and/or increased herbicide tolerance, in which the trait is measured with respect to a maize plant lacking such transgenic trait.

[0064] Plants of the invention may be used to produce progeny that contain maize event MON87429. As used herein, "progeny" includes any plant, seed, and cell comprising maize event MON87429 inherited from an ancestor plant, indicated by the plant comprising a DNA molecule having at least one sequence selected from SEQ ID NO:1-8 and SE ID NO:10. Plants, seeds, and cells may be homozygous or heterozygous for maize event MON87429. Progeny plants may be grown from seeds produced by a maize plant containing maize event MON87429 or from seeds produced by a maize plant fertilized with pollen containing maize event MON87429.

[0065] As used herein, a "plant part" of the invention is any part from a plant containing maize event MON87429. Plant parts include but are not limited to tissue samples, pollen, ovule, pod, flower, roots, stems, fibers, and leaves in whole or part. Plant parts may be viable or nonviable.

[0066] The invention provides a commodity product that is produced from plants containing maize event MON87429. Commodity products of the invention contain a detectable amount of DNA comprising a DNA sequence selected from the group consisting of SEQ ID NO:1-10. As used herein, a "commodity product" refers to any composition or product which is comprised of material from plant, seed, cell, or plant part comprising maize event MON87429. Commodity products include but are not limited to processed seeds, grains, plant parts, and meal. A commodity product of the

invention will contain a detectable amount of DNA corresponding to maize event MON87429. Detection of one or more of this DNA in a sample may be used for determining the content or the source of the commodity product. Any standard method of detection for DNA molecules may be used, including methods of detection disclosed herein.

[0067] Maize event MON87429 contains four expression cassettes that together provide tolerance to inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group; synthetic auxins; inhibitors of glutamine synthetase; and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS).

[0068] As used herein, inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group (referred to as "FOP herbicide(s)") include, but are not limited to, clodinafop, clodinafop-ethyl, clodinafop-propargyl, cyhalofop, cyhalofop-butyl, diclofop, diclofop-methyl, diclofop-P, diclofop-P-methyl, fenoxaprop, fenoxaprop-P, fenoxaprop-P-ethyl, fenthiaprop, fluazifop, fluazifop-butyl, fluazifop-P, fluazifop-P-butyl, fluroxypyr, haloxyfop, haloxyfop-ethyl, haloxyfop-methyl, haloxyfop-P, haloxyfop-P-methyl, isoxapyrifop, metamifop, propaquizafop, quizalofop, quizalofop-ethyl, quizalofop-P, quizalofop-P-ethyl, quizalofop-P-tefuryl, and trifop.

[0069] As used herein, synthetic auxins include, but are not limited to, benzoic acid herbicides, phenoxy acid herbicides, arylpicolinate herbicides, and pyridinyloxy acid herbicides. Examples of a benzoic acid herbicides include, but are not limited to, dicamba (3,6-dichloro-2-methoxybenzoic acid), dicamba salts, dicamba-butyl, dicamba-diglycolamine salt, dicamba-dimethylammonium, dicamba-diethanolammonium, dicamba-isopropylammonium, dicamba-potassium, dicamba-sodium, and dicamba-trolamine. Examples of phenoxy acid herbicides are 2,4-D (2,4-dichlorophenoxyacetic acid), 2,4-D-butyl, 2,4-D-butyl, 2,4-D-choline, 2,4-D-dimethylammonium, 2,4-D-diolsamin, 2,4-D-ethyl, 2,4-D-2-ethylhexyl, 2,4-D-isobutyl, 2,4-D-isocetyl, 2,4-D-isopropyl, 2,4-D-isopropylammonium, 2,4-D-potassium, 2,4-D-sodium, 2,4-D-trisopropanolammonium, 2,4-D-trolamine, clomeprop, dichlorprop, fenoprop, MCPA, MCPA-butyl, MCPA-dimethylammonium, MCPA-2-ethylhexyl, MCPA-isopropylammonium, MCPA-potassium, MCPA-sodium, MCPA-thioethyl, 2,4-DB, MCPB, MCPB-methyl, MCPB-ethyl-sodium, and mecoprop. Examples of arylpicolinate herbicides are halauxifen, halauxifen-methyl, and florpyrauxifen-benzyl. Examples of pyridinyloxy acid herbicides are triclopyr, fluroxypyr, aminopyralid, clopyralid, and picloram.

[0070] As used herein, inhibitors of glutamine synthetase include, but are not limited to, phosphinothricin, glufosinate, glufosinate salts, glufosinate-ammonium, glufosinate-sodium, glufosinate-P, L-glufosinate-ammonium, and L-glufosinate-sodium.

[0071] As used herein, inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) include, but are not limited to, glyphosate, glyphosate salts, glyphosate-isopropylammonium, glyphosate-ammonium, glyphosate-dimethylammonium, glyphosate-trimesium (=sulfosate), glyphosate-diammonium, glyphosate-potassium, and glyphosate-sodium.

[0072] As used herein, "herbicide tolerant" or "herbicide tolerance" or "tolerance" means the ability to be wholly or partially unaffected by the presence or application of one of more herbicide(s), for example to resist the toxic effects of

an herbicide when applied. A cell, seed, or plant is “herbicide tolerant” or has “improved tolerance” if it can maintain at least some normal growth or phenotype in the presence of one or more herbicide(s). A trait is an herbicide tolerance trait if its presence can confer improved tolerance to an herbicide upon a cell, plant, or seed as compared to the wild-type or control cell, plant, or seed. Crops comprising an herbicide tolerance trait can continue to grow in the presence of the herbicide and may be minimally affected by the presence of the herbicide. A protein confers “herbicide tolerance” if expression of the protein can confer improved tolerance to an herbicide upon a cell, plant, or seed as compared to the wild-type or control cell, plant, or seed. Examples of herbicide tolerance proteins are phosphinothricin N-acetyltransferase, dicamba monooxygenase coding sequence, the FT_T protein, and glyphosate tolerant 5-enolpyruvylshikimate-3-phosphate synthase coding sequence from *Agrobacterium* sp strain CP4. Herbicide tolerance may be complete or partial insensitivity to a particular herbicide and may be expressed as a percent (%) tolerance or insensitivity to a particular herbicide.

[0073] As used herein, a “weed” is any undesired plant. A plant may be considered generally undesirable for agriculture or horticulture purposes (for example, *Amaranthus* species) or may be considered undesirable in a particular situation (for example, a crop plant of one species in a field of a different species, also known as a volunteer plant). Weeds are commonly known in the art and vary by geography, season, growing environment, and time. Lists of weed species are available from agricultural and scientific societies and efforts (such as the Weed Science Society of America, the Canadian Weed Science Society, the Brazilian Weed Science Society, the International Weed Science Society, and the International Survey of Herbicide Resistant Weeds), government agencies (such as the United States Department of Agriculture and the Australia Department of the Environment and Energy), and industry and farmer associations (such as the National Corn Growers Association).

[0074] The invention provides methods for controlling weeds in an area for maize cultivation by applying at least one herbicide selected from the group consisting of (i) inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group such as quizalofop and haloxyfop; (ii) synthetic auxins such as dicamba and 2,4-D; (iii) inhibitors of glutamine synthetase such as glufosinate; and (iv) the 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitor glyphosate, where seeds or plants comprising maize event MON87429 are planted in the area before, at the time of, or after applying the herbicide and the herbicide application prevents or inhibits weed growth and does not injure the maize plants. The plant growth area may or may not comprise weed plants at the time of herbicide application. The herbicide(s) used in the methods of the invention can be applied alone or in combination with one or more herbicide(s) during the growing season. The herbicide(s) used in the methods of the invention can be applied in combination with one or more herbicide(s) temporally (for example, as a tank mixture or in sequential applications), spatially (for example, at different times during the growing season including before and after maize seed planting), or both. For example, a method for controlling weeds is provided that consists of planting seed comprising maize event MON87429 in an area and applying

an herbicidally effective amount over the growing season of one or more of dicamba, glufosinate, glyphosate, 2,4-D, or a FOP herbicide, alone or in any combination with another herbicide, for the purpose of controlling weeds in the area without injuring the plants containing maize event MON87429. Such application of herbicide(s) may be pre-planting (any time prior to planting seed containing maize event MON87429, including for burn-down purposes, that is application to emerging or existing weeds prior to seed plant), pre-emergence (any time after seed containing maize event MON87429 is planted and before plants containing maize event MON87429 emerge), or post-emergence (any time after plants containing maize event MON87429 emerge). Multiple applications of one or more herbicides, or a combination of herbicides together or individually, may be used over a growing season, for example, two applications (such as a pre-planting application and a post-emergence application, or a pre-emergence application and a post-emergence application) or three or more applications (such as a pre-planting application and two post-emergence applications).

[0075] Herbicide application in practicing the methods of the invention may be at the recommended commercial rate or any fraction or multiple thereof, such as twice the recommended commercial rate. Herbicide rates may be expressed as acid equivalent per pound per acre (lb ae/acre) or active ingredient per pound per acre (lb ai/acre), depending on the herbicide and the formulation. The herbicide application may be the recommended commercial rate or a fraction or multiple thereof. The use of acres in the herbicide application rates as provided herein is merely instructive; herbicide application rates in the equivalent dosages to any rate provided herein may be used for areas larger or smaller than an acre. The herbicide application comprises at least one herbicide selected from the group consisting of (i) inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group such as quizalofop and haloxyfop; (ii) synthetic auxins such as dicamba and 2,4-D; (iii) inhibitors of glutamine synthetase such as glufosinate; and (iv) the 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitor glyphosate. The plant growth area may or may not comprise weed plants at the time of herbicide application. An herbicidally effective amount of the FOP herbicides for use in the area for controlling weeds should consist of a range from about 0.01 lb ai/acre to about 1.0 lb ai/acre over a growing season (for example, quizalofop could be applied at a rate of about 0.034 lb ai/acre to about 0.083 lb ai/acre and haloxyfop could be applied at a rate of about 0.018 lb ai/acre to about 0.07 lb ai/acre). An herbicidally effective amount of the synthetic auxin phenoxy acid herbicides for use in the area for controlling weeds should consist of a range from about 0.1 lb ae/acre to about 10 lb ae/acre over a growing season (for example, 2,4-D could be applied at a rate of about 0.75 lb ae/acre to 1.0 lb ae/acre). An herbicidally effective amount of the synthetic auxin pyridinyloxy acid herbicides for use in the area for controlling weeds should consist of a range from about 0.05 lb ae/acre to about 5.0 lb ae/acre over a growing season (for example, fluroxypyr could be applied at a rate of about 0.14 lb ae/acre to about 0.49 lb ae/acre). An herbicidally effective amount of a synthetic auxin benzoic acid herbicide for use in the area for controlling weeds should consist of a range from about 0.1 lb ae/ac to as much as about 16 lb ae/ac over a growing season (for example, dicamba could be applied at

a rate of about 0.5 lb ae/acre to about 2.0 lb ae/acre). An herbicidally effective amount of glutamine synthetase inhibitors for use in the area for controlling weeds should consist of a range from about 0.1 lb ae/acre to as much as about 10 lb ae/acre over a growing season (for example, glufosinate could be applied at a rate of about 0.4 lb ai/acre to about 1.59 lb ai/acre). An herbicidally effective amount of EPSPS inhibitors for use in the area for controlling weeds should consist of a range from about 0.5 lb ae/ac to about 12 lb ae/ac over a growing season (for example, glyphosate could be applied at a rate of about 0.75 lb ae/acre to about 2.25 lb ae/acre).

[0076] The invention provides methods for controlling volunteer maize comprising maize event MON87429 in an area for crop cultivation by applying an herbicidally effective amount of at least one cyclohexanedione (DIM) herbicide, such as clethodim, sethoxydim, and tralkoxydim, where the herbicide application prevents growth of maize comprising maize event MON87429. An herbicidally effective amount of a DIM herbicide for use in the area for controlling volunteer maize could be applied at a rate of about 0.03 lb ai/acre to about 2.75 lb ai/acre over a growing season (for example, clethodim could be applied from about 0.0625 lb ai/acre to about 0.125 lb ai/acre and sethoxydim could be applied from about 0.188 lb ai/acre to 0.281 lb ai/acre).

[0077] Methods for producing plants and seeds containing maize event MON87429 are provided. Plants may be bred using any method known in the art, for example, descriptions of breeding methods that are commonly used can be found in W R Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison Wis. (1987). Plants may be self-pollinated (also known as “selfing”) or cross-pollinated (also known as “crossing”). Plants containing maize event MON87429 may be self-pollinated to generate a true breeding line of plants that are homozygous for maize event MON87429. Selfing results in progeny known as “inbred” and can be used to produce inbred lines that are genetically uniform. Alternatively, plants containing maize event MON87429 may be cross-pollinated (bred with another plant that is transgenic or nontransgenic) to produce a varietal or a hybrid seed. Seed and progeny plants made by the methods of the invention contain maize event MON87429. Application of one or more herbicide for which maize event MON87429 confers tolerance may be used to select progeny that contain maize event MON87429. Alternatively, progeny may be analyzed using diagnostic methods to select for plants or seeds containing maize event MON87429. Progeny may be varietal or hybrid plants; may be grown from seeds produced by a plant containing maize event MON87429 or from seeds produced by a plant fertilized with pollen from a plant containing maize event MON87429; and may be homozygous or heterozygous for maize event MON87429.

[0078] Methods for producing hybrid seed using maize event MON87429 are provided. Plants comprising maize event MON87429 have expression of the glyphosate tolerant protein CP4-EPSPS in all tissues except male reproductive tissues. This results in glyphosate tolerance in vegetative and female reproductive tissues and glyphosate sensitivity in male reproductive tissues. This glyphosate sensitivity can be used to induce male-sterility through the proper application of glyphosate. Glyphosate is a systemic herbicide that is

translocated from source to sink tissues in plants. Due to the rate of glyphosate metabolism in maize, application to plants comprising maize event MON87429 prior to the development of the male reproductive tissue may prevent pollen development, pollen shed, or anther extrusion. This glyphosate-induced male-sterility can be used to increase the efficiency of hybrid seed production, for example by eliminating or reducing the need to physically emasculate the maize plant used as a female in a given cross during hybrid seed production.

[0079] The invention provides a method of producing hybrid seed comprising (a) growing a plant comprising maize event MON87429, (b) applying an effective amount of glyphosate to the plant to induce male-sterility, wherein the herbicide application is prior to or during the development of the male reproductive tissue of the plant thereby inducing male-sterility in the plant; (c) fertilizing the plant with pollen from a second plant; and (d) harvesting hybrid seed from the plant. In one embodiment, the glyphosate is applied prior to or during the development at an effective amount of about 0.25 lb ae/acre to about 11.0 lb ae/acre total, applied in one or more application. In another embodiment, the step of fertilizing may be accomplished by allowing passive fertilization (for example through wind pollination), by other means such as mechanical or hand pollination, or by a combination of these. The herbicide application may be applied in one or more application prior to or during the development of the male reproductive tissue, such as at a stage selected from the group consisting of the V4, V5, V6, V7, V8, V9, V10, V11, V12, V13, and V14 stage of maize plant development and may prevent at least pollen development, pollen shed, or anther extrusion. The male-sterility may be partial or complete. In one embodiment, the effective amount of glyphosate would be about 0.5 lb ae/acre to about 2.5 lb ae/acre total (in one application or split into two or more applications) applied at the V4 through V8 stage or up to 100 growing degree units (GDU) before flowering.

[0080] Plants, progeny, seeds, cells, and plant parts of the invention may also contain one or more additional maize trait(s) or transgenic events, particularly those introduced by crossing a maize plant containing maize event MON87429 with another maize plant containing the additional trait(s) or transgenic events. Such trait(s) or transgenic events include, but are not limited to, increased insect resistance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, improved nutritional quality, hybrid seed production, and herbicide tolerance, in which the trait is measured with respect to a maize plant lacking such transgenic trait. Maize transgenic events are known to one of skill in the art; for example, a list of such traits is provided by the United States Department of Agriculture’s (USDA) Animal and Plant Health Inspection Service (APHIS) and can be found on their website at www.aphis.usda.gov. Two or more transgenic events may thus be combined in a progeny seed or plant by crossing two parent plants each comprising one or more transgenic event(s), collecting progeny seed, and selecting for progeny seed or plants that contain the two or more transgenic events; these steps may then be repeated until the desired combination of transgenic events in a progeny is achieved. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation.

[0081] A deposit of a representative sample of seed comprising maize event MON87429 has been made according to

the Budapest Treaty with the American Type Culture Collection (ATCC®) having an address at 10801 University Boulevard, Manassas, Va. USA, Zip Code 20110. The ATCC Patent Deposit Designation (accession number) for seeds comprising maize event MON87429 is PTA-124635 and the date of deposit was Jan. 12, 2018. The deposit will be maintained in the depository for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer.

[0082] As used herein, the term “comprising” means “including but not limited to”.

EXAMPLES

[0083] The following examples are included to more fully describe the invention. Summarized are the construction and testing of thirty-five expression constructs, the production of over fifteen thousand unique events, and the analysis of hundreds of thousands of individual plants over seven years through the rigorous molecular, agronomic, and field testing required for the creation and ultimate selection of maize event MON87429.

[0084] It should be appreciated by those of skill in the art that many modifications can be made in the specific examples which are disclosed and still obtain a similar result. Certain agents which are both chemically and physiologically related may be substituted for the agents described herein while achieving the same or similar results. All such substitutions and modifications apparent to those skilled in the art are deemed to be within the scope of the invention.

Example 1: Expression Cassette Testing, Construct Design, and R0 Plant Testing

[0085] This example describes the design and testing in maize plants of thirty-five different constructs. Each construct contained four expression cassettes, each expression cassette used for expressing a different transgene. This testing was done to select the best construct to use for expressing all four transgenes in maize. Each construct had a unique configuration, varying by expression cassette orientation and expression elements.

[0086] Various individual expression cassettes with different expression element combinations and transgenes were designed, cloned into plant transformation vectors, and tested for trait efficacy in maize plants to create a pool of the best individual expression cassettes. Using this pool of individual expression cassettes, 35 different constructs were designed so that each contained four expression cassettes (each expression cassette having the transgene for PAT, DMO, CP4-EPSPS, or FT) and could be used to test the four-way combination of the different expression cassettes. The expression cassette combinations varied by expression elements, protein coding sequence, and orientation. This resulted in the testing of two PAT expression cassettes, 6 DMO expression cassettes, 5 FT expression cassettes, 18 CP4-EPSPS expression cassettes.

[0087] The 35 four-expression cassette constructs were cloned into plant transformation vectors, and these vectors were used for *Agrobacterium*-mediated transformation of LH244 maize immature embryos using methods known in the art to produce 15,326 unique transformation events, each made by a random insertion of the transgene insert into the maize genome. R0 plants were then regenerated from the

transgenic cells, and rooted plants with normal phenotypic characteristics were transferred to soil for growth and further assessment.

[0088] The 15,326 R0 plants were analyzed for having a single copy of the transgenic insert and absence of vector backbone sequence. Plants with a single copy of the insert were advanced for herbicide tolerance efficacy testing. Single copy R0 plants were assessed in the greenhouse for tolerance to quizalofop (0.16 lb ai/ac of Assure II® herbicide) followed by a tank mix of glufosinate (0.98 lb ae/ac of Ignite® 280 herbicide), dicamba (2.0 lb ae/ac of Clarity® herbicide), or 2,4-D (2.0 lb ae/ac 2,4-D Amine 4® herbicide) (or a combination of any of these) sprayed at the V1/V2 growth stage. Plants that showed >30% injury were discarded.

[0089] From the initial 15,326 unique transformation events produced using the 35 transformation vectors, 1,945 unique events were selected after analyzing the copy number and herbicide spray data. Data are provided in Table 2. The R0 plants for the selected events were self-pollinated to produce R1 seed or crossed to produce F1 seed that was advanced to first season field trials.

TABLE 2

First Season Field Trial Inbred Efficacy		
Construct	R0 Events	Events Advanced
HT4-1	515	39
HT4-2	263	30
HT4-3	158	53
HT4-4	340	79
HT4-5	97	20
HT4-6	121	24
HT4-7	49	10
HT4-8	88	16
HT4-9	853	112
HT4-10	366	55
HT4-11	600	78
HT4-12	729	73
HT4-13	459	63
HT4-14	1026	151
HT4-15	224	44
HT4-16	1877	76
HT4-17	365	39
HT4-18	127	24
HT4-19	94	27
HT4-20	23	4
HT4-21	409	74
HT4-25	123	13
HT4-26	57	3
HT4-27	76	1
HT4-29	118	5
HT4-30	177	24
HT4-31	311	18
HT4-32	913	123
HT4-34	884	124
HT4-36	1364	185
HT4-37	1019	179
HT4-38	1021	129
HT4-39	98	15
HT4-50	250	26
HT4-51	132	9
Total	15326	1945

Example 2: First Season Field Trials

[0090] Field trials were conducted over many years with events advanced from the R0 analysis for each of the 35 constructs, and the performance of every plant for each

construct in their first season of field trials was then analyzed as a set. Each construct was thus represented by many unique events. This allowed a larger number of constructs to be tested in first season field trials while only advancing beyond the first season trial the top performing constructs. Separate first season field trials were conducted with R2 plants (homozygous for each event) for (1) inbred efficacy for glufosinate+dicamba, quizalofop, and 2,4-D tolerance, (2) Roundup Hybridization System (RHS) efficacy, and (3) herbicide pressure testing for tolerance to higher application rates of herbicides quizalofop, 2,4-D, glufosinate, and dicamba.

[0091] An inbred efficacy screen of plants to evaluate herbicide tolerance for glufosinate+dicamba, quizalofop, and 2,4-D was carried out. Herbicide treatments consisted of: a tank-mix application of glufosinate at 0.8 lb ai/acre plus dicamba at 2.0 lb ae/acre; quizalofop at 0.16 lb ai/acre; or 2,4-D at 2.0 lb ai/acre. Plots were visually rated for crop injury 10-14 days after herbicide treatment on a scale of 0-100 with “0” being no crop injury and “100” being complete crop destruction. Plant height (PHT), ear height (EHT), days to 50% silk (S50D), days to 50% pollen (P50D), shell weight (SHW), test weight (TWT), moisture (MST), and grain yield (YLD) were also collected. All data were subjected to analysis of variance and means separation at $p < 0.05$. Overall averages for multiple plants containing the same event were used, and inbred efficacy was summarized for glufosinate+dicamba, quizalofop, and 2,4-D as excellent (4), good (3), fair (2), or poor (1) or not applicable (NA). Data are provided in Table 3.

TABLE 3

First Season Field Trial Inbred Efficacy			
Construct	Glufosinate/Dicamba	Quizalofop	2,4-D
HT4-1	4	4	4
HT4-2	4	4	4
HT4-3	4	4	4
HT4-4	4	4	4
HT4-5	4	4	4
HT4-6	4	4	4
HT4-7	4	4	4
HT4-8	4	4	4
HT4-9	4	4	4
HT4-10	4	4	4
HT4-11	4	4	4
HT4-12	4	4	4
HT4-13	4	4	4
HT4-14	4	4	4
HT4-15	4	4	4
HT4-16	4	4	4
HT4-17	4	4	4
HT4-18	4	4	3
HT4-19	4	4	4
HT4-20	4	4	4
HT4-21	4	4	4
HT4-25	4	4	4
HT4-26	NA	NA	NA
HT4-27	NA	NA	NA
HT4-29	4	4	4
HT4-30	4	4	4
HT4-31	4	4	4
HT4-32	4	4	4
HT4-34	4	4	4
HT4-36	4	4	4
HT4-37	4	4	4
HT4-38	4	4	4
HT4-39	4	4	4

TABLE 3-continued

First Season Field Trial Inbred Efficacy			
Construct	Glufosinate/Dicamba	Quizalofop	2,4-D
HT4-50	4	4	4
HT4-51	4	4	4

[0092] An RHS efficacy screen of plants was carried out to identify differences in glyphosate tolerance and glyphosate-induced tassel sterility of inbred material. A single herbicide treatment was used for screening and consisted of glyphosate at 1.5 lb ae/acre applied to V2 followed by 0.75 lb ae/acre to approximately V8 (875 growing degree days) followed by approximately V10 (1025 growing degree days). Plots were visually rated for % crop injury (CIPV2, CIPV8, CIPV10, and CIPVT) 10-14 days after herbicide application on a scale of 0 to 100 plus a final injury rating at VT (after tassel emergence). Plots were also visually rated for % silk emergence [(SES9A (S90) and SES9C (S90 plus 4 days) SES9E (S90 plus 8 days)] and % anther extrusion [(AES9A (S90), AES9C (S90 plus 4 days), and AES9E (S90 plus 8 days)] on a similar scale of 0 to 100 with “0” being no silk emergence or anther extrusion and “100” being complete silk emergence or anther extrusion. Other agronomic parameters were collected, as in the inbred efficacy screen. Overall averages for multiple plants containing the same event were used, and glyphosate tolerance, tassel sterility, and yield were each summarized as excellent (4), good (3), fair (2), or poor (1) or not applicable (NA). Data are provided in Table 4.

TABLE 4

First Season Field Trial RHS Efficacy			
Construct	Glyphosate Tolerance	Tassel Sterility	Yield
HT4-1	4	2	NA
HT4-2	4	2	NA
HT4-3	2	2	NA
HT4-4	4	2	NA
HT4-5	2	2	NA
HT4-6	2	2	NA
HT4-7	4	2	NA
HT4-8	4	2	NA
HT4-9	2	2	NA
HT4-10	2	2	NA
HT4-11	2	2	NA
HT4-12	4	2	NA
HT4-13	4	2	NA
HT4-14	4	4	4
HT4-15	4	2	NA
HT4-16	3/2	4	1
HT4-17	4	2	NA
HT4-18	2	2	NA
HT4-19	1	NA	NA
HT4-20	1	NA	NA
HT4-21	4	2	NA
HT4-25	2	2	NA
HT4-26	NA	NA	NA
HT4-27	NA	NA	NA
HT4-29	1	NA	NA
HT4-30	4	2	NA
HT4-31	2	2	NA
HT4-32	4	4	4
HT4-34	4	4	4
HT4-36	4	2	NA
HT4-37	4	2	NA
HT4-38	4	2	NA
HT4-39	4	2	NA

TABLE 4-continued

First Season Field Trial RHS Efficacy			
Construct	Glyphosate Tolerance	Tassel Sterility	Yield
HT4-50	2	2	NA
HT4-51	2	2	NA

[0093] An herbicide pressure test was conducted to evaluate construct level crop tolerance to higher application rates of herbicides quizalofop, 2,4-D, glufosinate, and dicamba as follows. Quizalofop treatments consisted of: 1) quizalofop at 0.32 lb ai/acre (4x) plus 0.25% v/v non-ionic surfactant (NIS) applied to VE-V2 followed by V4 followed by V8; 2) quizalofop at 0.64 lb ai/acre (8x) plus 0.25% v/v NIS applied to VE-V2 followed by V4 followed by V8; or 3) quizalofop at 1.28 lb ai/acre (16x) plus 0.25% v/v NIS applied to VE-V2 followed by V4 followed by V8. 2,4-D treatments consisted of 1) 2,4-D amine at 2 lb ai/acre plus 0.25% v/v non-ionic surfactant (NIS) applied to VE-V2 followed by V4 followed by V8; 2) 2,4-D amine at 4 lb ai/acre plus 0.25% v/v NIS applied to VE-V2 followed by V4 followed by V8; 3) 2,4-D amine at 8 lb ai/acre plus 0.25% v/v NIS applied to VE-V2 followed by V4 followed by V8; or 4) 2,4-D amine at 16 lb ai/acre plus 0.25% v/v NIS applied to VE-V2 followed by V4 followed by V8. Glufosinate treatments consisted of: 1) glufosinate 1.0 lb ai/acre applied to V2 followed by V4 followed by V8; 2) glufosinate 2.0 lb ai/acre applied to V2 followed by V4 followed by V8; 3) glufosinate 4.0 lb ai/acre applied to V2 followed by V4 followed by V8; or 4) glufosinate 8.0 lb ai/acre applied to V2 followed by V4 followed by V8. Dicamba treatments consisted of: 1) dicamba at 2.0 lb applied to V2 followed by V4 followed by V8; 2) dicamba at 4.0 lb applied to V2 followed by V4 followed by V8; or 3) dicamba at 8.0 lb applied to V2 followed by V4 followed by V8. 4) dicamba at 16 lb applied to V2 followed by V4 followed by V8. Plots were rated visually for crop injury and agronomic parameters were collected, as in the inbred efficacy screen. Overall averages for multiple plants containing the same event were used, and herbicide tolerance efficacy was summarized for each of the four herbicides as excellent (4), good (3), fair (2), or poor (1) or not applicable (NA). Data are provided in Table 5.

TABLE 5

First Season Field Trial Herbicide Pressure Test				
Construct	Glufosinate	Dicamba	Quizalofop	2,4-D
HT4-1	4	4	3	3
HT4-2	4	4	4	4
HT4-3	4	4	4	4
HT4-4	4	4	4	4
HT4-5	4	4	4	4
HT4-6	4	4	4	4
HT4-7	4	4	4	4
HT4-8	4	4	4	4
HT4-9	4	4	4	4
HT4-10	4	4	4	4
HT4-11	4	4	4	4
HT4-12	4	4	4	4
HT4-13	4	4	4	4
HT4-14	4	4	4	4
HT4-15	4	4	4	4
HT4-16	4	4	4	4
HT4-17	4	4	4	4
HT4-18	4	4	4	4

TABLE 5-continued

First Season Field Trial Herbicide Pressure Test				
Construct	Glufosinate	Dicamba	Quizalofop	2,4-D
HT4-19	4	4	4	4
HT4-20	4	4	4	4
HT4-21	4	4	4	4
HT4-25	4	4	4	4
HT4-26	NA	NA	NA	NA
HT4-27	NA	NA	NA	NA
HT4-29	4	4	4	4
HT4-30	4	4	4	4
HT4-31	4	4	4	4
HT4-32	4	4	4	4
HT4-34	4	4	4	4
HT4-36	4	4	4	4
HT4-37	4	4	4	4
HT4-38	4	4	4	4
HT4-39	4	4	4	4
HT4-50	4	4	4	4
HT4-51	4	4	4	4

[0094] The data of the composite performance of R2 plants produced with each of 35 constructs was compiled and analyzed for (1) the inbred efficacy test for glufosinate+dicamba, quizalofop, and 2,4-D tolerance, (2) the RHS efficacy test for glyphosate tolerance, tassel sterility, and yield, and (3) the herbicide pressure testing for tolerance to higher application rates of herbicides quizalofop, 2,4-D, glufosinate, and dicamba. Using this data, 3 constructs (HT4-14, HT4-32, and HT4-34) were selected for advancement from the 35 constructs tested. Events for these 3 constructs were then advanced to second season field trials.

Example 3: Molecular Analysis

[0095] Molecular analysis was conducted concurrently with the field trials on events that were advanced. DNA amplification and sequencing were used to confirm the insert sequence, insert copy number, and absence of backbone in the insert. The insertion site in the maize genome for each event was mapped. Northern analysis was done to detect and measure mRNA transcripts of the pat, dmo, ft_t, and cp4-epsps genes. N-terminal protein sequencing of the PAT, DMO, FT_T, and CP4-EPSPS proteins purified from transgenic plants was done to confirm the recombinant protein sequence. Western blot analysis to detect the PAT, DMO, FT_T, and CP4-EPSPS proteins was done with transgenic plant samples. In depth Southern analysis was performed on genomic DNA from R1 plants to confirm copy number and the absence of backbone.

Example 4: Advanced Field Trials

[0096] Advanced field trials (second season testing and beyond) were conducted over many years with events advanced from the first season field trials for constructs HT4-14, HT4-32, and HT4-34. The performance of many individual plants for each event in each field trial was analyzed as a set. Each event was thus represented by many unique plants. This allowed the performance of each event to be analyzed under many conditions, in different locations and geographies, and for a variety of properties.

[0097] Field trials were conducted with inbred plants (homozygous for the event) and hybrid plants (hemizygous for the event) to assess (1) trait efficacy for commercial rates of glufosinate, dicamba, quizalofop, and 2,4-D tolerance, (2)

agronomic performance, (3) Roundup Hybridization System (RHS) efficacy and glyphosate, and (4) herbicide pressure testing for tolerance to higher application rates of quizalofop, 2,4-D, glufosinate, and dicamba herbicides.

[0098] In the Field Season 1 trials, 30 events were tested for construct HT4-14, 41 events were tested for construct HT4-32, and 21 events were tested for construct HT4-34. Using the composite data from these trials, events were selected for advancement. In the Field Season 2 trials, 15 events were tested for the HT4-14 construct, 38 events for the HT4-32 construct, and 38 events for the HT4-34 construct (this number included some events not tested in Field Season 1 due to because of seed shortages during that season). Events for construct HT4-14 were characterized by molecular analysis as described in Example 3 and this data was also used to select events. The composite data from these trials and the in depth molecular characterization of events for the HT4-14 construct was used to select 3 events for advancement for the HT4-14 construct. The composite data from these trials was used to select 24 events for advancement for the HT4-32 construct and to decide not to advance any events for the HT4-34 construct. In the Field Season 3 trials, the 3 events for construct HT4-14 and 24 events for the HT4-34 construct were tested. The composite data from these trials was used to select 2 events for advancement for the HT4-14 construct and to decide not to advance any events for the HT4-32. The Field Season 4 trials were used to compare the final two events in a large number of locations, under a variety of conditions, and in hybrid and inbred germplasm in order to produce the data necessary to select the superior event. Table 6 provides the number of unique events tested for each construct in the field trials conducted during each season.

TABLE 6

Advanced Field Trials Summary			
Milestone	HT4-14	HT4-32	HT4-34
Field Season 1	30	41	21
Field Season 2	15	38	38
Field Season 3	3	24	0
Field Season 4	2	0	0
Final event selection	1	0	0

[0099] The agronomic performance field trials were run during the same season as the trait efficacy field trials. All field trials used a randomized complete block design and were conducted at multiple locations. Field trials were conducted at locations in North America and South America. For both efficacy and agronomic field trials, agronomic scoring was collected throughout the field trial season, and at the end of the season yield was determined (efficacy yield or agronomic yield). Efficacy field trials were conducted to assess crop injury 10 to 14 days following herbicide application. The target crop injury rating was a score of less than 10% for advancement of an event. For agronomic field trials, the plots were maintained weed free and none of the test herbicides were applied during the growing season. The hybrid agronomic field trials included controls of a comparable hybrid (hybrid control) produced using the same parental maize lines used to make the transgenic hybrid cross, but not containing a transgenic event. Inbred controls were a comparable inbred to the transgenic inbred lines.

[0100] To compare the field trial data, meta-analysis was performed using the aggregate of all plants in the multi-season, multi-location field trial data. As an example, Table 7 illustrates over multiple seasons the number of replications (reps) for which an observation was repeated for two selected HT4-14 events and the total number of individual plants tested in each trial for each event.

TABLE 7

Field trial replications			
Milestone	Event	Total Reps	Total Plants
Field Season 1	EVENT 2	235	16,450
Field Season 1	MON87429	235	16,450
Field Season 2	EVENT 2	100	7,000
Field Season 2	MON87429	100	7,000
Field Season 3	EVENT 2	529	37,030
Field Season 3	MON87429	513	35,910
Field Season 4	EVENT 2	228	15,960
Field Season 4	MON87429	228	15,960
Field Season 4	EVENT 2	56	3,920
Field Season 4	MON87429	2465	172,550

[0101] Meta-analysis of the multiple hybrid efficacy field trials was completed for comparison of the hybrid injury ratings. As an example, Table 8 provides the injury rating over multiple seasons for two selected HT4-14 events scored at V8 (where V8 analysis encompasses the cumulative injury from V2, V4, V6, and V8 herbicide applications) with a statistical least significant difference at 95% confidence level (LSD at $p < 0.05$). Plants containing maize event MON87429 and plants containing Event 2 both performed well in these trials.

TABLE 8

Meta-analysis of injury rating from hybrid efficacy field trials			
Milestone	Event	V8 injury	LSD ($P < 0.05$)
Field Season 1	MON87429	1.3	2.1
Field Season 1	EVENT 2	1.1	2.1
Field Season 2	MON87429	1.7	3.1
Field Season 2	EVENT 2	2.6	3.1
Field Season 3	MON87429	0.08	3.4
Field Season 3	EVENT 2	0.13	3.4

[0102] Meta-analysis of the multiple hybrid efficacy field trials was completed for comparison of yield as bushels/acre (Bu/ac). As an example, Table 9 provides yield from hybrids over multiple seasons for two selected HT4-14 events with a statistical least significant difference at 95% confidence level (LSD at $p < 0.05$). Plants containing maize event MON87429 and plants containing Event 2 both performed well in these trials.

TABLE 9

Meta-analysis of yield from hybrid efficacy field trials			
Milestone	Event	Yield (Bu/ac)	LSD ($p < 0.05$)
Field Season 1	MON87429	222	12
Field Season 1	EVENT 2	222	12
Field Season 2	MON87429	197	15.4

TABLE 9-continued

Meta-analysis of yield from hybrid efficacy field trials			
Milestone	Event	Yield (Bu/ac)	LSD (p < 0.05)
Field Season 2	EVENT 2	197	15.4
Field Season 3	MON87429	210.9	9.1
Field Season 3	EVENT 2	211.5	9.1

[0103] Pressure testing field trials with herbicide applied at higher application rates than commercial use were conducted with hybrid and inbred plants containing a single transgenic event. The herbicides glufosinate (over the range of 1.6 to 6.4 lb ai/acre), dicamba (over the range of 2.0 to 16 lb ai/acre), quizalofop (over the range of 0.32 to 1.28 lb ai/acre), 2,4-D (over the range of 2.0 to 8.0 lb ai/acre), and glyphosate (at 3.0 lb ae/acre) were applied in field trials for pressure testing of the efficacy of the herbicide tolerance traits. At the end of the season, the hybrid pressure testing field trials were harvested and yield (Bu/ac) was determined. As an example, Table 10 provides yield data from hybrids and inbreds in different trials for two selected HT4-14 events with a statistical least significant difference at 95% confidence level (LSD at p<0.05). Plants containing maize event MON87429 and plants containing Event 2 both performed well in hybrid and inbred yield trials for all of the herbicide treatments. The results suggested that further field testing could identify an inbred yield advantage for maize event MON87429 over Event 2.

TABLE 10

Yield from hybrid/inbred pressure testing efficacy field trials				
Herbicide	Plant	Event	Yield (Bu/ac)	LSD (p < 0.05)
Glufosinate	Hybrid	MON87429	256	28.4
Glufosinate	Hybrid	EVENT 2	240	28.4
Dicamba	Hybrid	MON87429	264	39.2
Dicamba	Hybrid	EVENT 2	256	39.2
Quizalofop	Hybrid	MON87429	251	48.5
Quizalofop	Hybrid	EVENT 2	257	48.5
2,4-D	Hybrid	MON87429	261	38.4
2,4-D	Hybrid	EVENT 2	254	38.4
Glyphosate	Inbred	MON87429	90.1	46
Glyphosate	Inbred	EVENT 2	67.2	46

[0104] Hybrid agronomic field trials were conducted, agronomic measures were collected throughout the season, and agronomic yield was determined at the end of the season. Meta-analysis across the multi-season, multi-location hybrid agronomic field trials was used to compare the yield of the hybrid control and the hybrids. As an example, Table 11 provides yield data (Bu/ac) for two selected HT4-14 events with a statistical least significant difference at 95% confidence level (LSD at p<0.05). Plants containing maize event MON87429 and plants containing Event 2 both performed well in these trials, and no statistical difference in hybrid yield was found for plants containing either of these events when compared to the control plants.

TABLE 11

Meta-analysis of yield from hybrid agronomic field trials			
Milestone	Event	Yield (Bu/ac)	LSD (p < 0.05)
Field Season 1	Control - none	222	9
Field Season 1	MON87429	220	9
Field Season 1	EVENT 2	224	9
Field Season 2	Control - none	224.6	13.7
Field Season 2	MON87429	211	13.7
Field Season 2	EVENT 2	215	13.7
Field Season 3	Control - none	213.6	9.7
Field Season 3	MON87429	213.3	9.7
Field Season 3	EVENT 2	212.4	9.7

[0105] Inbred efficacy field trials were conducted for glyphosate tolerance and the Roundup Hybridization System (RHS) and yield was determined at the end of the season. Glyphosate was applied at a rate of 1.5 lb ae/acre for weed control followed by two sterility applications of glyphosate at 0.75 lb ae/acre at approximately V8 followed by 0.75 lb ae/acre at approximately V10. Meta-analysis across the multi-season, multi-location inbred efficacy field trials was used to compare the yield of plants. As an example, Table 12 provides yield data (Bu/ac) for two selected HT4-14 events with a statistical least significant difference at 95% confidence level (LSD at p<0.05). Field Season 2 and Field Season 3 trials both showed a statistically significant decrease in yield in plants containing Event 2 when compared to plants containing the Event MON87429. These data indicated the superior performance in inbred efficacy yield trials of plants containing maize event MON87429.

TABLE 12

Meta-analysis of yield from glyphosate treated inbred efficacy field trials			
Milestone	Event	Yield (Bu/ac)	LSD (p < 0.05)
Field Season 1	MON87429	89.9	14.3
Field Season 1	EVENT 2	87.5	14.3
Field Season 2	MON87429	119.6	18.0
Field Season 2	EVENT 2	99.0	18.0
Field Season 3	MON87429	105.6	10.5
Field Season 3	EVENT 2	90.13	10.5

[0106] Inbred agronomic field trials were conducted and yield was determined at the end of the season for untreated plants. The trials included controls of inbred lines comparable to the transgenic inbred lines. Meta-analysis across the multi-season, multi-location inbred agronomic field trials was conducted comparing yield for the paired control and the transgenic inbreds. As an example, Table 13 provides yield data (Bu/ac) for two selected HT4-14 events with a statistical least significant difference at 95% confidence level (LSD at p<0.05). No statistical difference in inbred agronomic yield was found between the control plants and plants containing maize event MON87429. In contrast, for Field Season 3 trials, there was a statistically significant decrease in yield in plants containing Event 2 when compared to control plants and plants containing maize event MON87429. These data indicated the superior performance in inbred agronomic yield trials of plants containing maize event MON87429.

TABLE 13

Meta-analysis of yield from inbred agronomic field trials			
Milestone	Event	Total (Bu/ac)	LSD (p < 0.05)
Field Season 1	Control - none	105.2	8.9
Field Season 1	MON87429	104.1	8.9
Field Season 1	EVENT 2	102.5	8.9
Field Season 2	Control - none	103.7	15.3
Field Season 2	MON87429	93.4	15.3
Field Season 2	EVENT 2	91.4	15.3
Field Season 3	Control - none	116.6	6.2
Field Season 3	MON87429	112.8	6.2
Field Season 3	EVENT 2	106.1	6.2

[0107] Hybrid efficacy trials were conducted at four locations in Argentina to evaluate plant tolerance to glufosinate, dicamba, quizalofop, haloxyfop, 2,4-D, and glyphosate. Plants containing MON87429 were crossed with plants containing both maize event MON88017 and maize event MON89034 to produce progeny containing all three events (MON87429×MON88017×MON89034). Herbicide treatments consisted of 1) a non-treated control; 2) glufosinate at 0.448 kg ai/ha applied at the V2 stage followed by the same application to the V6 stage; 3) glufosinate at 0.896 kg ai/ha applied at the V2 stage followed by the same application to the V6 stage; 4) dicamba at 0.56 lb ae/acre applied at the V2 stage followed by the same application to the V6 stage; 5) dicamba at 1.12 lb ae/acre applied at the V2 stage followed by the same application to the V6 stage; 6) quizalofop at 0.09 kg ae/ha applied at the V2 stage followed by the same application to the V6 stage; 7) quizalofop at 0.18 kg ae/ha applied at the V2 stage followed by the same application to the V6 stage; 8) haloxyfop at 0.1 kg ae/ha applied at the V2 stage followed by the same application to the V6 stage; 9) haloxyfop at 0.2 kg ae/ha applied at the V2 stage followed by the same application to the V6 stage; 10) 2,4-D at 1.12 lb ai/acre applied at the V2 stage followed by the same application to the V6 stage; 11) 2,4-D at 2.24 lb ae/acre applied at the V2 stage followed by the same application to the V6 stage; or 12) glyphosate at 2.24 lb ae/acre applied at the V2 stage followed by the same application to the V6 stage. Data collection consisted of crop injury 10 to 14 days after V2 and V6 herbicide applications and a final rating at VT, days to 50% pollen, days to 50% silk, plant height, ear height, shell weight, test weight, moisture, and grain yield. All data were subjected to analysis of variance and means separated at p<0.05.

[0108] Herbicide tolerance of plants containing MON87429×MON88017×MON89034 was excellent (<10% crop injury) over all rates of glyphosate, glufosinate, dicamba, quizalofop, haloxyfop, and 2,4-D tested. Herbicide treatment rates did not produce differences with respect to visual crop injury. Ear height was not significantly different for any of the treatments compared to the standard herbicide treatment 12 (glyphosate at 2.24 lb ae/acre). There was no significant reduction in plant height or test weight, increase in grain moisture, delay in maturity (measured as increase days to 50% pollen or silk), decrease in grain yield of the plants containing MON87429×MON88017×MON89034 within any treatment over that of non-treated plants. Hybrid plants produced through the cross of a plant containing MON87429 with a plant containing an event providing glyphosate tolerance in male tissues (such as commercially available maize events MON88017 or NK603) provide

excellent vegetative tolerance to glyphosate, glufosinate, dicamba, quizalofop, haloxyfop, and 2,4-D when applied at commercial label rates.

[0109] Three years of field trials were used to test for control of plants comprising maize event MON87429 using clethodim. These trials assessed the use in volunteer control methods of a DIM herbicide with plants containing maize event MON87429. Plants were treated with clethodim at commercial label rates and complete control of plants comprising maize event MON87429 was observed.

[0110] The data accumulated from the molecular analysis and from the field trials with inbred and hybrid plants assessing (1) trait efficacy for commercial rates of glufosinate, dicamba, quizalofop, and 2,4-D tolerance, (2) agronomic performance, (3) Roundup Hybridization System (RHS) efficacy and glyphosate tolerance, and (4) herbicide pressure testing for tolerance to higher application rates of quizalofop, 2,4-D, glufosinate, and dicamba herbicides was analyzed for all the events tested for constructs HT4-14, HT4-32, and HT4-34. Analysis of the cumulative data demonstrated the overall superior performance of maize event MON87429 compared to the other events and resulted in selection of this event for commercial purposes.

Example 5: Molecular Characterization of Maize Event MON87429

[0111] Maize event MON87429 was subjected to extensive molecular characterization upon selection as a commercial event. The transgenic insert of maize event MON87429 contains the elements and sequences described in Table 1.

[0112] DNA sequence analysis of maize event MON87429 was conducted. Southern blot analysis was conducted to confirm that plants containing maize event MON87429 contained a single, intact copy of the entire transgenic insert without any transformation vector backbone. Flanking DNA was sequenced on both the 5' and 3' ends of the insert, and the respective junctions were determined using sequence capture, enrichment, sequencing, inverse PCR, and genome walking techniques. The sequences of the flanking DNA for maize event MON87429 were mapped to the known maize genome physical assembly. The insertion site sequence information was used for bioinformatics analysis of the chromosomal location of the event. Insertion site integrity was determined by PCR across the wild-type allele using primers specific to the flanking regions of maize event MON87429. The wild-type insertion site was used to map the unique site of transgene integration for maize event MON87429 to the maize reference genome. To ensure that no alterations or mutations were introduced to any region of the transgene during transformation, the entire transgenic insert of maize event MON87429 was isolated from the plant and sequenced. Sequence information for the 5' junction, 3' junction, and transgenic insert are provided herein as SEQ ID NOs:1-10.

[0113] RNA analysis of plants comprising the construct of maize event MON87429 was conducted. Northern analysis was conducted on total RNA isolated from grain of plants containing maize event MON87429. This confirmed the transcript size and number for the pat, dmo, ft_t, and cp4-epsps mRNA products. RNA expression levels for CP4-EPSPS were also measured by real-time PCR using samples from plant tissues containing maize event MON87429. Rapid amplification of cDNA ends (RACE) was used to identify CP4-EPSPS cleavage products to confirm that

cleavage of the CP4-EPSPS transcript occurs only in tassels of plants comprising maize event MON87429 and that this is triggered by maize endogenous male-specific small interference RNAs (siRNAs) in a sequence-specific manner. Low molecular weight Northern analysis was performed to demonstrate that there are no CP4-EPSPS siRNAs that could compromise glyphosate tolerance in non-tassel tissues.

[0114] Protein analysis of plants comprising the construct of maize event MON87429 was conducted. N-terminal protein sequencing of the expressed PAT, DMO, FT_T, and CP4-EPSPS proteins was performed using immunopurified protein extracts from grain to confirm the authentic N-terminal amino acid sequence. Western blot analysis was conducted on protein extracts from grain containing maize event MON87429 to confirm that a single expected-sized protein was being produced for PAT, DMO, FT_T, and CP4-EPSPS, respectively. ELISAs was used to determine protein levels in the leaf, seed, roots, and pollen of plants for the PAT, DMO, FT_T, and CP4-EPSPS proteins.

Example 6: Detection of Maize Event MON87429

[0115] Detection of maize event MON87429 in a sample can be done using DNA, RNA, or protein detection techniques. Exemplary detection methods and materials are provided below. Detection may determine the presence or absence of maize event MON87429 in a sample. Detection may indicate the number of genomic copies of maize event MON87429 (that is, hemizygous, homozygous, or heterozygous) in a sample of genomic DNA.

[0116] An event specific endpoint Applied Biosystems™ TAQMAN® thermal amplification method (Thermo Fisher Scientific) was developed to identify maize event MON87429 in a sample. The DNA primers and probe used in the endpoint assay are primers SQ51062 (SEQ ID NO:11), SQ51053 (SEQ ID NO:12), and 6-FAM™ labeled probe PB50370 (SEQ ID NO:13). 6-FAM is a fluorescent dye product of Applied Biosystems (Foster City, Calif.) attached to the DNA probe. For TAQMAN MGB™ probes, the 5' exonuclease activity of Taq DNA polymerase cleaves the probe from the 5'-end, between the fluorophore and quencher. When hybridized to the target DNA strand, quencher and fluorophore are separated enough to produce a fluorescent signal, thus releasing fluorescence. SQ51062 and SQ51053 when used with these reaction methods and PB50370 produce a DNA amplicon that is diagnostic for maize event MON87429. The controls for this analysis should include a positive control containing maize event MON87429, a negative control from non-transgenic maize, and a negative control that contains no template DNA. Additionally, a control for the PCR reaction should optimally include Internal Control Primers and an Internal Control Probe, specific to a single copy gene in the maize genome. These assays are optimized for use with the Applied Biosystems GeneAmp® PCR System 9700 (Thermo Fisher Scientific) run at maximum speed, but other equipment may be used.

[0117] An example of conditions useful with TAQMAN methods for detection of maize event MON87429 is as follows. Step 1: 18 megohm water adjusted for final volume of 5 µl. Step 2: 2.28 µl of 2× Universal Master Mix (dNTPs, enzyme, buffer) to a 1× final concentration. Step 3: 0.05 µl Event Primer-1 (SQ51062) and Event Primer-2 (SQ51053) (resuspended in 18 megohm water to a concentration of 100 µM for each primer) to 0.9 µM final concentration. Step 4:

0.01 µl Event 6-FAM MGB Probe PB50370 (resuspended in 18 megohm water to a concentration of 100 µM) to 0.2 µM final concentration. Step 5: 0.05 µl Internal Control Primer-1 and Internal Control Primer-2 Mix (resuspended in 18 megohm water to a concentration of 100 µM for each primer) to 0.9 µM final concentration. Step 6: 0.01 µl Internal Control VIC™ Probe (resuspended in 18 megohm water to a concentration of 100 µM) to 0.2 µM final concentration. Step 7: 2.5 µl Extracted DNA (template) for each sample with one each of the following comprising: (a) Leaf Samples to be analyzed; (b) Negative control (non-transgenic DNA); (c) Negative water control (no template); and (d) Positive control maize containing maize event MON87429 DNA. Step 8: Thermocycler Conditions as follows: one cycle at 95° C. for 20 seconds; forty cycles of 95° C. for 3 seconds then 60° C. for 20 seconds; and final cycle of 10° C.

[0118] A zygosity assay is developed to determine whether a plant comprising maize event MON87429 is heterozygous or homozygous for the event or the wild-type allele. An amplification reaction assay can be designed using the sequence information provided herein. For example, such a PCR assay would include design of at least three primers: primer-1, primer-2, and primer-3, where primer-1 is specific to maize genomic DNA on the 3' flank of maize event MON87429; primer-2 is specific to maize event MON87429 transgenic insert; and primer-3 is specific to the wild-type allele. When used as a primer pair in an amplification reaction, primer-1 with primer-2 will produce a PCR amplicon specific for maize event MON87429. When used as a primer pair in an amplification reaction, primer-1 with primer-3 will produce a PCR amplicon specific for wild-type allele. In a PCR reaction performed on maize genomic DNA, the respective PCR amplicons generated from primer-1+primer-2 and that generated from primer-1+primer-3 will differ in sequence and size of the amplicon. When the three primers are included in a PCR reaction with DNA extracted from a plant homozygous for maize event MON87429, only the primer-1+primer-2 amplicon (specific for the maize MON87429 insertion) will be generated. When the three primers are included in a PCR reaction with DNA extracted from a plant heterozygous for maize event MON87429, both the primer-1+primer-2 amplicon (specific for the maize MON87429 insertion) and the primer-1+primer-3 amplicon (specific for wild-type allele or absence of the maize MON87429 insertion) will be generated. When the three primers are mixed together in a PCR reaction with DNA extracted from a plant that is null for maize event MON87429 (that is wild-type), only the primer-1+primer-3 amplicon (specific for wild-type allele) will be generated. The amplicons produced using the PCR reaction may be identified or distinguished using any method known in the art.

[0119] Another zygosity assay for maize event MON87429 is a TAQMAN thermal amplification reaction. For this type of assay, in addition to primers as described above, the assay would include two fluorescently labeled probes. Probe-1 would be specific for maize event MON87429, and probe-2 would be specific for a maize plant that is null for maize event MON87429 (wild-type), and where the two probes contain different fluorescent labels, for example the 6-FAM-label or VIC™-label. When used in a TAQMAN reaction, primer-1+primer-2+probe-1 will produce a first fluorescent signal specific for maize event

MON87429 and primer-1+primer-3+probe-2 will produce a second fluorescent signal specific for wild-type maize. When the three primers and two probes are included in a TAQMAN reaction with DNA extracted from a plant homozygous for maize event MON87429, only the first fluorescent signal (specific to primer-1+primer-2+probe-1) will be generated. When the three primers are included in a TAQMAN reaction with DNA extracted from a plant heterozygous for maize event MON87429, both the first fluorescent signal (specific to primer-1+primer-2+probe-1) and the second fluorescent signal (specific to primer-1+primer-3+probe-2) will be generated. When the three primers are mixed together in a TAQMAN reaction with DNA extracted from a plant which is null for maize event MON87429 (wild-type), only the second fluorescent signal (specific to primer-1+primer-3+probe-2) will be generated.

[0120] Another method to detect the presence of maize event MON87429 in a plant sample would be Southern analysis. One of skill in art would understand how to design Southern hybridization probe(s) specific for maize event MON87429 and a second southern hybridization probe specific for a maize plant which is null for maize event MON87429 (wild-type). With Southern analysis, a signal detected only from the first Southern hybridization probe will be indicative of a plant homozygous for maize event MON87429; a signal detected from both the first Southern hybridization probe and the second Southern hybridization probe will be indicative of a plant heterozygous for maize event MON87429; and a signal detected only from the second Southern hybridization probe will be indicative that the DNA was extracted from a plant that is null for maize event MON87429 (wild-type).

[0121] Another example of a detection kit comprises at least one antibody specific for at least one protein encoded by maize event MON87429. For example, such a kit may

utilize a lateral flow strip comprising reagents activated when the tip of the strip is contacted with an aqueous solution. Exemplary proteins sufficient for use in antibody production are ones encoded by the sequence provided as SEQ ID NO:10, or any fragment thereof.

[0122] A protein detection method is developed to determine whether a sample is from a plant, seed, cell, or plant part comprising maize event MON87429. At least one antibody specific for at least one protein encoded by maize event MON87429 is used to detect a protein encoded by maize event MON87429 in a sample. A detection kit comprising one or more antibodies specific for one or more proteins encoded by maize event MON87429 may utilize a lateral flow strip containing reagents activated when the tip of the strip is contacted with an aqueous solution. Samples of maize tissue may be ground up and protein extracted for analysis using water or an aqueous buffer (e.g., phosphate buffered saline containing detergent and bovine serum albumin). Following centrifugation, the aqueous supernatant is analyzed using the ELISA method in a sandwich format on a lateral flow strip containing an absorbent pad. Detection is activated by dipping the tip of the strip into the aqueous solution containing the sample to be tested. The aqueous solution is carried up the strip by capillary action and solubilizes gold labeled antibodies on the strip. The gold labeled antibodies are specific for at least one protein encoded by maize event MON87429 and will bind to an epitope on the protein in the sample to form an antibody-antigen complex. The gold labeled antibody-antigen complex is then carried up the strip to a nitrocellulose membrane. The membrane comprises a test line of immobilized antibodies that bind to a second, separate epitope on the protein encoded by maize event MON87429, causing a visible line to appear across the test strip if the protein encoded by maize event MON87429 is present in the sample.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 1

ccactcttgt ttggtttcat gtccgggaaa

30

<210> SEQ ID NO 2

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 2

cggtgcacaa actatagcaa gtgtgttcta

30

<210> SEQ ID NO 3

<211> LENGTH: 60

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 3

acttgcaaaa atagaccact cttgtttggt ttcattgtccg ggaaatctac atggatcagc 60

<210> SEQ ID NO 4
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 4

gcacaacaaa cgcaccggtg cacaaactat agcaagtgtg gtctattttt atatagcaag 60

<210> SEQ ID NO 5
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 5

agatactctt gtttggttac acttgcaaaa atagaccact cttgtttggt ttcattgtccg 60

ggaaatctac atggatcagc aatgagtatg atggtaata 100

<210> SEQ ID NO 6
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 6

atgacgtccc gcgacgtccc gcacaacaaa cgcaccggtg cacaaactat agcaagtgtg 60

gtctattttt atatagcaag tgaggcaga taaaataata 100

<210> SEQ ID NO 7
<211> LENGTH: 1350
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 7

atctacaggg cgcgcagcgg ctgacacatg ggacccacgg gcaaaggcag cggcagactc 60

gcgcgcgaga gcacgcaagc accgattgac gcgcgtgccc cttttgtcgg agaacgaggg 120

tgcgcgatcg tccgtgcatg gttgagacgg ccaggcgggg cccacccatt ggcgccatgc 180

cgcgcgagca gcagctggcc cgcacagggt caaagggtga gtgggctaga agcgaagacc 240

tagtccaagc gacggtttct tccttttttc tcttttttcc tttttccttt atattttctt 300

gttttatttt attttcatat ttttttttcc aaatctaaat ttgaactcaa gtgtgagatt 360

-continued

catactttga attaaatgct cacattcaaa tatcagtagg aatagaatat tttaactat	420
atgtttatatt tctctatttt atacaactctt ttcctttttt cattttttaa cctcaatttg	480
taaattatat ctaaattoca attttgggtca ttagtatact cctactaata tcattttatt	540
gttatgaaat gcacacacaa taaactccaa ctgtgatgaat agatatctat ttattcta	600
taattttatt gtttggtaga tgttcaaaat atgaacaca cacatatgtt tttctttctt	660
tttagaaaaa tggatatttt attgtgggac aagaactaaa agtcacctca aaattaaata	720
ctcatgtata atttgggata tctgaattta ctatttttat ttcttcatat atttaattta	780
ttattaattt tttttcttat tgggccttac acttatgaga tcatagaggc atgtggatca	840
gtggcataca agttgaaatt actgccaaaa atgtctgtca tacacaatcc gtgtgtttgg	900
tttgtggacc agcagagcct ggctactcta atccgtatgt agagaaccaa gctcccacga	960
gacagactca atgtatccga gatactcttg tttggttaca cttgcaaaaa tagaccactc	1020
ttgttttggt tcatgtcggg gaaactctaca tggatcagca atgagtatga tggatcaat	1080
ggagaaaaag aaagagtaat taccaatttt ttttcaattc aaaaatgtag atgtccgcag	1140
cgttattata aaatgaaagt acattttgat aaaacgacaa attacgatcc gtcgtattta	1200
taggcgaaag caataaacia attattctaa ttcggaaatc tttatttoga cgtgtctaca	1260
ttcacgtcca aatgggggct tagatgagaa acttcacgat ttggcgcgac taactaagca	1320
ctagcgtacg ggaccagat atcgaattca	1350

<210> SEQ ID NO 8

<211> LENGTH: 1069

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and transgene DNA

<400> SEQUENCE: 8

gatcgccgc acaacaaacg caccggtgca caaactatag caagtgtggt ctatttttat	60
atagcaagtg aggtcagata aaataataaa taaaatcatt aaattttata tagcaagtgt	120
ggctctattt tgcttgtaaa ctacagagaa tatatatatg ttaaatcggg agaacaattg	180
gctattttaa ctaaaagatt taacaaaaac aaagataaaa aatgtttttg aactgcgtgt	240
ctcgagagca gccagcatgc ttccgacctg catgagcagc gatgctcctg gtgaccgcac	300
gaacggaacc tggcccgatg gatgattttg catcaacagc gcctgtatgc aacgatcatc	360
caaccaaaca tacttcgctt tgaggatcca cgcctacaac gtccgcgctc gtccatgcaa	420
ccaaacacac ggaatgtggt ccatgtgtcc caattgaaag aagtgtattc ggtaacaac	480
ggaagtcata accgagccag atatatagat agaaccagat ctatcatacc aagaacaccc	540
ctccaagatt ctagactgca aggaaagatc cactcgtgcg aagacgacca agatgtataa	600
gatccaatgg agcaaccata cggaagaaga ggctacgtgg gagactgagg attatctatg	660
caaatactac ccgattgtgc tacctaagga agtcagtacg taaccatgcc ccagcccct	720
gccctccgat tccaaatata gaaaagatac tcttaatgaa aactgaatta agaaatgaaa	780
ttaagcgaag aaggacttcc ttctgaagtt gcaaagagga tggcattoga agagcagatt	840
tttctcgcga accttaaaaa gctaaccaat actatgcaag caagctataa ccacctctc	900
actctgggt gatctcgact cgaatctcgg ggagagattc ttttaagggg ggagagctgt	960

-continued

aacaccccag gtgttactta gggtttcccc cttagtagctt ccatttgtga cctattatca 1020

catgtgggtt ggtttaagaa aaaggcacca aacttgaggg gccaaagccc 1069

<210> SEQ ID NO 9

<211> LENGTH: 14008

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Transgene insert

<400> SEQUENCE: 9

ttcatgtccg ggaaatctac atggatcagc aatgagtagt atgggtcaata tggagaaaaa 60

gaaagagtaa ttaccaatct tttttcaatt caaaaatgta gatgtccgca gcgttattat 120

aaaatgaaag tacattttga taaaacgaca aattacgacg cgtcgtatct ataggcgaaa 180

gcaataaaca aattattcta attcggaaat ctttatttcg acgtgtctac attcacgtcc 240

aatggggggc ttagatgaga aacttcacga ttggcgcgca ctaactaagc actagcgtac 300

gggacccaga tatcgaattc aagctgaccg ccaccggcaa acaaccacga atttgtaatg 360

gtactaggca aattctccgt ttggcgggtg gtgccggcca attacacgtt ttgctgggtg 420

cctccgacaa aatttgccct ttaaaaacaa ttttataaga gaagctccgg agataaaaag 480

ccgtcaatgt tacaagagtg aagtcgtcta ctccctccat cccaaaaaat gtaattctaa 540

gtatgagttg tattattatt ttggacaaa aggagtatac cacaagaatg atatcatcgt 600

catgcttaga tccttttttag taaagcttga gcttctctaa aagtagagaa attagaaaaa 660

aatcacgttt ttgtggtctt gatttctagc ctccacaaaa tctttggttt tacatttttt 720

gtttgatttt gggttcagaa gtccttattt atatgtgcta gtttggcagc acttaaaatc 780

gttagagaga gcctaaacaa aagccttttc aaaacgacct tgagccagat tggttgatgg 840

ccaaaatttg attgtcaaaa cttaggcaag ccaagatttt agcagctatt tggtttggtg 900

ccaaaatttg ccaatgatct gttcttttgc cttttcaacc ggtttatcag ccgtacttca 960

gcttattctc tctcacagaa cactattgaa tcagccgaaa agccaccgca gaacaggacc 1020

agtatctcac aaatggcatg ccaaatatac tcaccgtcag tgagcccggt taacggcgctc 1080

gacaagtcta acggccacca accagcgaa caccagcgct aagctagcca agcgaagcag 1140

acggccgaga cgttgacacc ttggcgcggg catctctctg gccccctctc gagagtctcg 1200

ctccacctcc actggtggcg gtttccaagt ccgttcggcc tctgtctctc cctcacacgg 1260

cacgaaacgg tcacggcacc ggcagcacgg gggattcctt tcccaccgct ccttcccttt 1320

cccttctctg cccgcggttt taaatagcca gccccatccc cagcttctct ccccaacctc 1380

agcttctctc gttgttcgga gcgcacacac aaccgatccc ccaatccctc cgtctctctc 1440

cgcgagcctc gtcgatcccc gcttcaaggt acggcgatca tctccctttt ctctaccttc 1500

tcttctctag actaggtcgg cgatccatgg ttagggcctg ctagtctctg tctgtttttt 1560

ccgtggctgc gaggtacaat agatctgatg gcgttatgat ggttaacttg tcatactcct 1620

gcggtgtgcg gtcctatagtg cttttaggac atcaatttga cctgggtcgt tcgagatcgg 1680

cgatccatgg ttaggacctt aggcgggtga gtcgggttag atccgcgctg tttgtgttag 1740

tagatggatg cgacctttac ttcagacacg ttctgattgt taacttgta gcacctggga 1800

gtcctgggat ggttctagct ggttcgcaga tgagatcgat ttcagatctt gctgtatctt 1860

-continued

gtttcggttag	gttcctttta	atctatccgt	ggtattatgc	taacctatga	tatgggttcga	1920
tcgtgctagc	tacgtcctgt	gtcataattt	ttagcatgcc	cttttttggt	tggttttgtc	1980
tgattgggct	gtagatcaga	gtatactgtt	tcaaactacc	tactggatat	atttattaaa	2040
tttgaatctg	tatgtgtgtc	acatatatct	tcataattaa	aatggatgga	aagatatatg	2100
gataggatca	tgtgttgctg	tgggttttac	tggtactttg	ttagatatac	atgcttagat	2160
acatgaagca	acatgatggt	acagttcaat	aattcttggt	tacctataaa	acaaataagg	2220
atagggtgat	gttgctgtgg	gttttgctgg	tactttgtta	gatatatatg	cttagatata	2280
tgaagcaaca	tcctgctacg	gtttaataat	tattgtttat	atctaataga	caagcctgct	2340
ttttaattat	tttgatatac	ttggatgatg	gcatacagca	gctatgtgtg	gattttttaa	2400
taccagcat	catgagcatg	catgacctg	ccttagtatg	ctgtttatgt	gcttgagact	2460
tcttttttg	ttggtactca	ccttttgtag	tttggtgact	cttctgcagg	tgcaaccatg	2520
tctccggaga	ggagaccagt	tgagattagg	ccagctacag	cagctgatat	ggcgcgggtt	2580
tgtgatatcg	ttaaccatta	cattgagacg	tctacagtga	actttaggac	agagccacaa	2640
acaccacaag	agtggttgga	tgatctagag	aggttgcaag	atagataccc	ttggttggtt	2700
gctgaggttg	aggggtgtgt	ggctggtatt	gcttacgctg	ggccctggaa	ggctagggaac	2760
gcttacgatt	ggacagtga	gagtactgtt	tacgtgtcac	ataggcatca	aagggtgggc	2820
ctaggatcca	cattgtacac	acatttgctt	aagtctatgg	aggcgcaagg	ttttaagtct	2880
gtggttgctg	ttataggcct	tccaaacgat	ccatctgtta	ggttgcatga	ggctttggga	2940
tacacagccc	gggttacatt	gcgcgcagct	ggatacaagc	atggtggatg	gcatgatggt	3000
ggtttttggc	aaagggattt	tgagttgcca	gctcctccaa	ggccagttag	gccagttacc	3060
cagatctgat	acgcgctgct	agtagccaag	tacctacctc	gagggcacac	tgtaggcagt	3120
gtgccatatt	acagggtcag	attggccggg	acaaagagta	cactgcgatt	ttactatcct	3180
cggtagctg	gtactacagg	ttcacattca	catggttact	acctggcctg	tggtatgctg	3240
gtactacagg	tgttgctctg	aatagctcag	gcttgaccca	agtaagagca	gtgcaaattc	3300
ccaatttcca	gaaaacagga	ggacctccaa	aggacttcaa	tgtaattcaa	atttctttca	3360
gaggtccttt	tgcaaggggc	tgatgtataa	gtacttttat	aataatccaa	gccattgtgc	3420
ttctcaaaaa	aaggaaaaga	aaaaaatttg	ttgcgggccc	gccgtgacgg	ccacgagcga	3480
actcctgcag	gagcagactc	gcattatcga	tgagctctca	ccaaactggc	cctaggcatt	3540
aacctaccat	ggatcacatc	gtaaaaaaa	aacctacca	tgatcctat	ctgttttctt	3600
tttgccctga	aagagtgaag	tcatcatcat	atttaccatg	gcgcgcgtag	gagcgcttcg	3660
tcgaagagcc	ataggggggc	ggtagctgca	ccgtgggtgt	ttcctgttat	gtaatatcgg	3720
atgggggagc	agtcggctag	gttggtccca	tcggtagctg	tcgtccctca	gtgcgctaga	3780
tgcgcatgtg	ttgtcctcaa	aaactctttt	cttcttaata	acaatcatat	gcaaattttt	3840
tgcgatttcg	agaaaaaag	aagattctat	ctgttttttt	tttgaaatgg	ctccaattta	3900
taggaggagc	ccgtttaacg	gcgtcgacaa	atctaacgga	caccaaccag	cgaatgagcg	3960
aaccaccag	cgccaagcta	gccaaagcga	gcagacggcc	gagacgctga	cacccttgcc	4020
ttggcgcggc	atctccgctg	ctggctcgtc	ggctctggcc	ccttcgcgag	agttccggtc	4080
cacctccacc	tgtgtcggtt	tccaaactcg	ttccgccttc	gcgtgggact	tgttcggttc	4140

-continued

atccgttggc	ggcaccgga	aattgcgtgg	cgtagagcac	ggggccctcc	tctcacacgg	4200
cacggaacgg	tcacgagctc	acggcaccgg	cagcacggcg	gggattcctt	ccccaccacc	4260
gtctcttccc	tttcccttcc	tcgcccgcga	tcataaatag	ccacccctcc	cagcttccctt	4320
cgccacatcc	tctcatcctc	ttctctcgtg	tagcacgcgc	agcccgatcc	ccaatccctt	4380
ctcctcgcga	gcctcgtcga	tccctcgttt	caaggtatgg	ctatcgtcct	tcctctctct	4440
ctctttacct	tatctagatc	ggcgatccat	ggttagggcc	tgctagtctt	ccgttcgtgt	4500
ttgtcgatgg	ctgtgaggca	caatagatcc	gtcggcggtta	tgatgggttag	cctgtcatgc	4560
tcttgcgac	tgtggttcct	ttaggaaagg	cattaattta	atccctgatg	gttcgagatc	4620
ggatgatccat	ggttagtagc	ctaagctgtg	gagtcggggt	tagatccgcg	ctgttcgtag	4680
gcgatctgtt	ctgattgtta	actgtcaggt	acctgcgaat	cctcgggtgg	tctagctggt	4740
tcggagatca	gatcgattcc	attatctgct	atacatcttg	tttcggttgc	taggctccgt	4800
ttaatctatc	catcgtagta	tgttagcctt	tgatatgatt	cgatcgtgct	agctatgtcc	4860
tgtggactta	attgtcaggt	cctaattttt	aggaagactg	ttccaaacca	tctgctggat	4920
ttattaaatt	tggatctgga	tgtgtcacat	acaccttcat	aattaaaatg	gatggaaata	4980
tctcttatct	tttagatatg	gataggcatt	tatatgatgc	tgtgagtttt	actagtactt	5040
tcttagaata	tatgtacttt	tttagacgga	atattgatat	gtatacatgt	gtagatacat	5100
gaagcaacat	gctgctgtag	tctaataatt	cctgttcctc	taataatcaa	gtatgtatat	5160
gttctgtgtg	ttttattggg	atttgattag	atatatacat	gcttagatac	atacatgaag	5220
cagcatgctg	ctacagttta	atcattattg	tttatccaat	aaacaaacat	gctttttaat	5280
ttatcttgat	atgcttggtg	gacggaatat	gcagagattt	taagtaccca	gcatcatgag	5340
catgcatgac	cctgcgtag	tatgctgttt	atttgcttga	gactctttct	ttttagatata	5400
ctcacccctg	tttctggtga	tctactgca	ggtgaccggg	cgccatggcc	accgccacca	5460
ctaccgccac	cgtcgcgttc	tccggcgtgg	tgagcgtcgg	cactgagacg	cgcaggatct	5520
actccttcag	ccacctccag	ccttctgctg	cgttccccgc	taagccgtct	tcgttcaaga	5580
gcctgaagct	gaaacagtcc	gcacgcctta	cccggcgcc	ggaccatagg	ccattcggtg	5640
tcagggtgat	gctcaccttc	gttaggaacg	cctggtacgt	cgcgcgtctc	cctgaggagc	5700
tgagcgagaa	gcccttgggg	cgcacccatc	tagacactcc	gttagccctt	taccgccagc	5760
ctgacggcgt	agtggcgggc	ctgcttgaca	tctgcccga	taggttcgct	ccgctcagcg	5820
acggcatcct	cgtcaacggg	catcttcagt	gcccgtacca	cgggctggaa	tttgacggcg	5880
gtgggcagtg	tgtccacaac	cgcacgggca	acggcgacgc	gccagcttcc	ctcaacgtta	5940
ggctcgttccc	tgtgtctgag	cgcgacgcac	tgatctggat	ctggcctggc	gaccagctc	6000
tggccgatcc	aggagccatt	cccgaacttc	gttgccgcgt	ggaccagacc	tatcggaagg	6060
tcggcgggta	cgggcacgtc	gattgttaact	ataagctcct	tgtggacaac	cttatggatt	6120
tgggccacgc	tcagtacgtg	caccgggcta	acgctcagac	tgacgccttt	gacgctctcg	6180
aaaggggagg	catcgctggc	gacggagaga	ttcaggcgct	gatgaagatc	cctggaggga	6240
cgcctctctg	gctcatggcg	aagttttctc	gaggcgcgaa	cacgcccgtg	gacgctgga	6300
acgacatccg	ctggaataag	gtctccgcga	tgtgaaactt	catcgccgtt	gcgcccagg	6360
gcacacccaa	agagcagtca	atccacagca	gagggaacca	tattcttaca	ccggaaacgg	6420

-continued

aggctagttg ccactacttc ttcggctcgt cacggaattt cgggatatagac gatccggaga	6480
tggacggtgt tcttcgatct tggcaagcgc aagctctcgt caaggaagat aaggtggtcg	6540
tggaggctat cgagcgtagg cgcgcctacg ttgaggcgaa cgggtattagg cccgcgatgc	6600
tgtcctgcga cgaggccgca gttagagtgt cgcgcgagat agaaaagctg gagcagctag	6660
aggccgcctg attaatatag gccaaaggca tctatgactg aattgccaat gcaccagcct	6720
gtctacatga tgaataaata aagagtccat ccagtgtgat ggctcatgcc tgtgtgagt	6780
tgactgaatc catcagtgtg tgtgtgtgtt tgtgtcaacc atgtgtgaat cagggtgtcaa	6840
aaatcgtggc tggaaatcca tgtggtttct agctttatgt aaatgttgtt tgtgaaatat	6900
aaatatgtt ttgtgtatgt gaattttact ctctcatttt tctcttgca caccattct	6960
attatagtaa tttttttaag cggccgctga tgtatccgtc cagggtggtg tgtccaatca	7020
gtgaataatc tagttagtga agccagaagt ccatagtgcc ccttgctctg tcaccatata	7080
tccagttcaa ccgcaccaat ttgccatctc gaactgggtc atgttttatt cagggtggta	7140
aatgaatttt gccaatccaa tgtagttaga tatttccatg tcatttttagt acattttacca	7200
attttttata ttctggctag aaaaggagaa tggtgacgtc ttctggaaga tcaagatcaa	7260
ttatcaagta tcagcaacag cacctgaagg ttggagtga ttagttgtca ttgagaataa	7320
tgctagctat tcattgcact ggcattagag acagagaggg cgagccagtt tgacatggca	7380
aattagcaca gtcaaatcgt atacgtggtg acggaggagg gggcactatg aatttttgg	7440
gacggaggga ggggcactat gaatttttgg ctttgctgac gggcacgccc actatggatg	7500
aaattggaca aaatacgaat attcaaggat gaaagtggtc ggtttgatag ttcagggatg	7560
aaatgtgtct ttgggcaaac tttagaggac aagttgccta ttttgatta aacgaatata	7620
tttatatacc ccaaaaaaaa gaatacacat ctccactccg agccggcatg tgggggtccc	7680
actagtcagc cactgtatgg cgcgcactag ctcaacggcc acgaaccagc caaccaccag	7740
cgcaacctaa acggcgtaaa cgttgacggc atctctctct cgcctcctct cgaagcttcc	7800
gcaccgctcg ctggtcgtcg cccggcgccg ctctgtctgg actctttccg tggcggtctc	7860
cgcgaaattg cgtggtggag aggagagacg gaaccgtcac ggcactggat tccttcccca	7920
cccggcttgg ccggccctc ctgcctcca taaataggca cccgctctc gcctcctctc	7980
cccacctcat ctctccttt cccgtgaacc gtgaacacaa cccgaccag atccctctt	8040
gcgagcttgg tcgatccctc ctccgctga aggtacggag cttctcctcc cccttctct	8100
ctagatcggc gtgttatgtt gtttccgtgg ttgcttggtt ggatgaatcg aatgattctt	8160
agggcctagg aggtctgtta gatctgttgc gttctgttcc gtagatggat ttgggtgtaa	8220
gatcaggctg gttccgctgt ttaacttgt atgctagtgt gatttttggg aggtattgag	8280
ttgttaatct gggagtgtt gggaggttct cgtaggcgga ttgtagatga agtcgcccgc	8340
acgatttgcg tggttgttg ggtagctagg gttagatctg ctcggtttt tcattgttac	8400
ttattgagag ataatgtagc taacctttac ttgttcatct atgtatctcg tattcgtatt	8460
catctggttc gatggtgcta gatagatcg cctgatttgt ccgatcgaat tgggtagcat	8520
ccgcggttg tttggtagt ttctgattga tttgtcgtc tagatctgag tggaataata	8580
ttacatctca acatgttact agaaacttgg tttatagctc cggattttaca tgtttattct	8640
tatgtaaggt tttaaatgaa agatttatgc tactgtgtct cgttgatcct ttagcatcca	8700

-continued

cctgaggaac atgcatgcat ctgttacttc ttttgatata tgcttagata gttgttagta	8760
tatactgctg ttgttcgatg atccttcagg atgaacatgc atgatcatgt tacttgtttt	8820
tatatgcttc tgctgttcgt tgattcttta gtactaccta cctgatcatc ttgcatgttt	8880
cctgcttggt agagattaat tgattaggct taccttggtg cctggtgatt ctcccttgca	8940
ggtaggtacc cggaccgatg gcaacagcaa catcagcttc tctgttttca actgtttctt	9000
catcttactc caaagctagc tccataccac attcaagact ccaatctgtg aaattcaact	9060
cagtccttag ctccaccggt ctcaaataca cctctctcat ctccggatct gattcctctt	9120
ccttagccaa gactctacgc ggttccgtaa cgaaagcaca aacatctgac aagaagcctt	9180
acggattcaa aatcaacgct atgcacgcgg cgctgactcc tctcaccaac aagtatcgct	9240
ttatcgacgt gcagccgctg acaggcgctc tcgggtgcaga gattacaggc gtggatctgc	9300
gggagcctct cgatgacagc acttggaatg agatcctgga cgcctttcac acctaccaag	9360
tgatctactt tccgggtcaa gctatcacta acgagcagca catcgcgctc tcccgccggt	9420
tcggccctgt ggacccggtg ccgatcttaa agagtatcga gggctatcca gaggtgcaga	9480
tgatacggcg cgaggcgaac gagagcagcc ggttcatcgg agatgactgg cacaccgatt	9540
ccaccttctt ggacgctccg cctgcccgcg tggatgatgag agctatcgaa gtgccggagt	9600
atggaggatg cacaggcttc ctctccatgt acagtgcctg ggagacactc tcgcctacga	9660
tgcaagctac catcgaaggc ttaaacgtgg tccactcggc gacgaaggc ttcgggtcat	9720
tgtaccaggc gactaattgg cgcttctcga acaccagcgt gaaagtgatg gacgtggacg	9780
ccggagatag agagactgtg caccactcgc tcgtgacgca tcctgttacg ggaaggcgcg	9840
cactctactg caaccagggt tactgccaga agatccaggg aatgacggac gcggagtcca	9900
agtcctgtt gcaattcctt tacgagcacg ccaccaagtt cgacttcacc tgccgggtcc	9960
ggtaggaaga ggaccaagtc ctgggtgtggg acaacctgtg taccatgcac cgcgcgctcc	10020
cggactacgc tgggaaattc agatacctga cccgcaccac cgtggcggga gacaagccgt	10080
cgcgttgacg gtcggttaat taatcgagtg tgaagggaagt gaactgagtc tcaagaataa	10140
aaactgcaag tatggcgctc cattccatgg aagacctgca tgcactctat atatgtgctt	10200
ttatgttcaa gttggagaac tatgtgtgtt gtctttaagt tggagtactc taagctattc	10260
gatctggaga ttatgctgtt ttaataaaaa gtttgaagtg tatctgtcct gtttaattag	10320
tttcaagaaa gccaaagcgt tttgggttct aggatgaaca aagtcgtcca agatgcaaga	10380
agttctagag aggtagctac atttctggaa attgttgcac ctcccttctt tcaatatata	10440
tatattgtcc tatttcataa tgtcattcta gccagctatc tatattctat atatccattt	10500
ccgttgccat actttattca ctatgttgca tcgagtgagc tgcattgctc tgtactacat	10560
gttacttgat atgtgttggt cataaacaca cacattaata atgatcagat tgtgaaaaat	10620
acgcgtccga tcctacctgt cacttcatca aaaggacagt agaaaaggaa ggtggcacct	10680
acaaatgcca tcattgcgat aaaggaaagg ctatcattca agatgcctct gccgacagtg	10740
gtcccaaaaga tggaccccca cccacgagga gcatcggtga aaaagaagac gttccaacca	10800
cgtcttcaaa gcaagtggat tgatgtgata ctccactga cgtaagggat gacgcacaa	10860
cccactatcc ttcgcaagac ccttctctta tataaggaag ttcatttcat ttggagagga	10920
cacgctgaaa tcaccagtct ctctctacaa gatcggggat ctctagccct agaaccatct	10980

-continued

tccacacact	caagccacac	tattggagaa	cacacagga	caacacacca	taagatccaa	11040
gggaggcctc	cgccgcgcgc	ggtaccacc	ccgcccctct	cctctttctt	tctccgtttt	11100
tttttccgtc	tccgtctcga	tctttggcct	tggtagtgtt	ggtgggcgag	aggcggtctc	11160
gtgcgcgcgc	agatcgtg	gcgggagggg	cgggatctcg	cggctggggc	tctgcgcgcg	11220
gtggatccgg	cccggatctc	gcggggaatg	gggctctcgg	atgtagatct	gcgatccgcg	11280
gttgttgggg	gagatgatgg	gggggtttaa	atttcgcgcg	tgctaaacaa	gatcagggaag	11340
aggggaaaag	ggcactatgg	tttatatttt	tatatatttc	tgtgcttcg	tcaggcttag	11400
atgtgctaga	tctttctttc	ttctttttgt	gggtagaatt	tgaatccctc	agcattgttc	11460
atcggtagtt	tttcttttca	tgatttgtga	caaatgcagc	ctcgtgcgga	gcttttttgt	11520
aggtagaagt	gatcaaccat	ggcgcaagtt	agcagaatct	gcaatggtgt	gcagaaccca	11580
tctcttatct	ccaatctctc	gaaatccagt	caacgcaaat	ctcccttctc	ggtttctctg	11640
aagacgcagc	agcatccacg	agcttatccg	atttcgctcg	cgtggggatt	gaagaagagt	11700
gggatgacgt	taattggctc	tgagcttcgt	cctcttaagg	tcatgtcttc	tgtttccacg	11760
gcgtgcatgc	ttcatggagc	ttcatctagg	ccagctactg	ccaggaagtc	tagcgggctc	11820
agtggcaccg	tgcgcatccc	tgccgataaa	agtatttcac	acaggagctt	catgttcgga	11880
ggacttgcta	gtggagagac	gagaatcact	ggtttgcttg	agggcgaaga	tgttatcaac	11940
accggtaaag	cgatgcaagc	aatgggtgcc	agaatccgaa	aagagggcga	tacgtggatc	12000
atcgacgggtg	ttggtaacgg	aggattgtct	gctcccgaag	cgccacttga	ctttgggaac	12060
gcagctacgg	gggtccgtct	tactatggga	ctggtagggc	tgtatgactt	tgactctacc	12120
ttcatcgggtg	acgcgagcct	cactaagaga	ccaatgggac	gagtgtgaa	tcccctgagg	12180
gagatgggtg	tccagggtga	atctgaggat	ggtgatcgct	ttccggttac	tctgcgaggc	12240
cccaagaccc	ccacgccaat	cacgtacagg	gttccgatgg	cgtcagcaca	ggtcaagtca	12300
gcggtactcc	tgccgggcct	caacacacct	ggaatcacia	ccgtgattga	acccatcatg	12360
actagagacc	acacggagaa	gatgttgacg	ggtttcgcg	ctaactctaac	ggtcgaaacc	12420
gacgccgacg	gcgtgaggac	aatccgcttg	gagggcagag	gtaaaactgac	tgcccaagtc	12480
atcgatgtgc	ctggagatcc	ctcgtccaca	gcgtttcccc	tcgtagctgc	gttgctcgtc	12540
cctggatctg	atgtgacgat	cctgaatgtc	ctcatgaatc	caactagaac	cggcctcatc	12600
ctcacattgc	aggagatggg	tgtgacatc	gaggttatca	atcctagggt	ggcagggtgga	12660
gaggatgtgg	ccgatctg	cgtgcgttct	agtacactca	aaggcgtgac	cgtccctgag	12720
gatcgcgctc	catccatgat	cgacgagtac	cccattctcg	ccgttgctgc	tcggtttgcc	12780
gagggcgcaa	ctgtaatgaa	cgcccttgag	gagttgaggg	ttaaggagag	tgacaggctg	12840
tccgcggtgg	cgaatggcct	gaagctaaac	ggcgtggact	gcgacgaagg	tgaaacgtcc	12900
cttgtagtcc	gtggtcgccc	agacgggaag	gggttgggga	atgcttcggg	agctgctgtg	12960
gcgacgcacc	ttgatcatag	aatcgccatg	tcatttctgg	tgatgggact	tgtctccgag	13020
aatccggtga	ccgttgacga	tgctaccatg	atcgccacct	cctttcctga	gttcatggac	13080
ctcatggcag	gcttgggggc	caagatcgag	ctgtctgata	ctaaggccgc	ttgacacgtg	13140
agacgacgac	cgtgcatggt	gatgaatcac	agacgacgaa	cacgcagtgc	gcacgcgcgt	13200
cctctttgta	tcacacagca	tcaccttctt	cgctctacta	ctcccccgag	caatcaccca	13260

-continued

```

ccaataacac caaccatcaa cctccccgt cgccgcgcgc ttcaccgtcc tcccctcaca 13320
ccatagaact gcaaatgtcc gctgcaggg tggggcccat cgtggccagt tatecttagc 13380
tatccgtgtc agaatcatct tatcatcgag tcgagtcggt atcgtgtcca gtggctctct 13440
cgagtcgaga agccctctat ccatecatcc agtgttagggt gttcttcgtc cgtgatgtta 13500
ccatgaattg agttcgcttt ggttatggtg tttgaactgc ttgttgctat ctatcggaat 13560
gaaatgaaat agaagacaag gagaaaaaaa agagttcgaa agttttgttc gcataccata 13620
tatttccttc cggtgcgcgc tgtttattcc tcgctcagca gcaagattgt ttgatcgata 13680
ttgcagcaag caattacaca ataaatatat tgctacactg gtacttcaaa ctacactggt 13740
ggtcggtgat tttcaatagc atgaacctta attgaacatc tgtgtagctt acatctcctt 13800
cgaaagctgc aatgcttgag aacttggaag gaaattcttg tgatggcaga agctattcac 13860
tgtccttcgc tgcatttaca gtccatacag acacagcatt tccattttgc acaagataga 13920
gaacaacaat cagcctttta ggtcaatccc aagtgtgcat gacgtccgc gatcgccgc 13980
acaacaaacg caccggtgca caaactat 14008

```

```

<210> SEQ ID NO 10
<211> LENGTH: 16068
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of maize genomic DNA and
transgene DNA

```

```

<400> SEQUENCE: 10

```

```

atctacaggg cgcgagcgg ctgacacatg ggacccacgg gcaaaggcag cggcagactc 60
gcgcgcgaga gcacgcaagc accgattgac gcgcgtgccc cttttgtcgg agaacgaggg 120
tgcgcgatcg tccgtgcatg gttgagacgg ccaggcgggg cccaccatt ggcgccatgc 180
cgcgcgagca gcagctggcc cgcacagggtg caaagggtga gtgggctaga agcgaagacc 240
tagtccaagc gacggtttct tcttttttct tcttttttct ttttctctt atattttctt 300
gttttatttt attttcatct ttttatttct aaatctaaat ttgaactcaa gtgtgagatt 360
catactttga attaatgct cacattcaaa tatcagtagg aatagaatat ttttaactat 420
atgtttattt tctctatttt atacaatctt ttcctttttt cattttttaa cctcaatttg 480
taaatatat ctaaatcca attttggtca ttagtatact cctactaata tcattttatt 540
gttatgaaat gcacacacaa taaactccaa cttgatgaat agatatctat ttattctaatt 600
taattttatt gtttggtaga tgttcaaat atgaaacaca cacatattgt tttctttctt 660
tttagaaaaa tggatttttt attgtgggac aagaactaaa agtcacctca aaattaaata 720
ctcatgtata atttgggata tctgaattta ctatttttat ttcttcatat atttaattta 780
ttattaattt ttttcttat tgggccttac acttatgaga tcatagaggc atgtggatca 840
gtggcataca agttgaaatt actgccaaaa atgtctgtca tacacaatcc gtgtgtttgg 900
tttgtggacc agcagagcct ggctactcta atccgtatgt agagaaccaa gctcccacga 960
gacagactca atgtatccga gatactcttg tttggttaca cttgcaaaaa tagaccactc 1020
ttgtttggtt tcatgtccgg gaaatctaca tggatcagca atgagtatga tggatcaatat 1080
ggagaaaaag aaagagtaat taccaatttt ttttcaattc aaaaatgtag atgtccgcag 1140

```


-continued

cgttattata	aatgaaagt	acattttgat	aaaacgacaa	attacgatcc	gtcgtattta	1200
taggcgaaag	caataaacaa	attattctaa	ttcggaaatc	tttatttcga	cgtgtctaca	1260
ttcacgtcca	aatgggggct	tagatgagaa	acttcacgat	ttggcgcgac	taactaagca	1320
ctagcgtacg	ggaccagat	atcgaattca	agctgaccgc	caccggcaaa	caaccacgaa	1380
tttgtaatgg	tactaggcaa	attctccgtt	tggcgggtgtg	tgccggccaa	ttacacgttt	1440
ttgcgggtgc	ctccgacaaa	atttgccctt	taaaaacaat	tttataagag	aagctccgga	1500
gataaaaggc	cgtcaatgtt	acaagagtga	agtcgtctac	tccctccatc	ccaaaaaatg	1560
taattctaa	tatgagtgtt	attattattt	ttggacaaaa	ggagtatacc	acaagaatga	1620
tatcatcgtc	atgcttagat	cctttttagt	aaagcttgag	cttctctaaa	agtagagaaa	1680
ttagaaaaaa	atcacgtttt	tgtggtcttg	atttctagcc	tccacaaaat	ctttggtttt	1740
acattttttg	tttgattttg	gtttcagaag	tccttattta	tatgtgctag	ttgggcagca	1800
cttaaaatcg	ttagagagag	cctaaacaaa	agccttttca	aaacgacctt	gagccagatt	1860
ggttgatggc	caaaatttga	ttgtcaaaac	ttaggcaagc	caagatttta	gcagctattt	1920
ggtttggtac	caaaatttgc	caatgatctg	ttcttttgcc	ttttcaaccg	gtttatcagc	1980
cgtacttcag	cttattctct	ctcacagAAC	actattgaat	cagccgaaaa	gccaccgcag	2040
aacaggacca	gtatctcaca	aatggcatgc	caaataact	caccgtcagt	gagcccgctt	2100
aacggcgctc	acaagtctaa	cggccaccaa	ccagcgaacc	accagcgta	agctagccaa	2160
gcgaagcaga	cggccgagac	gttgacacct	tggcgcgggc	atctctctgg	ccccctctcg	2220
agagtccgc	tccacctcca	ctggtggcgg	tttccaagtc	cgttcgcct	cctgctctc	2280
ctcacacggc	acgaaacgt	cacggcaccc	gcagcacggg	ggattccctt	cccaccgctc	2340
cttcccttcc	ccttccctgc	ccgccgtttt	aaatagccag	ccccatcccc	agcttctctc	2400
cccaacctca	gcttctctcg	ttgttcggag	cgcacacaca	accgatccc	caatccctc	2460
gtctctctc	gcgagcctcg	tcgatcccc	cttcaaggta	cggcgatcat	cctcccttcc	2520
tctaccttct	cttctctaga	ctaggctggc	gatccatggt	tagggcctgc	tagttctggt	2580
cctgttttcc	cgtggctgcg	aggtacaata	gatctgatgg	cgttatgatg	gttaacttgt	2640
catactctcg	cgggtgtcgg	tctatagtgc	ttttaggaca	tcaatttgac	ctggctcggt	2700
cgagatcggc	gatccatggt	taggacctta	ggcgggtggg	tcggggtaga	tccgcgctgt	2760
ttgtgttagt	agatggatgc	gacctttact	tcagacacgt	tctgattggt	aacttgctag	2820
cacctgggag	tcctgggatg	gttctagctg	gttcgcagat	gagatcgatt	tcagtatctg	2880
ctgtatcttg	tttcgttagg	ttccttttaa	tctatccgtg	gtattatgct	aacctatgat	2940
atggttcgat	cgtgctagct	acgtcctgtg	tcataatttt	tagcatgccc	tttttggttt	3000
ggttttgtct	gattgggctg	tagatcagag	tatactgttt	caaactacct	actggatata	3060
tttattaaat	ttgaatctgt	atgtgtgtca	catatatctt	cataattaaa	atggatggaa	3120
agatatatgg	ataggtacat	gtgttgctgt	gggttttact	ggtaacttgt	tagatatata	3180
tgcttagata	catgaagcaa	catgatgtta	cagttcaata	attcttggtt	acctaataaa	3240
caaataagga	taggtgtatg	ttgctgtggg	ttttgctggg	actttggttag	atatatatgc	3300
ttagatatat	gaagcaacat	cctgctacgg	tttaataatt	attgtttata	tctaataagac	3360
aagcctgctt	tttaattatt	ttgatatact	tggatgatgg	catacagcag	ctatgtgtgg	3420

-continued

atttttaaat	accagcatc	atgagcatgc	atgaccctgc	cttagtatgc	tgtttatttg	3480
cttgagactt	ctttttttgt	tggactcac	ctttttagt	ttggtgactc	ttctgcaggt	3540
gcaaccatgt	ctccggagag	gagaccagtt	gagattaggc	cagctacagc	agctgatatg	3600
gccgcggttt	gtgatatcgt	taaccattac	attgagacgt	ctacagtga	ctttaggaca	3660
gagccacaaa	caccacaaga	gtggattgat	gatctagaga	ggttgcaaga	tagataccct	3720
tgggtggttg	ctgaggttga	gggtgttg	gctggtattg	cttacgctgg	gccctggaag	3780
gctaggaacg	cttacgattg	gacagttgag	agtactgttt	acgtgtcaca	taggcacaa	3840
agggtgggcc	taggatccac	attgtacaca	catttgctta	agtcctatga	ggcgcaaggt	3900
tttaagtctg	tggttgctgt	tataggcctt	ccaaacgac	catctgttag	gttgcagag	3960
gctttgggat	acacagcccg	gggtacattg	cgcgcagctg	gatacaagca	tgggtgatgg	4020
catgatgttg	gttttttgca	aagggatttt	gagttgccag	ctcctccaag	gccagttagg	4080
ccagttaccc	agatctgata	cgcgtgcta	gtagccaagt	acctacctcg	agggcacact	4140
gtaggcagtg	tgccatatta	caggttcaga	ttggccggga	caaagagtac	actgcgattt	4200
tactatcctc	ggtagctg	tactacaggt	tcacattcac	atggttacta	cctggcctgt	4260
ggtagctg	tactacaggt	gttgcctga	atagctcagg	cttgacccaa	gtaagagcag	4320
tgcaaattcc	caatttccag	aaaacaggag	gacctccaaa	ggacttcaat	gtaattcaaa	4380
tttctttcag	aggtcctttt	gcaaggggct	ggatgtaaa	tacttttata	ataatccaag	4440
ccattgtgct	tctcaaaaaa	aggaaaagaa	aaaaatttgt	tgccggcccg	ccgtgacggc	4500
cacgagcgaa	ctcctgcagg	agcagactcg	cattatcgat	ggagctctac	caaactggcc	4560
ctaggcatta	acctaccatg	gatcacatcg	taaaaaaaaa	acctaccat	ggatcctatc	4620
tgttttcttt	ttgccctgaa	agagtgaagt	catcatcata	tttaccatgg	cgcgcgtagg	4680
agcgttctgt	cgaagaccca	tagggggggc	gtactcgac	cgtggttgtt	tcctgttatg	4740
taatatcgga	tgggggagca	gtcggctagg	ttggtcccat	cgttactggt	cgtcccttag	4800
tgcgctagat	gcgcgatgtt	tgtcctcaaa	aactcttttc	ttcttaataa	caatcatacg	4860
caaatttttt	gcgtattoga	gaaaaaaga	agattctatc	tgtttttttt	ttgaaatggc	4920
tccaatttat	aggaggagcc	cgtttaacgg	cgtcgacaaa	tctaacggac	accaaccagc	4980
gaatgagcga	accaccacgc	gccaaagctag	ccaagcgaag	cagacggccg	agacgctgac	5040
acccttgcc	tggcgcggca	tctccgtcgc	tggctcgtg	gctctggccc	cttcgcgaga	5100
gttcgggtcc	acctccacct	gtgtcgggtt	ccaactccgt	tcgccttcg	cgtgggactt	5160
gttcggttca	tccgttggtg	gcatccggaa	attgcgtggc	gtagagcacg	gggccctcct	5220
ctcacacggc	acggaaccgt	cacgagctca	cggcaccggc	agcacggcgg	ggattccttc	5280
cccaccacgc	ctccttcctt	ttccttcctt	cgcgcgccat	cataaatagc	caccctcccc	5340
agcttctctc	gccacatcct	ctcatcatct	tctctcgtgt	agcacgcgca	gcccgatccc	5400
caatccccctc	tcctcgcgag	cctcgtcgat	ccctcgtctc	aaggtatggc	tatcgtcctt	5460
cctctctctc	tctttacctt	atctagatcg	gcgatccatg	gttagggcct	gctagttctc	5520
cgttcgtgtt	tgtcgatggc	tgtgaggcac	aatagatccg	tcggcggttat	gatggttagc	5580
ctgtcatgct	cttcgatctc	gtgggttcctt	taggaaaggc	attaatttaa	tcctgatgg	5640
ttcgagatcg	gtgatccatg	gttagtacct	taagctgtgg	agtcgggttt	agatccgcgc	5700

-continued

tgttcgtagg	cgatctgttc	tgattgttaa	cttgtcagta	cctgcgaatc	ctcggtggtt	5760
ctagctgggt	cggagatcag	atcgattcca	ttatctgcta	tacatcttgt	ttcgttgcc	5820
aggctccgtt	taatctatcc	atcgatgat	gttagccttt	gatatgatc	gatcgtgcta	5880
gctatgtcct	gtggacttaa	ttgtcaggtc	ctaattttta	ggaagactgt	tccaaaccat	5940
ctgctggatt	tattaaat	ggatctggat	gtgtcacata	caccttcata	attaaaatgg	6000
atggaaatat	ctcttatctt	ttagatatgg	ataggcattt	atatgatgct	gtgagtttta	6060
ctagtacttt	cttagaatat	atgtactttt	ttagacggaa	tattgatatg	tatacatgtg	6120
tagatacatg	aagcaacatg	ctgctgtagt	ctaataatc	ctgttcac	aataatcaag	6180
tatgtatatg	ttctgtgtgt	tttattggta	tttgattaga	tatatacatg	cttagataca	6240
tacatgaagc	agcatgctgc	tacagtttaa	tcattattgt	ttatccaata	aacaaacatg	6300
ctttttaatt	tatcttgata	tgcttgtagt	acggaatatg	cagagatttt	aagtaccag	6360
catcatgagc	atgcgatgac	ctgcgttagt	atgctgttta	tttgcttgag	actctttctt	6420
ttgtagatac	tcaccctgtt	ttctggtgat	cctactgcag	gtgaccggtc	gccatggcca	6480
cgcgccacc	taccgccacc	gctgcgttct	cggcggtggt	gagcgtcggc	actgagacgc	6540
gcaggatcta	ctccttcagc	cacctccagc	cttctgctgc	gttccccgct	aagccgtctt	6600
cgttcaagag	cctgaagctg	aaacagtccg	cacgccttac	cggcgccctg	gaccataggc	6660
cattcgttgt	cagggtgatg	ctcaccttcg	ttaggaacgc	ctggtacgtc	gccgctctcc	6720
ctgaggagct	gagcgagaag	cccttgggtg	gcaccatcct	agacactccg	ttagcccttt	6780
accgccagcc	tgacggcgta	gtggcgggcc	tgcttgacat	ctgcccgcat	aggttcgctc	6840
cgtcagcga	cggcatcctc	gtcaacgggc	atcttcagtg	cccgtagcac	gggctggaat	6900
ttgacggcgg	tgggcagtgt	gtccacaacc	cgcacggcaa	cggcgacagg	ccagcttccc	6960
tcaacgttag	gtcgttcctt	gttgctgagc	gcgacgcact	gatctggatc	tggcctggcg	7020
accagctctc	ggccgatcca	ggagccatc	cgaacttcgg	tgccgcgctg	gaccagcct	7080
atcggacggg	cggcggttac	gggcacgtcg	attgttaacta	taagctcctt	gtggacaacc	7140
ttatggattt	gggccacgct	cagtacgtgc	accgggctaa	cgctcagact	gacgcctttg	7200
accgtctcga	aagggaaggc	atcgctggcg	acggagagat	tcaggcgctg	atgaagatcc	7260
ctggaggcac	gccctctgtg	ctcatggcga	agtctctcag	aggcgcgaa	acgccgtgg	7320
acgcctggaa	cgacatccgc	tggaataagg	tctccgcgat	gctgaacttc	atcgccgttg	7380
cgcgcgaggg	cacacccaaa	gagcagtc	tcacacagc	agggaacct	attcttacac	7440
cggaaaccca	ggctagtgtc	caactattct	tcggctcgtc	acggaatttc	gggatagacg	7500
atccggagat	ggacgggtgt	cttcgatctt	ggcaagcgca	agctctcgtc	aaggaaagata	7560
agggtggtcgt	ggaggctatc	gagcgtaggc	gcgcctacgt	tgaggcgaa	ggtattaggc	7620
cgcgatgct	gtcctgcgac	gaggccgag	ttagagtgtc	gcgcgagata	gaaaagctgg	7680
agcagctaga	ggccgcctga	ttaattaagg	ccaaggcgat	ctatgactga	attgccaatg	7740
caccagcctg	tctacatgat	gaataaataa	agagtccatc	cagtgtgatg	gctcatgcct	7800
gtgtgagtgt	gactgaatcc	atcagtgtgt	gtgtgtgttt	gtgtcaacca	tgtgtgaatc	7860
agggtgtcaaa	aatcggtggc	ggaaatccat	gtggtttcta	gctttatgta	aatgtgtgtt	7920
gtgaaatata	aatattgttt	tgtgtatgtg	aattttactc	tctcattttt	ctcttgcaat	7980

-continued

caccattcta	ttatagtaat	ttttttaage	ggccgctgat	gtatccgtcc	acggtggtgt	8040
gtccaatcag	tgaataatct	agtttagtgaa	gccagaagtc	catagtgcc	cttgctctgt	8100
caccatatat	ccagttcaac	cgcaccaatt	tgccatctcg	aactgggtca	tgttttatct	8160
aggttggtaa	atgaattttg	ccaattcaat	gtagtttagat	atttccatgt	catttttagta	8220
cattttacc	aa	ttttttatat	tctggctaga	aaaggagaat	ggtgacgtct	8280
caagatcaat	tatcaagtat	cagcaacagc	acctgaaggt	tggagtgc	at	8340
tgagaataat	gctagctatt	cattgcactg	gcattagaga	cagagagggc	gagccagttt	8400
gacatggcaa	attagcacag	tcaaaactgga	tacgtgggta	cggagggagg	ggcactatga	8460
atttttggtg	acggagggag	gggcactatg	aatttttggc	tttgctgacg	ggacacgcca	8520
ctatggatga	aattggacaa	aatacgaata	ttcaaggatg	aaagtggctg	gtttgatagt	8580
tcagggatga	aatgtgtctt	tgggcaaaact	ttgaggacga	agttgcctat	ttgcattaa	8640
acgaatatat	ttatataccc	caaaaaaag	aatacacatc	tccactccga	gccggcatgt	8700
ggggtcccca	ctagtcagcc	actgtatggc	gccgactagc	tcaacggcca	cgaaccagcc	8760
aaccaccagc	gcaacctaaa	cggcgtaaac	gttgacgcca	tctctctctc	gccccgtctc	8820
gaagcttccg	caccgctcgc	tggctcgtgc	cgggcgcgcg	tcgtgctgga	ctctttccgt	8880
ggcggcttcc	gcgaaattgc	gtggtggaga	ggagagacgg	aaccgtcacg	gcactggatt	8940
ccttccccac	cggcttggc	cggccccctc	tcgcctccat	aaataggcac	cccgtcctcg	9000
cctctctctc	ccacctcacc	tctctcttcc	ccgtgaaccg	tgaacacaac	ccgacccaga	9060
tccccctctg	cgagcttctg	cgatccctcc	tccgcgtcaa	ggtaacggagc	ttctctctcc	9120
ccttctctct	tagatcggcg	tgttatgttg	tttccgtggt	tgcttggttg	gatgaatcga	9180
atgattctta	gggccttagga	ggctggttag	atctgttgcg	ttctgtttcg	tagatggatt	9240
ttggtgtaag	atcaggtcgg	ttccgctggt	taacttgtga	tgctagtgtg	atttttggga	9300
ggatttgagt	tgtaaatctg	ggagtgtgtg	ggaggttctc	gtaggcggat	tgtagatgaa	9360
gtcgcgccga	cgatttgcgt	ggcttgttgg	gtagctaggg	ttagatctgc	tcggattttt	9420
cattgttact	tattgagaga	taatgtagct	aacctttact	tgttcatcta	tgtatctcgt	9480
attcgtatct	atctggttctg	atgggtgctag	atagatgcgc	ctgatttgtc	cgatcgaatt	9540
gggtagcacc	cgcggcttgt	ttggtagtgt	tctgattgat	ttgtcgtctc	agatctgagt	9600
ggaataatat	tacatctcaa	catgttacta	gaaacttggg	ttatagctcc	ggatttaccat	9660
gtttattctt	atgtaagggt	ttaaatgaaa	gatttatgct	actgctgctc	gttgatcctt	9720
tagcatccac	ctgaggaaca	tgcatgcacc	tgttacttct	tttgatatat	gcttagatag	9780
ttgttagtat	atactgctgt	tgttcgatga	tccttcagga	tgaacatgca	tgatcatgtt	9840
acttgttttt	atatgcttct	gctgttctgt	gattcttttag	tactacctac	ctgatcatct	9900
tgcatgtttc	ctgcttggtta	gagattaatt	gattaggctt	acctgtgtgc	ctgggtgattc	9960
ttccttgca	gtgggtaccc	ggaccgatgg	caacagcaac	atcagcttct	ctgttttcaa	10020
ctgtttcttc	atcttactcc	aaagctagct	ccataccaca	ttcaagactc	caatctgtga	10080
aattcaactc	agtccttagc	ttcaccggtc	tcaaatcaac	ctctctcacc	tccggatctg	10140
attcctcttc	cttagccaag	actctacgcg	gttccgtaac	gaaagcacia	acatctgaca	10200
agaagcctta	cggattcaaa	atcaacgcta	tgcaacgggc	gctgactcct	ctcaccaaca	10260

-continued

agtatcgctt	tatcgacgtg	cagccgctga	caggcgctct	cggtgcagag	attacaggcg	10320
tggatctgcg	ggagcctctc	gatgacagca	cttggaatga	gatcctggac	gcctttcaca	10380
cctaccaagt	gatctacttt	cggggtcaag	ctatcactaa	cgagcagcac	atcgcgttct	10440
cccgcgggtt	cggccctgtg	gacccgggtg	cgatcttaaa	gagtatcgag	ggctatccag	10500
agggtgcagat	gatacggcgc	gaggcgaacg	agagcagccg	gttcacgga	gatgactggc	10560
acaccgattc	caccttcctg	gacgctccgc	ctgcgcgcgt	ggtgatgaga	gctatcgaag	10620
tgccggagta	tggaggtgac	acaggcttcc	tctccatgta	cagtgcctgg	gagacactct	10680
cgcctacgat	gcaagctacc	atcgaaggct	taaacgtggt	ccactcggcg	acgaaggctt	10740
tgggtcatt	gtaccaggcg	actaattggc	gcttctcgaa	caccagcggtg	aaagtgatgg	10800
acgtggacgc	cggagataga	gagactgtgc	acccactcgt	cgtgacgcac	cctgttacgg	10860
gaagcgcgcg	actctactgc	aaccagggtg	actgccagaa	gatccaggga	atgacggacg	10920
cggagtcgaa	gtccctgttg	caattccttt	acgagcacgc	caccaagtgc	gacttcacct	10980
gccgggtccg	gtggaagaag	gaccaagtcc	tgggtgtgga	caacctgtgt	accatgcacc	11040
gcgcgctccc	ggactacgct	gggaaattca	gatacctgac	ccgcaccacc	gtggcgggag	11100
acaagccgtc	gcgttgacgg	tccgttaatt	aatcgagtgt	gaagggaagtg	aactgagtct	11160
caagaataaa	aactgcaagt	atggcgctcc	attccatgga	agacctgcac	gcatcctata	11220
tatgtgcttt	tatgttcaag	ttagagaact	atgtgtgttg	tctttaagtt	ggagtactct	11280
aagctattcg	atctggagat	tatgctgttt	ttaataaaaag	tttgaagtgt	atctgtcctg	11340
tttaattagt	ttcaagaaag	ccaagcgtgt	ttgggttcta	ggatgaacaa	agtcgtccaa	11400
gatgcaagaa	gttctagaga	ggtagctaca	tttctggaaa	ttgttgcatc	ttcctttctt	11460
caatatatat	atattgtcct	atttcataat	gtcattctag	ccagctatct	atattctata	11520
tatccatttc	cgttgccata	ctttattcac	tatgttgcac	cgagtgcgct	gcattgctct	11580
gtactacatg	ttacttgata	tgtgtgtgtc	ataaacacac	acattaataa	tgatcagatt	11640
gtgaaaaata	cgcgtccgat	cctacctgtc	acttcatcaa	aaggacagta	gaaaagggaag	11700
gtggcaccta	caaatgccat	cattgcgata	aaggaaaggc	tatcattcaa	gatgcctctg	11760
cgcacagtgg	tcccaaagat	ggacccccac	ccacgaggag	catcgtggaa	aaagaagacg	11820
ttccaaccac	gtcttcaaag	caagtggatt	gatgtgatac	ttccactgac	gtaaggggatg	11880
acgcacaatc	ccactatcct	tcgcaagacc	cttcctctat	ataaggaagt	tcatttcatt	11940
tggagaggac	acgtgaaat	caccagtctc	tctctacaag	atcggggatc	tctagcccta	12000
gaaccatctt	ccacacactc	aagccacact	attggagaac	acacaggggac	aacacaccat	12060
aagatccaag	ggaggcctcc	gccgcgcgcg	gtaaccaccc	cgcctctctc	ctctttcttt	12120
ctccgttttt	ttttcgtctc	cgtgtctgat	ctttggcctt	ggtagtttgg	gtgggcgaga	12180
ggcggtctcg	tgcgcgccca	gatcggtgcg	cgggaggggc	gggatctcgc	ggctggggct	12240
ctcgccggcg	tggatccggc	cgggatctcg	cggggaatgg	ggctctcgga	tgtagatctg	12300
cgatccgcgc	ttgttggggg	agatgatggg	gggtttaaaa	tttccgccgt	gctaaacaag	12360
atcaggaaga	ggggaaaagg	gcactatggg	ttatatTTTT	atatatttct	gctgcttcgt	12420
caggcttaga	tgtgctagat	ctttctttct	tctttttgtg	ggtagaatTT	gaatccctca	12480
gcattgttca	tcggtagttt	ttcttttcat	gatttgtgac	aaatgcagcc	tcgtgcggag	12540

-continued

cttttttgta	ggtagaagtg	atcaaccatg	gcgcaagtta	gcagaatctg	caatggtgtg	12600
cagaacccat	ctcttatctc	caatctctcg	aaatccagtc	aacgcaaata	tcccttatcg	12660
gtttctctga	agacgcagca	gcateccacga	gcttatccga	tttcgctcgc	gtggggattg	12720
aagaagagtg	ggatgacgtt	aattggctct	gagcttcgtc	ctcttaaggt	catgtcttct	12780
gtttccacgg	cgtgcatgct	tcattggagct	tcattctaggc	cagctactgc	caggaagtct	12840
agcgggctca	gtggcacggt	gcgcatccct	ggcgataaaa	gtatttcaca	caggagcttc	12900
atgttcggag	gacttgctag	tggagagacg	agaatcactg	gtttgcttga	gggcgaagat	12960
gttatcaaca	ccggtaaagg	gatgcaagca	atgggtgccca	gaatccgaaa	agaggcgcat	13020
acgtggatca	tcgacggtgt	tggtaacgga	ggattgctcg	ctcccgaagc	gccacttgac	13080
tttgggaacg	cagctacggg	gtgccgtctt	actatgggac	tggtaggcgt	gtatgacttt	13140
gactctacct	tcattcggtga	gcgagacctc	actaagagac	caatgggacg	agtgtgtaat	13200
ccccgagggg	agatgggtgt	ccagggtgaaa	tctgaggatg	gtgatcgtct	tccggttact	13260
ctgcgaggcc	ccaagacccc	cacgccaata	acgtacaggg	ttccgatggc	gtcagcacag	13320
gtcaagttag	cggtaactct	ggcgggcctc	aacacacctg	gaatcacaa	cgtgattgaa	13380
cccatcatga	ctagagacca	cacggagaag	atgttgacgg	gtttcggcgc	taatctaacg	13440
gtcgaaacgg	acgccgacgg	cgtgaggaca	atccgcttgg	agggcagagg	taaaactgact	13500
ggccaagtca	tcgatgtgcc	tggagatccc	tcgtccacag	cgtttccctc	cgtagctgcg	13560
ttgctcgtcc	ctggatctga	tgtgacgac	ctgaatgtcc	tcattgaatc	aactagaacc	13620
ggcctcatcc	tcacattgca	ggagatgggt	gctgacatcg	aggttatcaa	tcctaggttg	13680
gcagggtggg	aggatgtggc	cgatctgcgc	gtgcgttcta	gtacactcaa	aggcgtgacc	13740
gtccctgagg	atcgcgctcc	atccatgac	gacgagtacc	ccattctcgc	cgttgctgct	13800
gcgtttgccc	aggggcgcaac	tgtaatgaac	ggccttgagg	agttgagggg	taaggagagt	13860
gacaggctgt	cgcgggtggc	gaatggcctg	aagctaaacg	gcgtggactg	cgcggaaggt	13920
gaaaacgtcc	ttgtagtccg	tgttcgccc	gacgggaagg	gggtggggaa	tgttcgggga	13980
gctgctgtgg	cgacgcacct	tgatcataga	atcgccatgt	catttctggg	gatgggactt	14040
gtctccgaga	atccgggtgac	cgttgacgat	gctaccatga	tcgccacctc	ctttcctgag	14100
ttcatggacc	tcattggcagg	cttggggggc	aagatcgagc	tgtctgatac	taaggccgct	14160
tgacacgtga	gacgacgacc	gtgcatgggt	atgaatcaca	gacgacgaac	acgcattgct	14220
catcgccgtc	ctctttgtat	cacacagcat	caccttcttc	gctctactac	tccccgagc	14280
aatcaccgac	caataacacc	aacctcaaac	ctcccccgct	gccgcgcctc	tcaccgtcct	14340
ccccctcac	catagaactg	caaatgtccg	cctgcagggt	ggggcccatc	gtggccagtt	14400
atccttagct	atccgtgtca	gaatcatctt	atcatcgagt	cgagtcgtta	tcgtgtccag	14460
tggctctctc	gagtcgagaa	gccctctatc	catccatcca	gtgttaggtg	ttcttcgtcc	14520
gtgatgttac	catgaattga	gttcgctttg	gttatgggtg	ttgaactgct	tgttgctatc	14580
tatcggaatg	aaatgaaata	gaagacaagg	agaaaaaaaa	gagttcgaaa	gttttgttcg	14640
cataccatat	atttccttcc	ggtgcgcgct	gtttattcct	cgctcagcag	caagattgtt	14700
tgatcgatat	tcgacgaagc	aattacacaa	taaatatatt	gctacactgg	tacttcaaac	14760
tacactgggt	gtcggtgatt	ttcaatagca	tgaaccttaa	ttgaacatct	gtgtagctta	14820

-continued

```

catctccttc gaaagctgca atgcttgaga acttggaag aaattcttgt gatggcagaa 14880
gctatttact gtccttcgct gcatttacag tccatacaga cacagcattt ccattttgca 14940
caagatagag aacaacaatc agccttttag gtcaatccca agtgtgcatg acgtcccgcg 15000
atcgcccgca caacaaacgc accggtgcac aaactatagc aagtgtgggc tatttttata 15060
tagcaagtga ggtcagataa aataataaat aaaatcatta aattttatat agcaagtgtg 15120
gtctattttt gcttgtaaac tacagagaat atatatatgt taaatcggtg gaacaattgg 15180
ctattttaaac taaaagattt aaaaaaaca aagataaaaa atgtttttga actgcggtgc 15240
tcgagagcag ccagcatgct tccgacctgc atgagcagcg atgctcctgg tgaccgcacg 15300
aacggaacct ggcccgatgg atgattttgc atcaacagcg cctgtatgca acgatcatcc 15360
aaccaaacat acttcgcttt gaggatccac gcctacaacg tccgcgctcg tccatgcaac 15420
caaacacacg gaatgtgttc catgtgtccc aattgaaaga agtgtattcg gttaacaacg 15480
gaagtcataa ccgagccaga tatagagata gaaccagatc tatcatacca agaacacccc 15540
tccaagattc tagactgcaa ggaaagatcc actcgtgcga agacgaccaa gatgtataag 15600
atccaatgga gcaaccatac ggaagaagag gctacgtggg agactgagga ttatctatgc 15660
aaatactacc ccgattgtct acctaaggaa gtcagtacgt aaccatgccc cagccccctg 15720
ccctccgatt ccaaatatag aaaagatact cttaatgaaa actgaattaa gaaatgaaat 15780
taagcgaaga aggacttctt tctgaagttg caaagaggat ggcatctgaa gagcagattt 15840
ttctcgcgaa ccttaaaaag ctaaccaata ctatgcaagc aagctataac cacctcctca 15900
ctcctgggtg atctcgactc gaatctcggg gcgagattct ttaaggggg gagagctgta 15960
acaccccagg tgttacttag ggtttccccc ttagtacctc catttgtgac ctattatcac 16020
atgtgggttg gtttaagaaa aaggcaccac acttgagggg ccaagccc 16068

```

```

<210> SEQ ID NO 11
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide
      PCR Primer

```

```

<400> SEQUENCE: 11

```

```

ttgctatata aaaatagacc acacttgct 29

```

```

<210> SEQ ID NO 12
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
      Primer

```

```

<400> SEQUENCE: 12

```

```

catgacgtcc cgcgatc 17

```

```

<210> SEQ ID NO 13
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
      Probe

```

-continued

<400> SEQUENCE: 13

cacaacaaac gcaccg

16

1. A recombinant DNA molecule comprising a sequence selected from the group consisting of SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:8, SEQ ID NO:7, SEQ ID NO:6, SEQ ID NO:5, SEQ ID NO:4, SEQ ID NO:3, SEQ ID NO:2, and SEQ ID NO:1.

2. The recombinant DNA molecule of claim 1, wherein
 - a) the recombinant DNA molecule is derived from a plant, seed, or cell comprising maize event MON87429, a representative sample of seed comprising the event having been deposited as ATCC Accession No. PTA-124635; or
 - b) the recombinant DNA molecule is in a plant, cell, seed, or plant part comprising maize event MON87429, a representative sample of seed comprising the event having been deposited as ATCC PTA-124635.

3. (canceled)

4. The recombinant DNA molecule of claim 1, wherein the recombinant DNA molecule is an amplicon diagnostic for the presence of maize event MON87429.

5. A DNA construct comprising four expression cassettes, wherein

- a) the first expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Erianthus ravennae*, (II) a phosphinothricin N-acetyltransferase coding sequence, and (III) a fructose-bisphosphate aldolase 3' UTR from *Setaria italica*;
- b) the second expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Coix lacryma-jobi*, (II) an albino and pale green 6 chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (III) a dicamba monooxygenase coding sequence, and (IV) a metallothionein-like protein 3' UTR from *Oryza sativa*;
- c) the third expression cassette comprises in operable linkage (I) a ubiquitin promoter, leader, and intron from *Arundo donax*, (II) a malate dehydrogenase chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (III) a FT_T protein coding sequence, and (IV) a no apical meristem protein 3' UTR from *Oryza sativa*; and
- d) the fourth expression cassette comprises in operable linkage (I) a CaMV 35S promoter and leader, (II) a chlorophyll a/b-binding protein leader from *Triticum aestivum*, (III) an actin 1 intron from *Oryza sativa*, (IV) a ShkG chloroplast transit peptide coding sequence from *Arabidopsis thaliana*, (V) a glyphosate tolerant 5-enolpyruvylshikimate-3-phosphate synthase coding sequence from *Agrobacterium* sp strain CP4, (VI) a male tissue specific siRNA target from *Zea mays*, and (VII) a glycine-rich RNA binding protein 3'UTR from *Oryza sativa*.

6. A DNA molecule having a sufficient length of contiguous nucleotides of the recombinant DNA molecule of claim 1 to function as a DNA probe specific for SEQ ID NO:10 in a sample of DNA derived from a maize plant, maize seed, or maize cell.

7. The DNA molecule of claim 6, wherein the DNA probe comprises SEQ ID NO:13.

8. A pair of DNA molecules comprising a first DNA molecule and a second DNA molecule, wherein the first and second DNA molecules each comprise a fragment of SEQ ID NO:10 and function as DNA primers when used together in an amplification reaction with DNA containing maize event MON87429 to produce the amplicon of claim 4.

9. The pair of DNA molecules of claim 8, wherein

- a) at least one DNA primer comprises a fragment of a sequence selected from the group consisting of SEQ ID NO:7 and SEQ ID NO:8; or
- b) the first DNA molecule comprises SEQ ID NO:11 and the second DNA molecule comprises SEQ ID NO:12.

10. (canceled)

11. A method of detecting the presence of maize event MON87429 in a sample of DNA derived from a maize plant, maize seed, or maize cell, the method comprising:

- a) contacting the sample with the DNA probe of claim 6;
- b) subjecting the sample and the DNA probe to stringent hybridization conditions; and
- c) detecting hybridization of the DNA probe to a DNA molecule in the sample,

wherein the hybridization of the DNA probe to the DNA molecule indicates the presence of maize event MON87429 in the sample of DNA.

12. A method of detecting the presence of maize event MON87429 in a sample of DNA derived from a maize plant, maize seed, or maize cell, the method comprising:

- a) contacting the sample with the pair of DNA molecules of claim 8;
- b) performing an amplification reaction sufficient to produce a DNA amplicon comprising a sequence selected from the group consisting of SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:8, SEQ ID NO:7, SEQ ID NO:6, SEQ ID NO:5, SEQ ID NO:4, SEQ ID NO:3, SEQ ID NO:2, and SEQ ID NO:1; and
- c) detecting the presence of the DNA amplicon,

wherein the presence of the DNA amplicon indicates the presence of maize event MON87429 in the sample.

13. A method of detecting the presence of maize event MON87429 in a sample derived from a maize plant, maize seed, or maize cell, the method comprising:

- a) contacting the sample with at least one antibody specific for at least one protein encoded by maize event MON87429; and
- b) detecting binding of the antibody to the protein in the sample,

wherein the binding of the antibody to the protein indicates the presence of maize event MON87429 in the sample.

14. The method of claim 13, further comprising at least a second antibody specific for a second protein encoded by maize event MON87429.

15. A kit for detecting the presence of maize event MON87429 comprising a DNA probe specific for SEQ ID NO:10, a pair of DNA primers that produces the amplicon

of claim 4, or at least one antibody specific for at least one protein encoded by maize event MON87429.

16. A plant, seed, cell, plant part, or commodity product comprising the DNA molecule of claim 1.

17. The plant, seed, cell, or plant part of claim 16, wherein the plant, seed, cell, or plant part comprises tolerance to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof.

18. The plant, seed, cell, or plant part of claim 17, wherein the plant, seed, cell, or plant part is tolerant to quizalofop, haloxyfop, dicamba, 2,4-D, glufosinate, and glyphosate.

19. A method for controlling weeds in a crop-growing area comprising:

- a) planting the plant of claim 16 in the crop-growing area; and
- b) applying to the crop-growing area, or any portion thereof, an effective amount of at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, to control the weeds in the area without injuring the maize.

20. The method of claim 19, wherein applying the effective amount of at least one herbicide comprises applying at least two or more herbicides selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) over a growing season.

21. The method of claim 19, wherein applying the effective amount of at least one herbicide comprises applying an herbicide selected from the group consisting of quizalofop, haloxyfop, dicamba, 2,4-D, glufosinate, and glyphosate, or any combination thereof.

22. The method of claim 21, wherein the effective amount of dicamba is about 0.5 lb ae/acre to about 2 lb ae/acre of dicamba over a growing season, wherein the effective amount of glufosinate is about 0.4 lb ai/acre to about 1.59 lb ai/acre over a growing season, wherein the effective amount of 2,4-D is about 0.75 lb ae/acre to 1.0 lb ae/acre over a growing season, wherein the effective amount of quizalofop is about 0.034 lb ai/acre to about 0.083 lb ai/acre over a growing season, and wherein the effective amount of haloxyfop is about 0.018 ai/acre to about 0.07 lb ai/acre over a growing season.

23-26. (canceled)

27. A method for controlling a volunteer maize plant comprising the recombinant DNA molecule of claim 1 in an area comprising applying an herbicidally effective amount of at least one cyclohexanedione (DIM) herbicide, where the herbicide application prevents growth of said maize plant.

28. The method of claim 27, wherein applying the herbicidally effective amount of at least one cyclohexanedione (DIM) herbicide comprises applying a cyclohexanedione (DIM) herbicide selected from the group consisting of clethodim, sethoxydim, and tralkoxydim.

29. A method of producing a plant that comprises tolerance to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, the method comprising:

- a) breeding a plant comprising maize event MON87429 with itself or a second plant to produce seed; and
- b) identifying progeny seed that comprise maize event MON87429.

30. The method of claim 29, wherein identifying progeny seed that comprise maize event MON87429 comprises:

- a) growing the progeny seed to produce progeny plants;
- b) treating the progeny plants with an effective amount of at least one herbicide selected from the group consisting of quizalofop, haloxyfop, dicamba, 2,4-D, glufosinate, and glyphosate, or any combination thereof; and
- c) selecting a progeny plant that is tolerant to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof.

31. The method of claim 29, wherein identifying progeny seed that comprise maize event MON87429 comprises detecting the presence of maize event MON87429 in a sample derived from the progeny seed or wherein identifying progeny seed that comprise maize event MON87429 comprises detecting the presence of at least one protein encoded by maize event MON87429 in a sample derived from the progeny seed.

32. (canceled)

33. A method of producing hybrid seed comprising:

- a) growing the plant of claim 16;
- b) applying an effective amount of glyphosate prior to or during the development of the male reproductive tissue of the plant thereby inducing male-sterility in the plant;
- c) fertilizing the plant with pollen from a second plant; and
- d) harvesting hybrid seed from the plant.

34. The method of claim 33, wherein applying the effective amount of glyphosate comprises application of glyphosate prior to or during the development at an effective amount of about 0.5 lb ae/acre to about 2.5 lb ae/acre.

35. The method of claim 33, wherein applying the effective amount of glyphosate comprises applying the glyphosate at a developmental stage selected from the group consisting of the V4, V5, V6, V7, V8, V9, V10, V11, V12, V13, and V14 stage of maize plant development.

36. A hybrid seed produced by the method of claim 33, wherein the hybrid seed comprises SEQ ID NO:10.

37. A method of determining zygosity of a plant comprising the recombinant DNA molecule of claim 1 comprising:

- a) contacting a sample comprising DNA derived from the plant with a primer set capable of producing a first amplicon diagnostic for the presence of said recombinant DNA molecule and a second amplicon diagnostic for the wild-type maize genomic DNA not comprising said recombinant DNA molecule;
- b) performing a nucleic acid amplification reaction; and

- c) detecting the first amplicon and the second amplicon, wherein the presence of both amplicons indicates the sample is heterozygous for said recombinant DNA molecule and the presence of only the first amplicon indicates the sample is homozygous for said recombinant DNA molecule.

38. The method of claim 37, wherein the primer set comprises SEQ ID NO:11 and SEQ ID NO:12.

39. A method of improving tolerance in a maize plant comprising tolerance to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, comprising:

- a) obtaining the DNA construct of claim 5;
- b) inserting the DNA construct into the genome of a maize cell;

- c) regenerating the maize cell into a maize plant; and
- d) selecting a maize plant comprising the DNA construct.

40. The method of claim 39, wherein the selecting comprises treating the maize plant with an effective amount of at least one herbicide selected from the group consisting of quizalofop, haloxyfop, dicamba, 2,4-D, glufosinate, and glyphosate.

41. A maize plant, maize seed, or maize cell comprising tolerance to at least one herbicide selected from the group consisting of inhibitors of acetyl CoA carboxylase (ACCase) in the aryloxyphenoxy propionate (FOP) group, synthetic auxins, inhibitors of glutamine synthetase, and inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), or any combination thereof, and obtainable by the method of claim 39, wherein the maize plant, maize seed, or maize cell comprises the DNA construct.

* * * * *